

JUL 29 1994

Mr. H. R. McLane
Howard Johnson's Enterprises, Inc.
% H.R. McLane, Inc.
7210 Red Road
Miami, Florida 33143-5321

Dear Mr. McLane:

Subject: Revised Labeling
Dursban Insecticide 0.7 Plus 16-3-5 Fertilizer
EPA Registration No. 32802-19
Your Submission Dated April 4, 1994

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable, subject to the comments listed below.

1. In the Statement of Practical Treatment for "If Inhaled," change: "Get medical attention immediately" to "Get medical attention." Immediate medical attention is not appropriate for an inhalation hazard classified as Toxicity Category III/IV.

2. Under the Directions for Use to control european crane fly larvae, the referenced grub control rate should be 50 lbs/7,575 sq. ft. (not 7,750 sq.ft.).

3. Correct the spelling of "Guarantee" in the fertilizer statement just below the Ingredients Statement. Additionally, change "contaminated" to "contaminate" in the last sentence of the Precautionary Statements.

A stamped copy of the label is enclosed for your records.

Sincerely,

DHE

Dennis H. Edwards, Jr.
Product Manager 19
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosure

292

DURSBAN® INSECTICIDE 0.7 Plus 16-3-5 Fertilizer

A Custom Lawn and Turf "Insect Control - Plus Fertilizer"

Chinch Bugs, Sod Webworms, Ants, Armyworms, Brown Dog Ticks, Crickets, Cutworms, Earwigs, Fleas, Sowbugs and White Grubs as listed which infest Home Lawns and other Ornamental & Recreational Turf areas

ACTIVE INGREDIENT:
CHLORPYRIFOS: [O,O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate] - 0.7%
INERT INGREDIENTS: - - - - - 99.3%
Total 100.0%

DURSBAN® is a Reg. Trademark of The DowElanco

Fertilizer Analysis and Gurantee found on Separate Tag.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
IF IN EYES: Flush eyes with plenty of water. Get medical attention.
IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Wash clothing before reuse.
IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated. Get medical attention immediately.
NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

See Side/Back Panel For Additional Precautionary Statements

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Do not take internally. Do not breathe dust. Avoid contact with eyes, skin or clothing. Wash contaminated clothing before reuse. Wash thoroughly after handling and before eating or smoking. When watering-in of the insecticide is directed keep children and pets off treated areas until the grass is dry. Do not taminated feed or foodstuffs.

ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters. This product is toxic to fish, birds, and wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Use only as directed.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store product in its original container in a cool, dry, locked place out of reach of children.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, out, if allowed by State and local authorities, by burning if burned stay out of smoke

(D 13) 19WTR PM5 11-22-93 vers EPA crtd 03-31-94.

SURFACE FEEDING INSECTS

For control of Chinch Bugs, Sod Webworms(Lawn Moth), Ants, Armyworms, Brown Dog Ticks, Crickets, Cutworms, Earwigs, and Sowbugs apply 50 pounds uniformly to cover 15,150 sq. ft. of turf (equivalent to 3.3 lbs. of product per 1,000 sq.ft.). Refer to Spreader Settings. Treat in the spring or when insects first appear troublesome and repeat as needed for control. The watering-in of the insecticide is not required for control of surface insects.

ROOT FEEDING INSECTS

For control of BILLBUG larvae and the "White Grub" larvae of JAPANESE BEETLE, EUROPEAN CHAFER, SOUTHERN CHAFER and HYPERODES WEEVIL apply 50 pounds to treat 7,575 sq.ft. of lawn(equivalent to 6.6 lbs. of product per 1,000 sq.ft.). Apply granules uniformly over the area and follow with a thorough watering to wash the insecticide into the soil. For best results treat when grubs are young and actively feeding near the soil surface which is usually in late July and early August. For Hyperodes (Turfgrass) Weevils apply in Mid-April and again in Mid-May at the same rate. Consult your Agricultural Extension Service for the best time to treat in your area. Keep children and pets off teated area until grass is dry

EUROPEAN CRANE FLY LARVAE

(Leather jackets)

For specific control of the Crane Fly Larvae feeding in western states turfgrass, apply 50 lbs. to cover 10,225 sq.ft. The larvae will also be controlled simultaneously when an application rate for Grubs (50 lbs/ 7,750 sq.ft.) has been made. Consult your Agricultural Extension Service for the best time to treat in your area. Keep children and pets off treated area until granules have been washed into soil and the grass is dry.

SPREADER INSTRUCTIONS

The following spreader settings are for use with new equipment. Always calibrate your spreader before applying product. Actual application rates will depend on the spreader's condition with regard to age and use, your walking speed and the evenness of the terrain, your spreading equipment will require adjustments. To calibrate, test apply by placing a measured amount for 500 sq.ft. into your spreader and apply to a pre-watered 500 sq.ft. area then adjust according to your spreader

SPREADER SETTINGS

| SPREADER | 3.3#/1,000 | 6.6 #/1,000 |
|--------------|------------|-------------|
| SPYKER-PRO | 3.4 to 3.6 | 3.4 |
| SCOTTS R7X | F to G | G to H |
| LESCO B'CAST | F to G | G to H |
| PRIZE CB-85 | 4.0 to 4.2 | 4.3 to 4.4 |
| " CBT-85 | 4.0 to 4.2 | 4.2 to 4.4 |
| CYCLONE PRO | 3.0 to 3.5 | 4.0 to 4.2 |

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens, or parks, or on golf courses or lawns and grounds.

EPA REG. NO. 32802-19

EPA EST. NO. 32802-WI-1; MO-1

The first letter of code indicates location of manufacturing plant.

NET WEIGHT 25 POUNDS (1 lb.; 4 lb.; 50 lb.)



HOWARD JOHNSON'S ENTERPRISES, INC.

P.O. BOX 2990 - MILWAUKEE, WI 53201

BEST AVAILABLE COPY