

BEST AVAILABLE COPY

page 1 of 2

DURSBAN® 1.14 GRANULAR

Lawn and Turf Insect Control

Kills: Sod Webworms, White Grubs, Ants, Imported Fire Ants, European Crane Fly Larvae (Leather jackets), Fleas, Brown Dog Ticks, Crickets, Chinch Bugs, Cutworms, Earwigs, Sowbugs, and Armyworms in Home Lawns and other Ornamental Turf.

ACTIVE INGREDIENT:
CHLORPYRIFOS: [O,O-diethyl O-(3,5,6 trichloro-2-pyridyl) phosphorothioate] 1.14%
INERT INGREDIENTS: 98.86%
Total 100.00%

DURSBAN® is a Reg. Trademark of The DowElanco

Directions For Use (Cont.)

SURFACE FEEDING INSECTS

SOD WEBWORMS, ANTS, FLEAS, BROWN DOG TICKS, CRICKETS, CHINCH BUGS, CUTWORMS, EARWIGS, SOWBUGS, ARMYWORMS

For control of Chinch Bugs, Sod Webworms(Lawn Moth), Ants, Fleas Armyworms, Brown Dog Ticks, Crickets, Cutworms, Earwigs, an Sowbugs apply this 50 pound bag to treat 25,000 sq.ft. of turf. Refer t Spreader Settings. Treat in the spring or when insects first appea troublesome and repeat as needed for control. Additional treatments a 30 to 60 day intervals may be needed for season long control of So Webworms. It is not necessary to wash granules into the soil for contro of surface insects.

USE AS A BAND TREATMENT AROUND THE HOME

To protect the dwelling from possible insect infestations a band o granules may be placed around the building to control Ants, Brown Do Ticks, Crickets, Earwigs, Fleas, Millipedes, Sowbugs, Springtails an Spiders. Or 5 pound of this product will treat a 5 foot wide band 100 fee long, arou the home, (4 ounces will treat a band 20 feet long). This wil act as a barrier against insects invading the home. Apply the protectiv band when insects have become troublesome or before they caus problems in the spring or early fall. Repeat as needed for control.

ROOT FEEDING INSECTS

WHITE GRUB CONTROL IN THE TURF

For root feeding insects such as Billbug Larvae, and the "White Grub" larvae of Japanese Beetle, European Chafer, Southern Chafer an Hyperodes Weevil apply this 50 lb. bag to treat 12,500 sq.ft. of lawn Referring to the Spreader Settings Chart apply the granules uniforml over the area and follow with a thorough watering to wash the insecticid into the soil where insects are feeding. For best control apply when th grubs are young and actively feeding near the soil surface in mid July an early August. For the Hyperodes (Turfgrass) Weevil apply in Mid-April and again in Mid-May at the same rate as above. Consult you Agricultural Extension Service for the best time to treat in your area. Kee children and pets off treated area until grass is dry.

EUROPEAN CRANE FLY LARVAE

(Leatherjackets)

For specific control of the Crane Fly Larvae feeding in western state turfgrass apply 10 lbs. to cover 3,335 sq.ft. The larvae will also b controlled simultaneously when an application rate for Grubs (10 lbs 2,500 sq.ft.) has been made. Consult your Agricultural Extension Servic for the best time to treat in your area. Keep children and pets off treat are until granules have been washed into soil and the grass is dry.

IMPORTED FIRE ANT

AS AN AID IN THE CONTROL AND REDUCTION OF FIRE ANT

As an aid in the control and reduction of Imported Fire Ants by individua ant mound treatment apply 4 level tablespoonfuls (2/3 ounce) o granules evenly in a circle around the outer edges (1 to 2 feet from center of mounds measuring from 3 to 12 inches in diameter. Finish th treatment then by applying 2 level tablespoonfuls (1/3 ounce) sprinkle over the center of the mound. Best results will be obtained if mound i not disturbed after treatment thus allowing ants to come and go and t be exposed to the insecticide treated area. Apply the granules when ant have become a nuisance and repeat as needed but not more than onc every 7 days.

SPREADER SETTINGS

SPREADER	2,000 to 4,000	4,000 to 8,000
SPYKER PRO	3.9 to 4.1	5.2 to 5.4
SCOTTS R7X	H to I	K to L
LESCO B'CAST	H to I	K to L
PRIZE CB-85	4.4 to 4.6	5.5 to 5.7
" CBT-85	4.4 to 4.6	5.5 to 5.7
CYCLONE PRO	3.9 to 4.1	5.2 to 5.4

These settings are based on a "Once Over" application.
EPA EST. NO. 32802-WI-1; MO-1

The first letter of code indicates location of manufacturing plant.

NET WEIGHT 50 POUNDS

HOWARD JOHNSON'S ENTERPRISES, INC.

P.O. BOX 2990 - MILWAUKEE WI 53201

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Wash clothing before reuse.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated. Get medical attention immediately.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Do not take internally. Do not breathe dust. Avoid contact with eyes, skin or clothing. Wash contaminated clothing before reuse. Wash thoroughly after handling and before eating or smoking. When watering-in of granules is directed keep children and pets off these treated areas until material is washed into the soil and the grass is dry. Do not contaminated feed or foodstuffs.

ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters. This product is toxic to fish, birds, and wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Use only as directed.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in its original container in a cool, dry, locked place out of reach of children.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens, or parks, or on golf courses or lawns and grounds.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label. (32-20WTR.PM4EPA corrected 01-31-94)

EPA REG. NO. 32802-20



Milwaukee, WI 53201

292

ACCEPTED
APR - 4 1994
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 32802-20

BEST AVAILABLE COPY

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100