

DURSBAN* 1.14

GRANULAR

GOLF COURSE TURF INSECT CONTROL

KILLS SOD WEBWORM, WHITE GRUBS,

EUROPEAN CRANE FLY LARVAE (LEATHER JACKETS),

ANTS, ARMYWORMS, BROWN DOG TICKS, CRICKETS,

CHINCH BUGS, CUTWORMS, EARWIGS,

AND SOWBUGS

FEB 2 2000
 U.S. PATENT OFFICE
 Registered under
 EPA Reg. No. 32802-20

ACTIVE INGREDIENT:

Chlorpyrifos: [O,O-Diethyl O-(3,5,6-trichloro-2-pyridinyl) phosphorothioate] 1.14%
 INERT INGREDIENTS: 98.86%
 TOTAL: 100.00%

Reg. No. 32802-20
 EPA Est. No. 32802-WI-1 (A), 32802-MO-1 (B)
 First letter of bag code shows establishment where manufactured.
 Manufactured by:

HOWARD JOHNSON'S ENTERPRISES, INC.
 700 W. Virginia Street, Milwaukee, WI 53204-1548

*DURSBAN and the DURSBAN logo are trademarks of DowElanco
 NET WEIGHT: 25 lb (or, 30, 36, 40, or 50lb)

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on golf course turf.

The maximum application rate is 1 lb active ingredient per acre.

SURFACE FEEDING INSECTS: For control of Chinch Bugs, Sod Webworms (Lawn Moth), Ants, Armyworms, Brown Dog Ticks, Crickets, Cutworms, Earwigs, Fleas, and Sowbugs, apply 2 pounds per 1000 square feet of turf. Refer to Spreader Settings. Treat in the spring or when insects first appear troublesome and repeat as needed for control. Additional treatments at 30 to 60 day intervals may be needed for season long control of Sod Webworms. It is not necessary to wash granules into the soil for control of surface insects.

ROOT FEEDING INSECTS:
WHITE GRUB CONTROL IN THE TURF: For root feeding insects such as Billbug Larvae, and the "White Grub" larvae of Japanese Beetle, European Chafer, Southern Chafer and Hyperodes Weevil apply 2 pounds per 1000 square feet of turf. Referring to the Spreader Settings chart apply the granules uniformly over the area and follow with a thorough watering to wash the insecticide into the soil where insects are feeding. For best control apply when the grubs are young and actively feeding near the soil surface in mid-July and early August.

For the Hyperodes (Turfgrass) Weevil apply in mid-April and again in mid-May at the same rate as above (2 pounds per 1000 square feet). Refer to the spreader setting chart. Consult your Agricultural Extension Service for the best time to treat in your area. When watering-in is directed keep public (especially children and pets) off treated areas until grass is dry.

EUROPEAN CRANE FLY LARVAE (LEATHER JACKETS): For specific control of the Crane Fly Larvae feeding in western states turfgrass apply 2 pounds per 1000 square feet. The larvae will also be controlled simultaneously when an application rate for Grubs (2 pounds per 1000 square feet) has been made. Consult your Agricultural Extension Service for the best time to treat in your area. When watering-in is directed keep public (especially children and pets) off treated areas until grass is dry.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

KEEP OUT OF REACH OF CHILDREN
CAUTION:

See (back or side) panel for additional precautionary statements and first aid information.

PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION:

May be harmful if swallowed. Do not take internally. Do not breathe dust. Avoid contact with eyes, skin, or clothing. Wash contaminated clothing before reuse. Wash thoroughly after handling and before eating or smoking. When watering-in is directed keep public (especially children and pets) off treated areas until grass is dry. Do not contaminate feed or foodstuff.

FIRST AID

IF SWALLOWED: Call a physician or poison control center immediately. Induce vomiting by giving person 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF INHALED: Remove person to fresh air. Apply artificial respiration if indicated. Get medical attention immediately.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Wash clothing before reuse.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists. Note to Physician: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine, by injection only, is an antidote.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to birds, fish, and aquatic invertebrates. Birds feeding in treated areas may be killed. Clean up spilled product to reduce exposure to wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment or disposing of equipment washwaters.

SPREADER SETTINGS to apply at the 2 lbs pounds per 1000 square feet

	2.0 lbs
Spyker 222 rotary	3.75
Lesco	10(E)
Cyclone Pro	3.4
Scotts R8A	G

(This chart may contain additional spreader brands/models, spreader settings, and instructions for spreader use and calibration, as appropriate for the product)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store in original container in a cool, dry, place out of reach of children.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.