

COMFILE B:32OFTNLP.LAB #3  
EPA FILE SYMBOL: 32802-



Reg 4 32802-23

PM16

PAGE 1 OF 4

15 POUNDS TREATS UP  
10,000 SQ.FT. FOR SOD WEBWORMS  
5,000 SQ.FT. FOR WHITE GRUBS

STOP GRUB PLUS  
WITH  
20 - 3 - 5 FERTILIZER

**ACCEPTED**  
NOV 1 1985  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 32802-23

(R)  
CONTAINS 1.5% OFTANOL  
INSECTICIDE

INSECTICIDE PLUS 20-3-5 LAWN FOOD

FOR SOD WEBWORMS, CHINCH BUGS, BILLBUG LARVAE, WHITE GRUB LARVAE,  
HYPERODES WEEVIL LARVAE AND IMPORTED FIRE ANTS

ACTIVE INGREDIENT:

1-Methylethyl 2-[[ethoxy[(1-methylethyl)-  
amino] phosphinothioyl] oxy] benzoate\* . . . 1.5%  
INERT INGREDIENTS: . . . . . 98.5%  
Total 100.0%

\*U,S Patent No. 3,621,082 Canadian Patent No. 825,198  
(R) Oftanol is a Reg. TM of the Parent Company of Farb- und Fabricken Bayer  
GmbH, Leverkusen

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce  
vomiting by giving victim 1 or 2 glasses of water and touching back of  
throat with finger. Do not induce vomiting or give anything by mouth to an  
unconscious or convulsing person.  
IF IN EYES: Flush eyes with plenty of water. Get medical attention.  
IF ON SKIN: Wash affected are with soap and warm water. Get medical attention

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 32802

EPA EST. NO. 32802-WI-1  
MO-1

NET WEIGHT 15 POUNDS

Reg # 32802-23  
PM 16

COMPILE B:32OFTNLP.LA2 #3  
EPA FILE SYMBOL: 32802

GRUB STOP PLUS (1.5% OFTANOL +) PAGE 2 OF 4  
20-3-5 FERTILIZER

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Wash contaminated clothing with soap and hot water before re-use. Keep children and pets off treated areas until insecticide has been washed to soil level and grass is dry.

**ENVIRONMENTAL HAZARDS**

Do not contaminate water by cleaning of equipment or disposal of wastes. This pesticide is toxic to fish and wildlife. Do not apply directly to water or wetlands. Keep out of lakes, streams and ponds. Apply this product only as directed.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. **STORAGE.** Store product in original container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not re-use empty bag. Wrap and put in trash.

**GUARANTEED FERTILIZER ANALYSIS**

Total Nitrogen (N) ..... 20%  
1.00% Ammoniacal Nitrogen  
14.00% Urea Nitrogen  
5.00% Sulfur Coated Slow Release Nitrogen  
Available Phosphoric Acid (P2O5). 3%  
Soluble Potash (K2O). . . . . 5%

Derived from: Urea, Sulfur-Coated Urea, Ammoniated Phosphate and Muriate of Potash. Calcium Carbonate equivalent 856 lbs. per ton.

When used at the rate of 15 lbs. per 5,000 Sq.Ft. this product supplies the lawn with 0.6 lbs. (N) nitrogen per 1,000 Sq.Ft. Water in after application for most effective insect control and fertilization. When applied to 10,000 Sq.Ft. of lawn supplies 0.3 lbs. (N) Nitrogen per 1,000 Sq.Ft.

WILL CONTROL THE FOLLOWING INSECTS AS LISTED.

SURFACE FEEDING -	IMPORTED FIRE ANTS	**ROOT FEEDING-
	CHINCHBUGS	- -List of White Grubs-
	SOD WEBWORM LARVAE	CHAFERS; PHYLLOPHAGA spp;
ROOT FEEDING -	BILLBUG LARVAE	BLACK TURF GRASS ATAENIUS;
	HYPERODES WEEVIL LARVAE	GREEN JUNE BEETLE; JAPANESE
	WHITE GRUB LARVAE**	BEETLE; ORIENTAL BEETLE;
		ASIATIC GARDEN BETTLE.

For best results, follow the timing and watering instructions below for the specific insects you need to control.

Ref # 32802-23  
PM 14

-----SURFACE INSECT CONTROL AND FERTILIZING THE LAWN-----

----- 15 POUNDS TREATS AND FEEDS UP TO 10,000 SQ.FT. OF LAWN-----

For preventive insect control apply 15 lbs. to cover \*10,000 Sq.Ft. If pests are present and damaging turf apply at rate of 15 lbs. for 5,000 Sq.Ft. - -  
FOR CHINCH BUGS AND SOD WEBWORM LARVAE: Apply when insect damage is first observed and immediately apply enough water to wet the soil 1/4 to 1/2 inches deep without causing run-off. Reapply as necessary for pest reinfestations. - - Restrictions - Do not apply more than 6 lbs./1,000 Sq.Ft. /year or an equivalent amount or 1.45 ounces of Active Ingredient /1,000 Sq. Ft. regardless of formulation used. Minimum interval between applications must be 30 days. If worms are present repeat for second generation Sod Weboworm larvae in 30 to 45 days. \*See foot note on 15 lbs/10,000 sq.ft. rate.

-----SOIL FEEDING INSECT CONTROL AND FERTILIZING THE LAWN-----

----- 15 POUNDS TREATS AND FEEDS 5,000 SQ.FT. OF LAWN-----

BILLBUG LARVAE: Apply when adult activity is observed in the spring, usually late March to Mid-May.  
HYPERODES WEEVIL LARVAE: Apply in the spring between full bloom periods of Forsythia and Flowering Dogwood, usually April 15 to May 15.  
WHITE GRUB LARVAE: (Such as Japanese Beetle, Black Turf Grass Ataenius, Chafers, Phyllophaga spp., Green June Beetle, Oriental Beetle and Asiatic Garden Beetle. PREVENTIVE APPLICATIONS: Apply in the spring (April thru Mid-May CURATIVE APPLICATION: Apply in July or early August when visible damage or high Grub populations have been observed.  
IMPORTANT RESTRICTIONS - For Billbugs, Hyperodes Weevil and White Grub Larvae do not apply more than 6 lbs. of this product /1,000 Sq.Ft. per year or an equivalent amount of 1.45 Oz. of Active Ingredient per 1,000 Sq.Ft. regardless of formulation used. Minimum interval between applications must be 30 days. IMPORTANT...IMMEDIATELY AFTER SOIL INSECT TREATMENT APPLY ENOUGH WATER TO WET THE SOIL 1 TO 1 1/2 INCHES DEEP WITHOUT CAUSING RUN-OFF.

IMPORTED FIRE ANT: Area Fire Ant Control- Apply 3 lbs./1,000 Sq.Ft. of turf uniformly over the area to be treated for residual control. Water immediately after application using 1/2 to 1 inch of water. If established Fire Ant Mounds are present use the mound drench treatment prior to area application. If a mound drench treatment is made only one area treatment may be made.

-----Established Fire Ant Mounds - DRENCH TREATMENTS-----

WET CONDITIONS-Distribute 1/2 ounce (1 1/2 tbsp) per 9 Sq.Ft. & immediately sprinkle gently with 3 gallons of water.

DRY CONDITIONS-Treat as noted for wet conditions, however, to insure penetration apply an additional 3 gallons of water to the treated mound area after treatment. Make only one mound drench treatment per season.

APPLICATION EQUIPMENT AND TECHNIQUES

For best results apply uniformly with a spreader designed to apply granular herbicides and insecticides to turf. Both drop type and rotary spreaders may be used. Avoid using spreaders that apply in narrow concentrated bands. Calibrate equipment prior to use according to manufacturers directions. Check frequently to be sure equipment is working properly. Follow directions. Apply when the wind is calm to prevent misses. Follow watering instructions.

-----SUGGESTED SPREADER SETTINGS FOR APPLYING 15 LBS/ 5,000 SQ.FT.-----

SPREADER	SETTING	Settings may vary with the age and condition of your spreader. Ask your dealer for settings to use on other brands of spreaders.
Cyclone	4	
Ortho (Rotary)	4	
Ortho (Drop)	6 1/2	
Scotts PF.1	6 1/2	

-----\*Consult Spreader Manual for 1/2 rate of 15 lbs./10,000 Sq.Ft.-----

Ref # 32802-25  
PM 16

STOP GRUB PLUS (Cont.)

GENERAL APPLICATION RESTRICTIONS

For effective insect control and as a general practice or as indicated above make sure to "water-in" the insecticide after each application. Do not mow the treated area before irrigation has been completed or before insecticide has been watered in (by rain). Do not collect clippings after first mowing after application. Do not allow product to come in continuous contact with bare skin. Do not graze treated areas or use clippings from treated areas for feed or forage. Keep children and pets off treated areas until insecticide has been washed in and the grass is dry. Do not apply more than twice per year.


NOTICE: Buyer and user assumes all risk and liability of use, and/or handling of this material not in accordance with the terms of this label.

-----  
ALTERNATE NOTICE  
-----

NOTICE: The material herein conforms to its chemical description and is reasonably fit for the purposes stated on this label when used in accordance with the directions under normal conditions. ALL OTHER WARRANTIES AND ALL IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE WHICH EXCEED THE FOREGOING REPRESENTATION ARE HEREBY DISCLAIMED BY SELLER. Buyer assumes all risks for use contrary to label instructions or for use under abnormal conditions. Sellers liability is limited to the purchase price.

-----  
{ STOP GRUB PLUS - A COMPLETE INSECTICIDE-FERTILIZER PRODUCT

KILLS WHITE GRUB LARVAE, SOD WEBWORMS, CHINCH BUGS, AND OTHER INSECTS AS LISTED ..... SUPPLIES YOUR LAWN WITH A BALANCED FERTILIZER FEEDING.



15 POUNDS TREATS UP  
10,000 SQ.FT. FOR SOD WEBWORMS  
5,000 SQ.FT. FOR WHITE GRUBS

STOP GRUB PLUS  
WITH  
20 - 3 - 5 FERTILIZER

ACCEPTED

NOV 1 1985

(R)

CONTAINS 1.5% OFTANOL  
INSECTICIDE

22802-13

INSECTICIDE PLUS 20-3-5 LAWN FOOD

FOR SOD WEBWORMS, CHINCH BUGS, BILLBUG LARVAE, WHITE GRUB LARVAE,  
HYPERODES WEEVIL LARVAE AND IMPORTED FIRE ANTS

ACTIVE INGREDIENT:

1-Methylethyl 2-[[ethoxy((1-methylethyl)-	
amino] phosphinothioyl] oxy] benzoate*	1.5%
INERT INGREDIENTS:	98.5%
Total	100.0%

\*U,S Patent No. 3,621,082 Canadian Patent No. 825,198  
(R) Oftanol is a Reg. TM of the Parent Company of Farbenfabriken Bayer  
GmbH, Leverkusen

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention.

IF ON SKIN: Wash affected are with scap and warm water. Get medical attention

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 32802

EPA EST. NO. 32802-WI-1  
MO-1

NET WEIGHT \_\_15\_\_ POUNDS

**BEST AVAILABLE COPY**

2054

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Wash contaminated clothing with soap and hot water before re-use. Keep children and pets off treated areas until insecticide has been washed to soil level and grass is dry.

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. This pesticide is toxic to fish and wildlife. Do not apply directly to water or wetlands. Keep out of lakes, streams and ponds. Apply this product only as directed.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE. Store product in original container in a cool, dry, locked place out of reach of children. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. HOUSEHOLD: Do not re-use empty bag. Wrap and put in trash.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) ..... 20%  
1.00% Ammoniacal Nitrogen  
14.00% Urea Nitrogen  
5.00% Sulfur Coated Slow Release Nitrogen  
Available Phosphoric Acid (P2O5). 3%  
Soluble Potash (K2O). . . . . 5%

BEST AVAILABLE COPY

Derived from: Urea, Sulfur-Coated Urea, Ammoniated Phosphate and Muriate of Potash. Calcium Carbonate equivalent 856 lbs. per ton.

When used at the rate of 15 lbs. per 5,000 Sq.Ft. this product supplies the lawn with 0.6 lbs. (N) nitrogen per 1,000 Sq.Ft. Water in after application for most effective insect control and fertilization. When applied to 10,000 Sq.Ft. of lawn supplies 0.3 lbs. (N) Nitrogen per 1,000 Sq.Ft.

WILL CONTROL THE FOLLOWING INSECTS AS LISTED.

SURFACE FEEDING -	IMPORTED FIRE ANTS	**ROOT FEEDING- - - -
	CHINCHBUGS	- -List of White Grubs- -
ROOT FEEDING -	SOD WEBWORM LARVAE	CHAFERS; PHYLLOPHAGA Spp.;
	BILLBUG LARVAE	BLACK TURF GRASS ATHANUS;
	HYPERODES WEEVIL LARVAE	GREEN JUNE BEETLE; JAPANESE
	WHITE CRUB LARVAE**	BEETLE; ORIENTAL BEETLE;
		ASIATIC GARDEN BEETLE.

For best results, follow the timing and watering instructions below for the specific insects you need to control.

COMFILE # SCOTT'S LAG #3 GRUB STOP PLUS (1.5% OFTANOL+) PAGE 2 OF 4  
EPA FILE SYMBOL:32802- 20-3-5 FERTILIZER

-----SURFACE INSECT CONTROL AND FERTILIZING THE LAWN-----  
----- 15 POUNDS TREATS AND FEEDS UP TO 10,000 SQ.FT. OF LAWN-----  
For preventive insect control apply 15 lbs. to cover \*10,000 Sq.Ft. If pests are present and damaging turf apply at rate of 15 lbs. for 5,000 Sq.Ft. - -  
FOR CHINCH BUGS AND SOD WEBWORM LARVAE: Apply when insect damage is first observed and immediately apply enough water to wet the soil 1/4 to 1/2 inches deep without causing run-off. Reapply as necessary for pest reinfestation. - - Restrictions - Do not apply more than 6 lbs./1,000 Sq.Ft. /year or an equivalent amount of 1.45 ounces of Active Ingredient /1,000 Sq. Ft. regardless of formulation used. Minimum interval between applications must be 30 days. If worms are present repeat for second generation Sod Webworm larvae in 30 to 45 days \*See foot note on 15 lbs/10,000 sq.ft. rate.

-----SOIL FEEDING INSECT CONTROL AND FERTILIZING THE LAWN-----  
----- 15 POUNDS TREATS AND FEEDS 5,000 SQ.FT. OF LAWN-----  
BILLBUG LARVAE: Apply when adult activity is observed in the spring, usually late March to Mid-May.  
HYPERODES WEEVIL LARVAE: Apply in the spring between full bloom periods of Forsythia and Flowering Dogwood usually April 15 to May 15.  
WHITE GRUB LARVAE: (Such as Japanese Beetle, Black Turf Grass Ataenius, Chafers, Phyllophaga spp., Green June Beetle, Oriental Beetle and Asiatic Garden Beetle.PREVENTIVE APPLICATIONS: Apply in the spring (April thru Mid-May CURATIVE APPLICATION: Apply in July or early August when visible damage or high Grub populations have been observed.  
IMPORTANT RESTRICTIONS - For Billbugs, Hyperodes Weevil and White Grub Larvae do not apply more than 6 lbs. of this product /1,000 Sq.Ft. per year or an equivalent amount of 1.45 Oz. of Active Ingredient per 1,000 Sq.Ft. regardless of formulation used. Minimum interval between applications must be 30 days.  
IMPORTANT...IMMEDIATELY AFTER SOIL INSECT TREATMENT APPLY ENOUGH WATER TO WET THE SOIL 1 TO 1 1/2 INCHES DEEP WITHOUT CAUSING RUN-OFF.

IMPORTED FIRE ANT: Area Fire Ant Control- Apply 3 lbs./1,000 Sq.Ft. of turf uniformly over the area to be treated for residual control.Water immediately after application using 1/2 to 1 inch of water. If established Fire Ant Mounds are present use the mound drench treatment prior to area application. If a mound drench treatment is made only one area treatment may be made.

-----Established Fire Ant Mounds - DRENCH TREATMENTS-----  
WET CONDITIONS-Distribute 1/2 ounce (1 1/2 tbsp) per 9 Sq.Ft.& immediately sprinkle gently with 3 gallons of water.

DRY CONDITIONS-Treat as noted for wet conditions, however, to insure penetration apply an additional 3 gallons of water to the treated mound area after treatment. Make only one mound drench treatment per season.

**APPLICATION EQUIPMENT AND TECHNIQUES**

For best results apply uniformly with a spreader designed to apply granular herbicides and insecticides to turf. Both drop type and rotary spreaders may be used. Avoid using spreaders that apply in narrow concentrated bands. Calibrate equipment prior to use according to manufacturers directions. Check frequently to be sure equipment is working properly. Follow directions. Apply when the wind is calm to prevent misses. Follow watering instructions.

-----SUGGESTED SPREADER SETTINGS FOR APPLYING 15 LBS/ 5,000 SQ.FT.-----

SPREADER	SETTING	Settings may vary with the age and condition of your spreader. Ask your dealer for settings to use on other brands of spreaders.
Cyclone	4	
Ortho (Rotary)	4	
Ortho (Drop)	6 1/2	
Scotts PF.1	6 1/2	

-----\*Consult Spreader Manual for 1/2 rate of 15 lbs./10,000 Sq.Ft.-----

STOP GRUB PLUS (Cont.)

GENERAL APPLICATION RESTRICTIONS

For effective insect control and as a general practice or as indicated above make sure to "water-in" the insecticide after each application. Do not mow the treated area before irrigation has been completed or before irrigation has been watered in (by rain). Do not collect clippings after first mowing after application. Do not allow product to come in continuous contact with bare skin. Do not graze treated areas or use clippings from treated areas for feed or forage. Keep children and pets off treated areas until insecticide has been washed in and the grass is dry. Do not apply more than twice per year.

NOTICE: Buyer and user assumes all risk and liability of use, and/or handling of this material not in accordance with the terms of this label.

-----  
ALTERNATE NOTICE  
-----

NOTICE: The material herein conforms to its chemical description and is reasonably fit for the purposes stated on this label when used in accordance with the directions under normal conditions. ALL OTHER WARRANTIES AND ALL IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE WHICH EXCEED THE FOREGOING REPRESENTATION ARE HEREBY DISCLAIMED BY SELLER. Buyer assumes all risks for use contrary to label instructions or for use under abnormal conditions. Seller's liability is limited to the purchase price.

-----  
STOP GRUB PLUS - A COMPLETE INSECTICIDE-FERTILIZER PRODUCT  
-----

KILLS WHITE GRUB LARVAE, SOIL WEBWORMS, CHINCH BUGS, AND OTHER INSECTS AS LISTED ..... SUPPLIES YOUR LAWN WITH A BALANCED FERTILIZER FEEDING.