



WITH 20-3-5 FERTILIZER
CONTAINS 1.5% OFTANOL IN INSECTICIDE

FOR SOD WEDWORMS, CHINCH BUGS, BILL BUG LARVAE, WHITE GRUB LARVAE,
HYPERODES WEEVIL LARVAE AND IMBOTTED FIRE ANTS

ACTIVE INGREDIENTS

1. Methyl ethyl 2-(ethoxy-1-hydroxyethyl)-amino phosphorothioic acid benzothiazolyl ester 1.5%

INERT INGREDIENTS 98.5%

U.S. Patent No. 3,621,061; Canadian Patent No. 1,142,195
(R) Oftanol is a Reg. TM of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center in the state. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention.

IF ON SKIN: Wash affected area with soap and warm water. Get medical attention.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 32802-23 EPA EST. NO. 32802-WI (A)
MO-1 (B)

The first letter of code on bottom of tape indicates location of manufacturing plant.

NET WEIGHT 45 POUNDS

Manufactured by:
HOWARD JOHNSON'S ENTERPRISES, INC.
VIROQUA, WISCONSIN 54665
NEOSHO, MISSOURI 64850

GUARANTEED FERTILIZER ANALYSIS

TOTAL NITROGEN (N)	20%
14.00% Urea Nitrogen	
5.00% Sulfur Coated Urea Nitrogen	
Available Phosphoric Acid (P2O5)	3%
Soluble Potash (K2O)	5%

Contains 14.00% Urea Sulfure Coated Urea, Ammoniated Phosphate, 5.00% Potash, Calcium Carbonate equivalent 8% (lbs per 100 lbs)

PRECAUTIONS AND RESTRICTIONS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION Harmful if swallowed. Avoid contact with eyes. If contact occurs, wash thoroughly with soap and water after handling and before eating or smoking. Wash thoroughly with soap and hot water before reuse. Keep out of reach of children. Do not feed or allow animals to graze on treated areas until soil and grass is dry.
 Do not contaminate water by storage, handling, or disposal. Do not apply directly to water or wetlands.
 Do not apply directly to water or wetlands.

BEST AVAILABLE COPY

SURFACE FEEDING	CRICKETS	ROOT FEEDING	BEETLE LARVAE
	GRASSHOPPERS		HYPERODES WEEVIL LARVAE
	SOIL WEBWORMS		WHITE GRUB LARVAE
	CHINCHES		
	FLYING ANT		
	THREATENING		
	BEETLE		
	BEETLE		
	BEETLE		
	BEETLE		

SURFACE INSECT CONTROL - FEEDING ON THE LAWN

Apply this product to lawns infested with crickets, grasshoppers, chinch bugs, flying ants, and beetles. Apply in the evening or early morning. Do not apply if rain is expected within 24 hours. Do not apply to wet lawns. Do not apply to lawns that have been treated with other insecticides within 14 days.

SOIL FEEDING INSECT CONTROL - FEEDING IN THE LAWN

Apply this product to lawns infested with soil webworms, hyperodes weevil larvae, and white grub larvae. Apply in the evening or early morning. Do not apply if rain is expected within 24 hours. Do not apply to wet lawns. Do not apply to lawns that have been treated with other insecticides within 14 days. Do not apply to lawns that have been treated with other soil insecticides within 14 days.

SUGGESTED SPREADER SETTINGS FOR APPLICATIONS OF 15 LBS/1000 SQ FT		
SPREADER	SETTING	Settings may vary with the age and condition of your spreader. Ask your dealer for settings to use on other brands of spreaders.
Cyclone	4	
Ortho (Rotary)	4	
Ortho (Drop)	6 1/2	
Scotts PF 1	6 1/2	

* Consult Spreader Manual for 1/2 rate of 15 lbs/10,000 sq ft

GENERAL APPLICATION RESTRICTIONS

For effective insect control and as a general practice or as indicated above make sure to water in the insecticide after each application. Do not re-irrigate treated area before irrigation has been completed or before insecticide has been watered in by rain. Do not collect clippings after first mowing after application. Do not allow product to come in contact with the bare skin. Do not graze treated areas or use clippings from treated areas for feed or forage. Keep children and pets off treated area until soil has been washed in and the grass is dry. Do not apply more than twice per year.

STORAGE AND DISPOSAL
 Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Store product in original container in a cool dry place out of reach of children. **PESTICIDE DISPOSAL:** Waste or extra from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag in application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOW TO REFILL:** Do not reuse empty bag. Wrap and put in trash.