

PM 25 32802-28 Page 1 of 1

SEED SAFE - TURF CARE

TUPERSAN®—Pre-emergence Herbicide Plus Turf Food may be used on newly seeded lawns.

When used as directed—Will not harm your desirable lawn grass seeds—Yet kills annual weed grasses such as SMOOTH and HAIRY CRABGRASS—FOXTAIL WEED GRASSES—and BARNYARDGRASS (WATERGRASS).

USE ON home lawns, golf courses, parks, roadsides, athletic fields, all ornamental and recreational turfgrass.

This Herbicide is not for use on Bermudagrass or Bentgrass.

TUPERSAN®—Registered Trademark of the E.I. DuPont de Nemours Company
U.S. Patent #3,309,192

ACTIVE INGREDIENT:

Siduron (1-(2-Methylcyclohexyl-3-Phenylurea))
3.71%

INERT INGREDIENTS: 96.29%
Total 100.00%

Howard Johnson's Enterprises, Inc.
Milwaukee, WI 53201

EPA REG. NO. 32802-28

EPA EST. NO. 32802-WI-1 (A)

32802-MO-1 (B)

The first letter of code indicates location of manufacturing plant.

NET WEIGHT 50 POUNDS

Fertilizer guarantee found on separate tag.

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENTS OF PRACTICAL TREATMENTS

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Get medical attention if irritation persists.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. May irritate eyes, nose, throat and skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Do not contaminate feed or foodstuffs. Do not use on turf to be used for feed or food purposes.

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. This product is toxic to fish. Do not apply to any body of water. Do not apply where run-off is likely to occur. Do not apply when weather favors drift from treated areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Store product in original container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **HOUSEHOLD:** Do not re-use empty bag. Wrap and put in trash.

DIRECTIONS FOR USE (Cont.)

WHERE TO USE

For use on BLUEGRASS—FESCUE—REDTOP—PERENNIAL RYEGRASS—ZOYSIA—SMOOTH BROME—ORCHARDGRASS AND CERTAIN BENTGRASS STRAINS as PENNCROSS, SEASIDE, HIGHLAND, ASTORIA, NIMISILA, C1, C7 and C-19. Do not use on other Bentgrass varieties since injury may occur—Do not use on Bermudagrass—This product will not control Poa Annua.

This product will not control established weeds that are already growing.

BEST AVAILABLE COPY

Not for use on turf grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

HOW MUCH AND WHEN TO USE

USE TO COVER 20,000 SQ. FT. FOR NEW SPRING SEEDINGS—Apply 50 lbs. to treat and feed 20,000 sq. ft. of lawn where bare ground or very thin turf is seeded in the spring of the year. Apply as the final operation after seeding in the spring and 2 weeks prior to weed seed germination. Follow good cultural practices of keeping area moist to help grass seed germinate and also to activate the pre-emergent herbicide and to get maximum benefit from the lawn food.

USE TO COVER 10,000 SQ. FT. FOR NEW "FALL" SEEDING OR SEEDING TO ESTABLISHED LAWNS IN THE SPRING—Apply 50 lbs. to treat and feed 10,000 sq. ft. of turf. Apply in spring prior to the expected emergence of the annual weed grasses. This will vary from early February in the south to early March in the north. Follow the application with a thorough watering (1/2 inch) to activate the herbicide.

HOW TO APPLY

To insure best weed control and proper fertilization the lawn should be free of leaves, grass clippings or thatch which might interfere with the pre-emergence action of the herbicide. Complete all cultural practices such as aeration before application. Apply with a properly calibrated Fertilizer Spreader. If it does not rain within 3 days after application, water the area thoroughly to get desired results.

MAKE SURE YOUR SPREADER IS OPERATING CORRECTLY

Spreader settings are for use with new equipment. Since proper application is based on the spreader's condition with regard to age and use, your walking speed and the evenness of the terrain, your spreader may require adjustment.

TO CHECK YOUR SPREADER

Using a Cyclone B-1 Spreader to apply 50 lbs. per 10,000 sq. ft. place 2.5 lbs. of this product in your spreader using No. 4 1/2 setting and apply to 500 sq. ft. With a 20 inch wide spreader this will equal 3 spreader widths wide (3 passes) by 100 feet long. If you run out of granules before you cover the pre-measured area, close the spreader setting by 1/2 number. If you have granules left over, open (increase) the spreader setting by 1/2 number. For further help refer to your spreader guide.

APPLICATION CHART

For optimum pre-emergence weed control apply at the appropriate time.

APPROPRIATE TIMES TO APPLY THIS PRE-EMERGENT HERBICIDE

JAN. FEB. MAR. APR. MAY JUNE JULY AUG. SEPT. OCT. NOV. DEC.

Northern Lawns

Southern Lawns

1. For Crabgrass, Foxtail, & other Grassy Weeds "NORTHERN STATES"
2. For Crabgrass, Foxtail, & other Grassy Weeds "SOUTHERN STATES"

RESEEDING THE TREATED AREAS

This product may be applied to the soil at the same time as sowing grass seed. When used as directed on listed turfgrasses this product will not affect the germination of those seeding grasses.

WEEDS CONTROLLED WITH THIS PRODUCT

Annual Weed Grasses such as Crabgrass (smooth and hairy), Foxtail (green and yellow), Barnyardgrass (watergrass).

SPREADER SETTINGS

SPREADER	50 LBS. PER 20,000 SQ. FT.	50 LBS. PER 10,000 SQ. FT.
SPYKER PRO	3.6-3.8	4.4-4.6
SCOTTS R7X	G-H	J-K
LESCO B'CAST	G-H	J-K
PRIZE CB-85	4.0-4.2	4.8-5.0
CBT-85	4.0-4.2	4.8-5.0
CYCLONE PRO	3.6-3.8	4.4-4.6
LELY B'CAST	2.3-2.5	3.4-3.6

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

AUG 10 1994

RECEIVED
SEP 21 1994
EPA Reg. No. 32802-28