

PM 11 24/04 208

10f2

AUG 21 1995

5488273  $\frac{301}{37}$

488277  $\frac{300}{18}$

Howard Johnson's Enterprises, Inc.  
c/o H.R. McLane, Inc.  
7210 Red Rd., Suite 206  
Miami, FL 33143-5321

Attention: Mr. H.R. McLane

Subject: Trichlorfon 6.2G  
EPA Registration No. 32802-29  
Your submissions of May 19 and 24, 1995

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. Place the "Storage and Disposal" text after the use directions and before the "NOTICE".

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

Robert A. Forrest  
Product Manager (14)  
Insecticide-Rodenticide Branch  
Registration Division (H7504C)

- Enclosures:
1. Stamped label
  2. A-29 Enclosure

Peacock WP#6:A:32802-29:305-5407,-6600:8/21/95

10f2

2 of 2

# TRICHLORFON 6.2G TURF GRANULES

## FOR CONTROL OF LAWN AND TURF INSECTS SUCH AS WHITE GRUB SPECIES, SOD WEBWORMS AND CUTWORMS.

For use in home lawns, parks, apartment & condominium complexes, golf courses, athletic fields and other ornamental & recreation turf. Does Not Destroy Beneficial Insect Populations When Used As Directed. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

AUG 21 1995

ACTIVE INGREDIENT:  
Dimethyl (2,2,2-trichloro-1-hydroxy-ethyl) phosphonate- - 6.2%  
INERT INGREDIENTS:- - 93.8%  
Total 100.0%

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 32802-29

### KEEP OUT OF REACH OF CHILDREN CAUTION

#### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention. **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and warm water. Wash clothing before reuse. **IF INHALED:** Remove victim to fresh air. Apply artificial respiration if indicated. Get medical attention immediately. **NOTE TO PHYSICIAN:** Prolonged exposure will cause cholinesterase depression. Atropine Sulfate is total. 2-PAM is also antidotal and may be administered in conjunction with atropine.

#### PRECAUTIONARY STATEMENTS

##### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** May be harmful if swallowed. Do not take internally. Do not breathe dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling and before eating or smoking. Keep children and pets off treated areas until material has been washed to soil surface and grass is dry. Do not contaminate feed or foodstuffs. Do not use treated areas or clippings from treated areas for feed or forage.

##### ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. This product is toxic to fish, birds and wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Use only as directed on this label. Do not use on crops used for food or forage.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in its original container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not reuse empty container. Wrap and put in trash.

**NOTICE:** Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label. (32-29am3.pm5 \ 05-24-95)

## SURFACE FEEDING INSECTS

**SOD WEBWORMS (Lawn Moth)** Sod Webworms are capable of destroying large areas of turf. They live through the winter as a larvae and change to a moth in May. Eggs are deposited very soon throughout the lawn and new worms hatch in May. By killing this first brood it is possible to reduce the larvae population of later broods and therefore reduce turf damage.

#### HOW TO APPLY

Turf should be free of troublesome thatch which encourages insects. Rake or detatch the area and remove the debris to expose insect hiding places. Water the lawn and cut to proper height before application

#### HOW MUCH TO APPLY

For control of Sod Webworms (Lawn Moth Larvae) and Cutworms apply at the rate of 2 lbs. per 1,000 sq.ft. of turf. Refer to chart for respective spreader settings

## WHITE GRUB LARVAE CONTROL IN THE TURF

**FOR ROOT FEEDING "White Grub" insect larvae of JAPANESE BEETLE, EUROPEAN CHAFER, AND SOUTHERN CHAFER apply 3 lbs. to treat 1,000 sq.ft. of turf.** Make sure your spreader is properly calibrated to apply this product. Irrigate the turf after treatment as soon as possible to wash the insecticide into the root zone where the insects are feeding. Apply when the White Grub Larvae are young and actively feeding near the soil surface in mid July and early August. Consult your Agricultural Extension Service for the best time to treat for "Grubs" in your area. A second treatment at the same rate may be needed for mature large sized grubs.

#### WHEN TO APPLY

Treat in the spring (Mid-May to June) or when your lawn program calls for a Surface Insect Control Application. Repeat as the program indicates or as needed for control. Sod webworms may have 3 to 4 generation of larvae in 1 year which may require additional treatments at 30 to 40 day intervals during the summer months. Lightly sprinkle the area to move the insecticide down into the insect feeding zone. Do not allow children or pets on treated areas until material has been sprinkled in and the grass is dry.

#### HOW TO APPLY - IMPORTANT

Do not attempt to control White Grubs in turf that has over 1/2 to 3/4 inch thatch build up since heavy thatch will prevent the insecticide from penetrating down to the area where the insects are feeding. Therefore, troublesome thatch must be removed before treating for White Grubs. If thatch is not removed before treatment, poor Grub control will result.

#### SPREADER SETTING

The following spreader setting are for use with new equipment. Your used spreader may need calibrating. Since proper application is based on the spreader's condition with regard to age and use, your walking speed and the evenness of the terrain, your spreading equipment may require adjustments. If you have reason to believe that your spreader is not calibrated correctly test apply by placing the amount for 500 sq.ft. into you spreader and apply to your pre-measured 500 sq.ft. (5 ft. wide by 100 feet long, equals 3 passes with a 20 inch wide spreader, or by other suitable calibrating methods.

#### SPREADER SETTING CHART

SPREADER	2 # / 1,000 sq.ft.	3 # / 1,000 sq.ft.
Ames Drop	11.5	14.0
Ames Rotary	11.5	13.0
Cyclone B-1	3.7	4.1
Earthway Drop	11.5	14.0
Earthway Rotary	1.5	13.0
Ortho Drop	5.0	7.0
Ortho Rotary	4.0	6.5
Prec Drop	9.5	11.0
Prec Rotary	5.5	8.0
Scotts Drop	5.25	8.0

EPA REG. NO. 32802-29

NET WEIGHT 15 POUNDS

EPA EST NO. 32802-WI-1

# HOWARD JOHNSON'S ENTERPRISES, INC.

P.O. BOX 2990 - MILWAUKEE, WI 53201

2