

ALL SEASON

TRIAMINE WEED AND FEED 25-8-3

Accept - treatments in Spring or Fall - not worth arguing

This product is a premium grade herbicide and turf feed combination which controls a wide spectrum of broadleaf weeds while providing both fast and long lasting action.

ACTIVE INGREDIENTS:

- *Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid..... 0.313%
- **Dimethylamine Salt of 2-(2-Methyl-4-chlorophenoxy) propionic Acid..... 0.314%
- ***Dimethylamine Salt of 2-(2,4-Dichlorophenoxy) propionic Acid..... 0.310%

Inert Ingredients..... 99.065%

Total..... 100.00%

EQUIVALENT TO:

- *0.26% of 2,4-Dichlorophenoxyacetic Acid
- **0.26% of 2-(2-Methyl-4-Chlorophenoxy) propionic Acid
- ***0.26% of 2-(2,4-Dichlorophenoxy) propionic Acid

EPA REG. NO. 32802-

EPA EST. NO. 32802-WI-1(A)
32802-MO-1(B)

The first letter of code on bottom tape indicates the location of the manufacturing plant.

KEEP OUT OF REACH OF CHILDREN - 12 pt. type

WARNING- 18 pt. type

IF SWALLOWED: Get Medical Attention.

IF ON SKIN: Wash Skin With Soap And Water.

IF IN EYES: Flush Eyes With Water For 15 Minutes And Get Medical Attention.

ACCEPTED

FEB 24 1986


Under the Federal Insecticide, Fungicide, and Herbicide Act as amended, for the pesticide registered under EPA Reg. No. 32802-31

NET WEIGHT 20

Covers 5000 Square Feet

MANUFACTURED BY:

Howard Johnson's Enterprises, Inc.
Viroqua, Wisc. 54665
Neosho, Mo. 64850



PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING - 18 pt type

Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash clothing after handling. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear goggles or safety glasses.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply when conditions favor drift from target areas. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GENERAL DIRECTIONS

TRIAMINE WEED AND FEED 25-3-3 is a stable mixture of three oxenoxy herbicides. It is for use on lawns, turf (not pastures), aprons, tees, and rounds.

DIRECTIONS

Read entire label before using this product. This product may be used on bluegrass, fescue, rye, bermuda, bahia, and zoysia lawns. DO NOT USE on dichondra, carpetgrass, and St. Augustine grass. DO NOT USE this product where desirable clovers are present or allow this product to "wind drift" on flowers, vegetables, ornamentals, shrubs, trees, or other desirable plants.

WHEN TO APPLY: Best results are obtained when weeds are growing rapidly in spring or early fall. IMPORTANT: Apply when grass is moist (after a rain, sprinkling, or dew). Do not water lawn after material is applied. Water will wash off weed killing material. For most effective broadleaf weed killing action, avoid unnecessary disturbance of treated areas for 48 hours after application. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded lawn until after mowings.

HOW TO APPLY: This 20 pound bag should be used on 3,000 square feet. Apply evenly with a lawn spreader either as one pass application or a two pass application, criss-crossing the area; half the rate each time. Lawns that are heavily over-run by weeds and that are in poor condition, may require more than one application of this product. A suggested second application should be made in approximately 30 days.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed, raise setting for higher speed.

SPREADER SETTINGS:	MAKE	SETTING
	Cyclone	5 1/4 - 5 3/4
	Scotts 35	5 1/2
	Scotts	C-1 J-1
	Central	7 - 8
	A-Marc	7 - 8
	Montgomery wards	6 3/4

This product is designed for control of broadleaf weeds in lawns such as:

Black Medic	Lambsquarter	Speedwell
Buckhorn	Mallow	Source
Burdock	Morning glory	white clover
Chickory	Pigweed	wild Carrot
Chickweed	Plantain	wild Onion
Clover	Poison Ivy	wild Lettuce
Dock	Poison Oak	Yarrow
Dandelion	Ruslane	Other Broadleaf
English Daisy	radweed	weeds
Ground Ivy	Red Clover	
Heal-All	red sorrel	
Knockweed	Stitchwort	

STORAGE AND DISPOSAL

STORAGE: Always store product in original container and in locked area.

CONTAINER DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash. Do not reuse bag.

Notice: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other products, and conditions of application equipment will vary, seller makes no claims other than those stated on this label.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N).....	25%
1.8% Ammonical Nitrogen	
16.5% Urea Nitrogen	
6.7% Sulfur Coated urea	
Available Phosphoric Acid (P2O5).....	3%
Soluble Potash (K2O).....	3%

Plant Food Derived From: Urea, Sulfur Coated Urea, Ammoniated Phosphate, Muriate of Potash. Potential Acidity Equivalent to 1100 Lbs. Calcium Carbonate Per Ton.