

OCT 4 1991

Mr. Michael Archer
 Director of Technical Services
 Howard Johnson's Enterprises, Inc.
 P.O. Box 2990
 Milwaukee WI 53201-2400

BEST AVAILABLE COPY

Dear Mr. Archer:

Subject: All Season Sprayable Weed & Feed
 EPA Registration No. 32802-48
 All Season Trifluralin Weed & Feed
 EPA Registration No. 32802-51
 Howard Johnson's Weed & Feed
 EPA Registration No. 32802-3
 Re: Amended Labeling
 Your Submission Dated September 30, 1991

The labeling referred to above submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the label changes under the "Environmental Hazard" Section of the label. Replace the statement "Do not apply directly to water or wetland" with the statement "Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark."

Also add the groundwater contamination statement, under the Environmental Hazard Section, "Most cases of groundwater contamination involving phenoxy herbicide such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies."

CONCURRENCES

SYMBOL	WH/RSK						
SURNAME	MAT						
DATE	10/2/91						

Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/ loading equipment on an impervious pad to contain spills will help prevent groundwater contamination. A stamped copy is enclosed for your records.

Sincerely

Jeanne J. Miller
Product Manager
EPA Registration Division
Registration Division

ENCLOSURE

BEST AVAILABLE COPY

2 of 6

ALL SEASON TRIAMINE WEED & FEED 25-3-3

Premium Herbicide kills Dandelions, Clover
Ground Ivy & Over 40 Other Weeds.

ACTIVE INGREDIENTS:

1. Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid.....	0.3132
2. Dimethylamine Salt of 2-(2-Methyl-4-chlorophenoxy)propionic Acid.....	0.3142
3. Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid.....	0.3102
INERT INGREDIENTS.....	99.0632
TOTAL.....	100.0002

ASSAYED BY ISOMER SPECIFIC ASSAY PROC METHOD NO. D01-5:

- 1. 0.26% of 2,4-Dichlorophenoxyacetic Acid
- 2. 0.26% of 2-(2-Methyl-4-chlorophenoxy)propionic Acid
- 3. 0.26% of 2-(2,4-Dichlorophenoxy)propionic Acid

Keep Out of Reach of Children
CAUTION

STATEMENTS OF PRACTICAL TREATMENT

- IF SWALLOWED: Get medical attention
- IF ON SKIN: Wash skin with soap and water
- IF IN EYES: Flush eyes with water for 15 minutes and get medical attention

See other panels for additional precautionary statements.

COVERS 10,000 sq. ft.
NET WT. 40 LBS.

EPA REG. NO. 32802-31 EPA EST. NO. 32802 WI-1 (A)
32802 MO-1 (B)

The first letter of code on bag seal indicates the location of the manufacturing plant.

Manufactured by: Howard Johnson's Enterprises, Inc.
Milwaukee, WI 53201

OCT 04 1991

BEST AVAILABLE COPY

32802-31

4-3-91

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	23.2
1.9% Ammoniacal Nitrogen	
23.1% Urea Nitrogen*	
Available Phosphoric Acid (P ₂ O ₅)	3.2
Soluble Potash (K ₂ O)	3.2
Sulfur (S)	2.2
Iron (Fe)	0.2
* 6.7% Slowly available nitrogen derived from sulfur coated urea.	

Plant food derived from: Urea, ammoniated phosphate, sulfur coated urea and muriate of potash. Sulfur derived from sulfur coated urea. Iron derived from ferric oxide.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution: Avoid inhalation of dust and contact with eyes, skin or clothing. Wear long pants, long sleeved shirt and shoes when applying this product. Wash thoroughly after handling. Do not allow persons or pets on treated areas until dusts have settled.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides, or seeds.

PRODUCT DISPOSAL: To dispose of bags which contain unused product, securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

BEST AVAILABLE COPY

5 of 6

DIRECTIONS FOR USE, CONT.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

WEED & FEED will control the most prevalent broadleaf weeds occurring in home lawns while providing a blend of nutrients that will make your lawn lush and green. This product may be used on bluegrass, fescue, ryegrass, Bermuda, St. Augustine, centipede grass, bahia, and zoysia lawns. Do not use on dichondria, carpetgrass or bentgrass lawns. Keep away from inside the drip line of shrubs and ornamentals and do not apply or allow to drift onto vegetable or flower gardens.

WHEN TO APPLY:

- (1) Best results occur when weeds are growing rapidly.
- (2) Apply when grass is moist, after a rain, irrigation, heavy dew, or other watering methods.
- (3) Avoid mowing or watering the lawn for 48 hours after application. This gives the broadleaf killer time to be absorbed into the weed leaves.
- (4) Do not reseed lawns for four weeks after application. Do not apply to newly seeded lawns until after three mowings.
- (5) In cases of severe weed infestation, hard to kill weeds (such as chickweed, or less than optimal weed killing conditions occur, weeds are not actively growing, cool weather occurs, or product was washed off the weeds), it may be necessary to retreat the lawn three to four weeks after the initial application of WEED & FEED. This application could take the place of the normal summer TURF FOOD application.
- (6) When used during hot (over 85 degrees F) and/or dry conditions apply WEED & FEED at one half the recommended rate to lessen the chance of burning your lawn.
- (7) Under conditions of adequate moisture, high temperature, and high humidity, temporary discoloration may occur if over application occurs on St. Augustine grass.

When used according to instructions, WEED & FEED will control the following weeds in your lawn.

BEST AVAILABLE COPY

Black Medic	English Daisy	Plantain	Speedwell
Buckhorn	Ground Ivy	Poison Ivy	Spurge
Burdock	Heal All	Poison Oak	White Clover
Chickory	Knotted	Purslane	Wild Carrot
Chickweed	Lambsquarter	Ragweed	Wild Onion
Clover	Mallow	Red Clover	Wild Lettuce
Dock	Morning Glory	Red Sorrel	Wild Garlic
Dandelion	Pioneed	Stitchwort	Yarrow

And other similar Broadleaf weeds

6/2/6

HOW TO APPLY:

Apply WEED & FEED with a lawn spreader either in one pass or by applying at half-rate in two passes. If spread using the two pass method, run the second pass at right angles to the first.

The following rates are approximate. Check your spreader manual for additional instructions. The condition of your spreader will greatly affect its rate of application. Walk at a normal pace while applying. Check your spreader on a 200 square foot section and adjust your setting to apply one pound of WEED & FEED to this area.

Wash out your spreader after each use and allow it to dry thoroughly. Treat any exposed metal areas with oil to prevent any corrosion.

SPREADER SETTINGS:

MAKE	SETTING
Excitone	5 1/4 - 5 3/4
Scotts 35	5 1/2
Scotts	1 J
Central	7 B
K Mart	7 B
Montgomery Ward (1091, 1093, 1095)	6 3/4

Contact your local retailer for additional information regarding establishing a total lawn care program tailored to your needs.

NOTICE: Manufacturer warrants that this product conforms to all label guarantees and will perform as indicated when used as directed. Manufacturer will not honor any claims resulting from product usage not defined or recommended on the label.

9-91

BEST AVAILABLE COPY