Tom J. Schmit, Mgr. Reg. Affairs

APR 24 1985

S. - EPA Use Only

L SEASON TRIAMINE®WEED & FEED

Premium Herbicide Kills Dandelions, Clover, Ground Ivy & Over 40 Other Weeds

Kills weeds quickly & makes your lawn lush and green. Kills Ugly Dandelions, Chickweed, Clover, Ground Ivy, Ragweed, Plantain & other common broadleaf weeds. Fast-acting nutrients make your lawn greener & lusher. Use in late spring or early summer when weeds are growing. Applyto wet leaves for the best control.

ACTIVE INGREDIENTS:

Dimethylamine Salt of

2,4-Dichlorophenoxyacetic acid -0.313%

**Dimethylamine Salt of 2-(2-Methyl-4-

chlorophenoxy) propionic Acid -

***Dimethylamine Salt of 2-(2,4-Dichloro

phenaxy) propionic Acid -**INERT INGREDIENTS:**

0.310% 99.063%

0.314%

100,000% Total Equivalent to: 0.26% of 2,4-Dichlorophenoxyacetic Acid 0.26% of 2-(2-Methyl-4-Chlorophenoxy) propionic Acid 0.26% of 2-(2.4-Dichlorophenoxy) propionic Acid

EPA REG. NO. 32802-31 EPA EST. NO. 32802-WI-1(A) 32802-MO-1(B) The first letter of code on bag seal indicates the location of the manufacturing plant.

KEEP OUT OF REACH OF CHILDREN CAUTION

IF SWALLOWED: Get medical attention. IF ON SKIN: Wash skin with scap and water. IF IN EYES: Flush eyes with water for 15 minutes and get medical Attention.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) - variable % variable % Ammonical Nitrogen variable % Urea Nitrogen

Available Phosphoric Acid (P₂0_c) - var Soluble Potash (K₂0) - variable % Iron (Fe) variable % variable % variable%

Plant Food Derived From: Urea, Ammoniated Phosphate, Sulfur Coated Urea, and Muriate of Potash. Sulfur Derived From Sulfur Coated Urea. Iron Derived From Ferric Oxide.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Avoid inhalation of dust and contact with eyes, skin or clothing. When using this product, wear long pants, long-sleeved shirt, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor wind drift of product from target area since this product will injure beans, cucumbers, grapes, roses, tomatoes, and most vegetables, flowers and ornamental plants. Do not apply where run-off is likely to occur. Do not contaminate water used for irrigation or domestic purposes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals: Do not store near feed or feedstuffs. To prevent cross-contamination, do not store near other fertilizers or near herbicides, insecticides, fungicides or seeds. PRODUCT DISPOSAL: To dispose of bags which contain unused product. securely wrap bag in several layers of newspaper and discard in trash. CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

NET WEIGHT 20 (Also 18, 36; 40; or 50) lbs. Covers 5,000 Square Feet (5,000 sq.ft. for 18 & 20, 10,000 sq.ft. for 36 & 40, 12,500 sq.ft. for 50 lbs)

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

ALL SEASON TRIAMINE® WEED AND FEED will control the most prevalent broadleaf weeds occurring in home lawns while providing a blend of nutrients that will make your lawn lush and green. This product may be used on bluegrass. fescue, ryegrass, Bermuda, St. Augustine, centipede grass Bahia and zoysia, lawns. Do not use on dichondra and Carpetgrass or Bentgrass lawns. Keep away from inside the drip line of shrubs and ornamentals and do not apply to allow to drift onto vegetables or flower gardens.

WHEN TO APPLY

- Best results are obtained when weeds are growing rapidly.

 Apply when grass is moist, after a rain, irrigation, heavy dew, or other watering methods.
- Avoid moving or watering the lawn for 48 hours after application. This gives the broadleaf killer time to be absorbed into the weed leaves.
- Do not reseed lawns for 4 weeks after application. Do not apply to newly seeded lawns until after 3 mowings.

 5. In cases of severe weed infestations, hard to kill weeds such as chickweed, or
- less than optimal weed killing conditions, (e.g. weeds are not actively growing, cool weather occurs, or product was washed off the weed), it may be necessary to retreat the lawn three to four weeks after the initial application of Weed & Feed This application could take the place of the normal summer TURF FOOD application.
- The maximum rate to turf is 0.46 pounds of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.
- When used during hot (over 85 degrees F) and/or dry conditions apply WEED & FEED at one-half the recommended rate to lessen the chance of burning your
- Under conditions of adequate moisture, high temperature, and high humidity, temporary discoloration may occur if over-application occurs on St. Augustinegrass.

When used according to instructions. ALL SEASON TRIAMINE® WEED & FEED will control the following weeds in your lawn.

Black Medic **English Daisy Plantains** Ground Ivy Buckhorn Poison Ivv Burdock Heal-All Poison Oak Chickweed Knotweed Pursiane Ragweed Chicory Lambsquarter Clover Red Clover Mallow Dandelion Red Sorrel Morning-glory Dock Speedwell Piaweed

Spurge Stitchwort White Clover Wild Carrot Wild Garlic Wild Lettuce Wild Onion Yarrow (and many more broadleaf weeds)

HOW TO APPLY

Apply ALL SEASON TRIAMINE® WEED & FEED with a lawn spreader either as one pass or by applying at half-rate in two passes. If spread using the twopass method, run the second pass at right angles to the first.

The following rates are approximate Check your spreader manual for additional instructions. The condition of your spreader will greatly affect its rate of application. Walk at a normal pace while applying. Check your spreader on a 250 sq.ft. section and adjust your setting to apply one pound of WEED & FEED to this area. Wash out your spreader after each use and allow it to dry thoroughly. Treat any exposed metal areas with oil to prevent corrosion.

SPREADER SETTINGS	
Cyclone Scott 35 Scotts Central K-Mart Montgomery Wards	51/453/4 51/2 IJ 7-8 ((() 63/4

Contact your local retailer for additional information regarding establishing a total lawn care program tailored to your needs. ر د ر د د

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility withouts? products, and conditions of application equipment will vary. Seller makes population of the standard of the st claims other than those stated on this label.

(32802-31.PM5 \ Acceptd w/c 04-22-94 \ Notif Amd. [as sod] Rev. 04-05-96)



HOWARD JOHNSON'S ENTERPRISES, INC. P.O. BOX 2990 - MILWAUKEE, WI. 53201