

HI-TECH CRABGRASS CONTROL

Reg # 32802-33

With Team[®] Herbicide

AM-23



181

Team[®] is a Registered Trademark of the Dow/Elanco Products Company

Not for use on turf grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

For Control of Crabgrass and Most Other Annual Weed Grasses on Home Lawns, Golf Courses, Parks and Other Ornamental & Recreational Turf.

FOR USE ON ESTABLISHED: Bluegrass (Perennial)—Ryegrass (Perennial)—Fescue—Bahagrass—Bermudagrass—Centipede-grass—St. Augustine Grass—Zoysiagrass

ACTIVE INGREDIENTS:

N-butyl-N-ethyl-alpha, alpha, alpha, trifluoro-2,6-dinitro-p-toluidine	1.00%
TRIFLURALIN; Alpha, alpha, alpha, trifluoro-2,6-dinitro-N,N-dipropyl-o-toluidine	0.50%
INERT INGREDIENTS:	98.50%
Total	100.00%

Howard Johnson's Enterprises, Inc.
Milwaukee, WI 53203

EPA REG. NO. 32802-33

EPA EST. NO. 32802-WI-1 (A)
32802-MO-1 (B)
41723-PA-1 (S)

The first letter of code indicates location of manufacturing plant.

NET WEIGHT 50 POUNDS

**KEEP OUT OF REACH OF CHILDREN
WARNING**

STATEMENTS OF PRACTICAL TREATMENTS

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Call a physician.
IF SWALLOWED: Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid Alcohol.
IF ON SKIN: Remove contaminated clothing. Wash with plenty of soap and water. Get medical attention if irritation persists.
IF INHALED: If breathing difficulty occurs, move individual to fresh air and get medical attention.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING: Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust or contact with skin or clothing. The granular formulation may cause skin sensitization reaction in certain individuals. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and chemical resistant gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not allow children or pets on lawn until 1/2" of water has been applied to lawn and the water has dried (or, if wet conditions persist, until 24 hours have passed).

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, or marshes). Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in its original container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not re-use empty bag. Wrap and put in trash.

WHERE TO APPLY

When applied according to these Directions this product can be used safely on the following turfgrass varieties: PERENNIAL BLUEGRASS VARIETIES, PERENNIAL RYEGRASS VARIETIES, BERMUDAGRASS, BAHAGRASS, CENTIPEDEGRASS, FESCUE VARIETIES, ST. AUGUSTINE GRASS & ZOYSIAGRASS lawns and ornamental turf. Will not harm most nearby ornamental trees and shrubs.

WEEDS CONTROLLED IN ESTABLISHED TURF

This product contains a combination of Beneth and Trifluralin pre-emergent herbicides plus Lawn Fertilizer. It should be applied to established turf for the control of CRABGRASS (Smooth and Hairy), GOOSEGRASS (Silver Crab or Crowfoot), BARNYARDGRASS (Watergrass), FOXTAIL (Green and Yellow), and ANNUAL BLUEGRASS (Poa Annual) and provides a complete lawn fertilization.

WHERE, WHEN AND HOW TO APPLY THIS PRODUCT: Apply to turf prior to germination of weed seeds since established weeds will not be controlled. Apply only to healthy established turf grass lawns.

IMPORTANT PRECAUTIONS

DO NOT: 1 Apply in the spring to Bermuda Grass which has been overseeded with winter grasses (annual rye) since thinning will occur
2 Do not use on Dichondra, nor to newly sprigged areas of Bermudagrass, St. Augustine Grass or Centipede Grass, until these turfs are well established, nor to Bahagrass or Putting Greens.
3 Apply in spring to Turfgrass planted the previous fall. This product may thin established Annual Bluegrass (Poa Annual) & Fine-Leaved Fescue lawns when used at higher rates than 132 lbs. per acre (2.0 lb. A.I. per Acre). Do not overdose

WHEN AND HOW TO APPLY

Successful pre-emergence control of the annual grass weeds on this label requires proper timing of application. This product must be applied one to two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grassy weeds listed on this label, weed control results may be erratic or poor.

WATER THE LAWN AFTER APPLICATION: It is not necessary but optimum weed control performance will be obtained if treated areas are irrigated with 1/2 to 1 inch water soon after application. Any cultural practices that disturb the soil such as verticuting, should be done prior to application of this product.

**SPRING GERMINATING SUMMER ANNUAL GRASSES:
CRABGRASS, FOXTAIL, BARNYARDGRASS:**

A. Northern cool season grasses—For pre-emergence control, apply 100 to 132 lbs. per acre (equal to 1.5 to 2.00 lbs. Active Ingredient per acre) applied in the spring one to two weeks prior to the onset of conditions favorable for annual weedy grass seed germination.

B. Southern warm season turfgrasses—For pre-emergence control, apply 132 to 200 lbs. per acre (equal to 2 to 3 lbs. Active ingredient per acre), in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weedy grass seed germination.

LATE SPRING, EARLY SUMMER GERMINATING ANNUAL GRASSES & GOOSEGRASS (SILVER CRAB):

Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than the crabgrass, foxtails and other summer annual grass species. Applications of this product should, therefore, be timed accordingly.

A. Northern cool season turfgrasses—For pre-emergence control, apply 132 lbs. per acre (equal to 2 lbs. Active ingredient per acre) of this product in late spring or early summer one to two weeks prior to the expected germination period of goosegrass.

B. Southern warm season turfgrasses—For pre-emergence control, apply 132 to 200 lbs. per acre (equal to 2 to 3 lbs. Active ingredient per acre) of this product in late spring or early summer one to two weeks prior to the expected germination of goosegrass.

FULL SEASON CONTROL IN NORTHERN AND SOUTHERN TURFGRASS

In areas of the United States where pre-emergence control of both early germinating annual grasses (Crabgrass, Foxtails, and Barnyardgrass) and the later germinating Goosegrass is desired, two applications of this product at the recommended rate spaced 8 to 10 weeks apart may be made to provide full season control of these annual grass species.

POA ANNUA (ANNUAL BLUEGRASS)

In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Proper fertilization, irrigation and soil incorporated re-seeding will encourage existing desirable turfgrasses and newly planted seedling turfgrasses to fill in these thin areas previously occupied by annual bluegrass.

A. Northern cool season turfgrasses—For pre-emergence control, apply 132 lbs. per acre (equal to 2 lbs. Active ingredient per acre) of this product per acre in late summer or early fall and again in late winter or spring one to two weeks prior to expected germination of annual bluegrass.

B. Southern warm season turfgrasses—For pre-emergence control, apply 200 lbs. per acre (equal to 3 lbs. Active ingredient per acre) of this product per acre in the late summer or early fall one to two weeks prior to expected germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application at 200 lbs. per acre can be made 10 to 12 weeks after the initial treatment.

RESEEDING TREATED AREAS

1. Delay reseeding 6 weeks when applying at the rate of 100 to 132 lbs. of product per acre (equal to 1.5 to 2.0 lbs. Active ingredient per acre.)

2. Delay reseeding 12 to 16 weeks when applying at the rate of 200 lbs. of product per acre (equal to 3.0 lbs. of Active ingredient per acre)

WHEN RESEEDING—INCREASE THE RECOMMENDED AMOUNT OF SEED BY 25%.

The following spreader settings are for use with new equipment. Your spreader may need calibrating since proper application is based on the spreader's condition with regard to age and use, your walking speed and the unevenness of the terrain. Poor weed control will result from uneven, or improper application.

SPREADER SETTINGS

SPREADER	1.50 LBS A./ACRE	2.0 LBS A./ACRE	3.0 LBS A./ACRE
SPYKER PRO	3.5-3.7	3.8-3.9	4.2-4.4
SCOTTS R7X	F-G	H-I	J-K
LESCO B'CAST	F-G	H-I	J-K
PRIZE CB-85	4.0-4.2	4.0-4.2	4.0-4.2
CBT-85	4.0-4.2	4.0-4.2	4.0-4.2
CYCLONE PRO	3.5-3.7	3.8-3.9	4.2-4.4
LELY B'CAST	2.3-2.5	2.6-2.8	3.1-3.3

50 LBS TO TREAT 22,000 SQ. FT.=100 LBS./ACRE=1.5 LBS A.I./ACRE

50 LBS TO TREAT 17,000 SQ. FT.=130 LBS./ACRE=2.0 LBS A.I./ACRE

50 LBS TO TREAT 11,000 SQ. FT.=200 LBS./ACRE=3.0 LBS A.I./ACRE

NOTICE: Buyer and user assumes all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

AUG 10 1994

