

JAN 10 1988

EPA 32802-35 PUB NO 830

EPA REG. No. 32802-35

20 LB BAG COVERS up to 5,000 Sq Ft

LAWN SERVICE CRABGRASS CONTROL

Contains a 22-3-11 Complete Lawn Food with Iron & Sulfur This Premium Grade Crabgrass Preventer and Lawn Food Combination is designed to control Crabgrass and other Common Annual Grasses while fertilizing your lawn.

Nitrogen found in this product is partially derived from a Slow Release Source that, while providing a Steady Source of Greening, Eliminates the Burning often associated with Certain High Analysis Fertilizers

ACTIVE INGREDIENTS:

N-butyl-N-ethyl-alpha, alpha, alpha, trifluoro-2,6-dinitro-p-toluidine 0.56%
TRIFLURALIN, Alpha, alpha, alpha, trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine 0.29%

INERT INGREDIENTS 99.15%
Total 100.00%

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENTS OF PRACTICAL TREATMENTS

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Call a physician.
IF SWALLOWED: Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, large quantities of water. Avoid Alcohol.
IF ON SKIN: Remove contaminated clothing. Wash with plenty of soap and water. Get medical attention if irritation persists.
IF INHALED: If breathing difficulty occurs, move individual to fresh air and get medical attention.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
EPA REG. NO. 32802-35 EPA EST. NO. 32802-WI-1 MQ-1

Contact your local retailer for additional information regarding establishing your total lawn care program.
FIRST LETTER OF CODE ON BAG BOTTOM TAPE INDICATES MFG PLAN

NET WEIGHT 20 POUNDS (40 lb.)

HOWARD JOHNSON'S ENTERPRISES, INC.
MILWAUKEE, WI 53201

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust and contact with skin or clothing. The granular formulation may cause skin sensitization reactions in certain individuals. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not allow children or pets on lawn until 1/2 inch of water has been applied to the lawn and the water has dried for, if wet conditions persist, until 24 hours have passed.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, or marshes). Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in its original container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Household: Do not reuse empty bag. Securely wrap original bag in several layers of newspaper and discard in trash.

DIRECTIONS FOR USE (Cont.)

This product gives your lawn protection against the germination of Crabgrass, Foxtail, and Barnyardgrass, and at the same time provides a springtime boost of nutrients needed to grow a green, healthy lawn. May be used on Perennial Ryegrass, Perennial Bluegrass, Fescue, Bermudagrass, Bahiagrass, St. Augustine Grass, Carpetgrass and Zoysia grass lawns. For satisfactory results make sure you follow the recommendations listed below. Will not harm most nearby ornamental trees and shrubs.

IMPORTANT

To be effective, this product must be applied to lawns one to two weeks before the germination and emergence of Crabgrass, Foxtail, and Barnyardgrass seedlings. Actively growing weeds present at the time of application will not be controlled with this material.

APPLICATION INSTRUCTIONS

- Do not apply to bermudagrass overseeded to winter annual grasses or thinning will occur.
- Do not use on Dichondra, or newly sprigged areas of St. Augustine-grass, Bermudagrass or Centipede grass unless these areas are well established.
- Do not apply turfgrass seeded the previous fall.
- Water the lawn with 1/4 to 1/2 inch of water after application to allow the material to form the "barrier" on the soil surface necessary for adequate weed control.
- Do not rake, thatch, aerate or otherwise disturb this "barrier" after application.
- Use higher spreader setting for lawns that have a history of severe weed problems.
- Sometimes cool, rainy weather during the summer months will simulate "spring crabgrass germinating weather", thus causing dormant annual grass seedlings to germinate. With these situations, a retreatment with 8 to 10 weeks after initial applications, will prevent the reoccurrence of annual grasses.
- Delay reseeding of treated areas for 8 to 10 weeks after application.

NORMAL SPRING APPLICATION TIMES

Northern Lawns ... Mid March to May 1 st
Southern Lawns ... Mid January to April 1 st
Application times may vary according to local weather patterns

HOW MUCH TO APPLY

Types of Lawns	Comment	Apply 20 lbs. to Cover
Southern Lawns	Serious Weed Problem	2,500 Sq Ft
Southern Lawns	Maintenance Only	3,000 Sq Ft
Northern Lawns	Serious Weed Problem	3,000 Sq Ft
Northern Lawns	Maintenance Only	5,000 Sq Ft

SPREADER SETTINGS

The following settings are approximate. Check your spreader manual for additional instructions. The condition of your spreader will greatly affect its rate of application. Walk at a normal pace while spreading. After application, treat all metal parts of your spreader that have been exposed to fertilizer with a light coating of oil to help prevent corrosion.

Spreader	Spreader Settings for Square Feet		
	2,500 Sq Ft	3,000 Sq Ft	5,000 Sq Ft
Precision	8	6	4
PSB	8	4	4
Scots	7	4	4
Cyclone	8	4	4
Central	8	4	4

Apply Crabgrass Preventer Plus with a lawn spreader either in one pass or by applying at a half rate in two passes. Half rate settings are approximately two marks below full rate suggestions. See your spreader manual for further suggestions. If spreading using the two pass method, run the second pass at right angles to the first.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	27%
12% Ammoniacal Nitrogen	
3.7% Controlled Slow Release Nitrogen	
17.1% Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	3%
Soluble Potash (K ₂ O)	11%
2.0% Sulfur	0.7% Iron

Plant food derived from Urea, Ammoniacal Phosphate, Murate of Potash, Controlled Slow Release Nitrogen derived from Sulfur Coated Urea, Sulfur derived from Sulfur Coated Urea, Iron derived from Ferric Oxide.

NOTICE: The manufacturer warrants that this product conforms to all label guarantees, and will perform as indicated when used as directed. Manufacturer will not honor any claims resulting from product usage not defined or recommended on the label.