

PM 23

32802-35

11-3-98

10/4

"Occupational Use" Label Version

LAWN SERVICE CRABGRASS CONTROL

WITH TEAM* HERBICIDE FOR PRE-EMERGENCE
CONTROL OF CRABGRASS, FOXTAIL, GOOSEGRASS,
ANNUAL BLUEGRASS (*POA ANNUA*) AND OTHER
ANNUAL WEED GRASSES AND BROADLEAF WEEDS
IN ESTABLISHED LAWNS AND ORNAMENTAL TURF

ACTIVE INGREDIENTS:

benefin:

N-butyl-*N*-ethyl- α,α,α -trifluoro-2,6-dinitro-*p*-toluidine 0.58%

trifluralin:

α,α,α -trifluoro-2,6-dinitro-*N,N*-dipropyl-*p*-toluidine 0.29%

INERT INGREDIENTS: 99.13%

TOTAL: 100.00%

Contains 0.435 pounds active ingredient per 50 pound bag

EPA Reg. No. 32802-35

EPA Est. No. 32802-WI-1 (A), 32802-MO-1 (B)

F Letter of bag code shows establishment where manufactured.

S Bag attached to bag for fertilizer guaranteed analysis.

Manufactured by:

HOWARD JOHNSON'S ENTERPRISES, INC.

700 W. Virginia Street, Milwaukee, WI 53204-1548

*TEAM is a trademark of Dow AgroSciences LLC

KEEP OUT OF REACH OF CHILDREN

WARNING: See back panel for further cautions,
first aid information and directions for use.

NET WEIGHT:

ACCEPTED

NOV 3 1998

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 32802-35

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury.
Harmful if swallowed or absorbed through skin. Do not get in eyes or
clothing. Avoid contact with skin. Prolonged or frequently repeated
skin contact may cause allergic reactions in some persons.

PERSONAL PROTECTIVE EQUIPMENT (PPE): Applicators and
other handlers must wear goggles or face shield, long-sleeved shirt,
long pants, shoes and socks, chemical resistant gloves when
handling. Wash hands before eating, drinking, chewing gum, using
tobacco, or using the toilet. Remove contaminated clothing and
wash clothing before reuse. Follow manufacturer's instructions for
cleaning/maintaining PPE. If no such instructions for washables, use
detergent and hot water. Keep and wash PPE separately from other
laundry.

USER SAFETY RECOMMENDATIONS: Users should remove
clothing immediately if pesticide gets inside. Then wash
thoroughly and put on clean clothing. User should remove PPE
immediately after handling this product. Wash the outside of
gloves before removing. As soon as possible, wash thoroughly
and change into clean clothing.

FIRST AID

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of
water for 15 minutes. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF SWALLOWED: Do not induce vomiting. Call physician or Poison
Control Center. Give person a large quantity of milk, egg white,
gelatin solution, or if these are not available, large quantities of water.
Avoid alcohol.

Note to physician: Probable mucosal damage may contraindicate the
use of gastric lavage.

ENVIRONMENTAL HAZARDS: This pesticide is extremely toxic to
freshwater, marine and estuarine fish, and aquatic invertebrates
including shrimp and oyster. Do not apply in a manner that will
directly expose canals, lakes, streams, ponds, marshes or estuaries
to aerial drift. Do not apply directly to water, or to areas where
surface water is present or to intertidal areas below the mean high
water mark. Do not contaminate water when disposing of equipment
washwaters.

DIRECTIONS FOR USE: It is a violation of Federal law to use this
product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod,
or for commercial seed production, or for research purposes.

Entry restrictions: Do not enter or allow others to enter the treated area
until dusts have settled.

GENERAL INFORMATION: This product contains turf food plus a
preemergence herbicide for pre-emergence control of annual weed
grasses and broadleaf weeds in established turf. Apply to turf before
germination of weed seeds, since established weeds will not be
controlled.

WEEDS CONTROLLED:

Annual grasses: annual bluegrass (*Poa annua*), smooth crabgrass
(*Digitaria ischaemum*), large or hairy crabgrass (*Digitaria sanguinalis*),
goosegrass (silver crabgrass or crowfootgrass) (*Eleusine indica*),
barnyardgrass (watergrass) (*Echinochloa crus-galli*), green foxtail
(*Setaria viridis*), yellow foxtail (*Setaria glauca*).

Broadleaf weed: common chickweed (*Stellaria media*), prostrate
knotweed (*Polygonum aviculare*), common lambsquarter
(*Chenopodium album*), pigweed (*Amaranthus spp.*), common purslane
(*Portulaca oleracea*).

Weeds suppressed: Texas panicum (*Panicum texanum*) Florida pusley
(*Richardia scabra*), prostrate spurge (*Euphorbia humistrata*), spotted
spurge (*Euphorbia maculata*), creeping spurge (*Euphorbia serpens*),
creeping woodsorrel (*Oxalis corniculata*), woodsorrel (*Oxalis spp.*).

(Label for EPA Reg. No 32802-35, PAGE 1 OF 4)

92718 (Occupational use)

WHERE TO APPLY: This product may be applied to the following established turfgrass species:

Established cool season turfgrass: bentgrass fairways, Kentucky bluegrass, perennial ryegrass, fescues.

Established warm season turfgrass: bahia, bermuda, centipede, tall fescue, St. Augustine, zoysia.

Important precautions: Do not apply this product to golf course putting greens, dichondra, or newly sprigged area of bermuda, St. Augustine, centipede or zoysiagrass. Apply only to healthy, established turf. Do not apply in spring to turfgrass planted the previous fall. Will not harm most nearby ornamental trees, shrubs and flowers. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

WHEN TO APPLY: Successful preemergence weed control requires proper timing of the application. This product must be applied 1 to 2 weeks before the germination of weed seeds, since established weeds will not be controlled. All annual weed grasses do not germinate at the same time. For most effective weed control, apply at the proper time:

Northern areas:

For crabgrass, foxtail, barnyardgrass, etc., apply Mar. 1 through June 1

For goosegrass, which germinates later, apply May 15 through July 1

For annual bluegrass, apply July 1 through Oct. 1

Southern areas:

For crabgrass, foxtail, barnyardgrass, etc., apply Feb. 1 through June 1

For goosegrass, which germinates later, apply April 1 through July 1

For annual bluegrass, apply Oct. 1 through Feb. 1

APPLICATION RATES: Apply in spring applications to control spring-to-summer germinating annual weeds, or fall applications (to control fall-germinating weeds). A single application will usually provide season-long weed control. Apply this product at the rate of 4 to 8 pounds per 1000 square feet. Make a single application 1 to 2 weeks before the expected germination of weed seeds.

To control goosegrass or annual bluegrass, apply 5.25 to 8 pounds per 1000 square feet, in late spring or early summer 1 to 2 weeks prior to onset of conditions favorable to weed seed germination. Better control may be possible by using the higher rate.

Apply this product evenly over the turf, using a properly calibrated granular fertilizer spreader which will uniformly distribute the product. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur. For the most effective weed control, water or irrigate the treated areas soon after application. Cultural practices that disturb the soil, such as aerification, should be done before application.

SPLIT APPLICATIONS FOR FULL SEASON WEED CONTROL: A series of split applications is recommended when the initial application is made well in advance of weed germination, or where an extended period of weed control is required (areas with longer growing seasons). In cool season turfgrass areas and the northern half of warm season turfgrass areas, apply 4 pounds per 1000 square feet plus a second application of 4 pounds per 1000 square feet. Make first application in spring 1 to 2 weeks prior to expected time of weed germination, and make the second application 8 to 10 weeks later.

In the southern half of warm season turfgrass areas, apply 5.25 pounds per 1000 square feet plus a second application of 5.25 pounds per 1000 square feet. Make first application in spring 1 to 2 weeks prior to expected time of weed germination, and make the second application 10 to 12 weeks later.

RESEEDING: This product will inhibit germination of grass seed if seeded too soon after application. When reseeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. Delay reseeding treated areas for a minimum of:

8 to 10 weeks after last application of 4 pounds/1000 square feet

10 to 12 weeks after last application of 5.25 pounds/1000 square feet

12 to 16 weeks after last application of 8 pounds/1000 square feet

When soil testing or professional advice indicates the need for additional plant nutrients, make additional applications of fertilizer products over the growing season.

This product may stain driveways, walks, patios, pools, etc. Wash or sweep off thoroughly immediately after each use.

204

SPREADER SETTINGS to apply at the listed pounds per 1000 square feet			
	4 lbs	5.25 lbs	8 lbs
Spreader A	X	X	X
Spreader B	X	X	X
Spreader C	X	X	X
Spreader D	X	X	X

(May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in a locked storage area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GUARANTEE: This product is unconditionally guaranteed.

(Alternate wording for warranty statement:)

NOTICE: Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

(Alternate wording for warranty statement:)

WARRANTY AND DISCLAIMER: Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label. Manufacturer/seller makes no other warranty, expressed or implied, and shall not be liable for indirect or consequential damages.

(Alternate wording for warranty statement:)

WARRANTY AND DISCLAIMER: Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label. No other warranty, expressed or implied, is made concerning the merchantability or fitness for a particular purpose. Manufacturer/seller's liability shall be limited to the purchase price of the product. In no case shall manufacturer/seller be liable for consequential or incidental damages or loss.

(Alternate wording for warranty statement:)

WARRANTY AND DISCLAIMER: Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. No other warranty, expressed or implied, is made concerning the merchantability or fitness for a particular purpose. Manufacturer/seller's liability shall be limited to the purchase price of the product. In no case shall manufacturer/seller be liable for consequential or incidental damages or loss. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label. Products not accepted under these terms should be returned within 20 days in the original, unopened container for return of purchase price.

"Household Use" Label Version

LAWN SERVICE CRABGRASS CONTROL

WITH TEAM® HERBICIDE FOR
PRE-EMERGENCE CONTROL OF CRABGRASS,
FOXTAIL, GOOSEGRASS, ANNUAL BLUEGRASS AND
OTHER ANNUAL WEED GRASSES & BROADLEAF WEEDS
IN ESTABLISHED LAWNS

PREVENTS CRABGRASS & OTHER WEEDS (AS LISTED)
FROM SPROUTING IN YOUR LAWN

ACTIVE INGREDIENTS:

benefin:

N-butyl-N-ethyl- α,α,α -trifluoro-2,6-dinitro-p-toluidine 0.52%

trifluralin:

α,α,α -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine 0.29%

INERT INGREDIENTS: 99.13%

TOTAL: 100.00%

EPA Reg. No. 32802-35

EPA Est. No. 32802-WI-1 (A), 32802-MO-1 (B)

First letter of bag code shows establishment where manufactured.

Manufactured by:

HOWARD JOHNSON'S ENTERPRISES, INC.

700 W. Virginia Street, Milwaukee, WI 53204-1548

*TEAM is a trademark of Dow AgroSciences LLC

KEEP OUT OF REACH OF CHILDREN

WARNING: See back panel for further cautions.
first aid information and directions for use.

NET WEIGHT:

GUARANTEED ANALYSIS

Total Nitrogen (N) X%
Available Phosphate (P₂O₅) Y%
Soluble Potash (K₂O) Z%

(May contain additional fertilizer Guaranteed Analysis information as required by
fertilizer composition and state registration and labelling requirements.)

ACCEPTED

NOV 3 1998

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 32802-35

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury.
Harmful if swallowed or absorbed through skin. Do not get in eyes or
clothing. Avoid contact with skin. Prolonged or frequently repeated
skin contact may cause allergic reactions in some persons. Wear
goggles or face shield, long-sleeved shirt, long pants, shoes and
socks, chemical resistant gloves when handling. Wash hands before
eating, drinking, chewing gum, using tobacco, or using the toilet.
Remove clothing immediately if pesticide gets inside. Then wash
thoroughly and put on clean clothing wash clothing before reuse.

FIRST AID

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of
water for 15 minutes. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF SWALLOWED: Do not induce vomiting. Call physician or Poison
Control Center. Give person a large quantity of milk, egg white,
gelatin solution, or if these are not available, large quantities of water.
Avoid alcohol.

Note to physician: Probable mucosal damage may contraindicate the
use of gastric lavage.

ENVIRONMENTAL HAZARDS: This pesticide is extremely toxic to
freshwater, marine and estuarine fish, and aquatic invertebrates
including shrimp and oyster. Do not apply in a manner that will
directly expose canals, lakes, streams, ponds, marshes or estuaries
to aerial drift. Do not apply directly to water. Do not contaminate
water when disposing of equipment washwaters.

DIRECTIONS FOR USE: It is a violation of Federal law to use this
product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod,
or for commercial seed production, or for research purposes.

Entry restrictions: Do not allow people or pets to enter the treated area
until dusts have settled.

GENERAL INFORMATION: This product contains turf food plus a
preemergence herbicide for pre-emergence control of annual weed
grasses and broadleaf weeds in established turf. Apply to turf before
germination of weed seeds, since established weeds will not be
controlled.

WEEDS CONTROLLED:

Annual grasses: annual bluegrass (*Poa annua*), smooth crabgrass
(*Digitaria ischaemum*), large or hairy crabgrass (*Digitaria sanguinalis*),
goosegrass (silver crabgrass or crowfootgrass) (*Eleusine indica*),
barnyardgrass (watergrass) (*Echinochloa crus-galli*), green foxtail
(*Setaria viridis*), yellow foxtail (*Setaria glauca*).

Broadleaf weed: common chickweed (*Stellaria media*), prostrate
knotweed (*Polygonum aviculare*), common lambsquarter
(*Chenopodium album*), pigweed (*Amaranthus spp.*), common purslane
(*Portulaca oleracea*).

Weeds suppressed: Texas panicum (*Panicum texanum*) Florida pusley
(*Richardia scabra*), prostrate spurge (*Euphorbia humistrata*), spotted
spurge (*Euphorbia maculata*), creeping spurge (*Euphorbia serpens*),
creeping woodsorrel (*Oxalis corniculata*), wood sorrel (*Oxalis spp.*).

WHERE TO APPLY: This product may be applied to the following
established turfgrass species:

Established cool season turfgrass: bentgrass fairways, Kentucky
bluegrass, perennial ryegrass, fescues.

Established warm season turfgrass: bahia, bermuda, centipede, tall
fescue, St. Augustine, zoysia.

Important precautions: Do not apply this product to golf course putting
greens, dichondra, or newly sprigged area of bermuda, St. Augustine,
centipede or zoysiagrass. Apply only to healthy, established turf. Do
not apply in spring to turfgrass planted the previous fall. Will not harm
most nearby ornamental trees, shrubs and flowers. Do not apply this
product in a way that will contact any person or pet, either directly or
through drift. Keep people and pets out of the area during application.

(Label for EPA Reg. No 32802-35, PAGE 3 OF 4)


92719 (Household use)

WHEN TO APPLY: Successful preemergence weed control requires proper timing of the application. This product must be applied 1 to 2 weeks before the germination of weed seeds, since established weeds will not be controlled. All annual weed grasses do not germinate at the same time. For most effective weed control, apply at the proper time:

Northern areas:
For crabgrass, foxtail, barnyardgrass, etc., apply Mar. 1 through June 1
For goosegrass, which germinates later, apply May 15 through July 1
For annual bluegrass, apply July 1 through Oct. 1

Southern areas:
For crabgrass, foxtail, barnyardgrass, etc., apply Feb. 1 through June 1
For goosegrass, which germinates later, apply April 1 through July 1
For annual bluegrass, apply Oct. 1 through Feb. 1

(alternate graph to show appropriate application dates:)



APPLICATION RATES: Apply in spring applications to control spring-to-summer germinating annual weeds, or fall applications (to control fall-germinating weeds). A single application will usually provide season-long weed control. Apply this product at the rate of 4 to 8 pounds per 1000 square feet. Make a single application 1 to 2 weeks before the expected germination of weed seeds.

To control goosegrass or annual bluegrass, apply 5.25 to 8 pounds per 1000 square feet, in late spring or early summer 1 to 2 weeks prior to onset of conditions favorable to weed seed germination. Better control may be possible by using the higher rate.

Apply this product evenly over the turf, using a properly calibrated granular fertilizer spreader which will uniformly distribute the product. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur. For the most effective weed control, water or irrigate the treated areas soon after application. Cultural practices that disturb the soil, such as aerification, should be done before application.

SPI™ APPLICATIONS FOR FULL SEASON WEED CONTROL: A second or split application is recommended when the initial application is made well in advance of weed germination, or where an extended period of weed control is required (areas with longer growing seasons). In cool season turfgrass areas and the northern half of warm season turfgrass areas, apply 4 pounds per 1000 square feet plus a second application of 4 pounds per 1000 square feet. Make first application in spring 1 to 2 weeks prior to expected time of weed germination, and make the second application 8 to 10 weeks later. In the southern half of warm season turfgrass areas, apply 5.25 pounds per 1000 square feet plus a second application of 5.25 pounds per 1000 square feet. Make first application in spring 1 to 2 weeks prior to expected time of weed germination, and make the second application 10 to 12 weeks later.

RESEEDING: This product will inhibit germination of grass seed if seeded too soon after application. When reseeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. Delay reseeding treated areas for a minimum of:

- 8 to 10 weeks after last application of 4 pounds/1000 square feet
- 10 to 12 weeks after last application of 5.25 pounds/1000 square feet
- 12 to 16 weeks after last application of 8 pounds/1000 square feet

When soil testing or professional advice indicates the need for additional plant nutrients, make additional applications of fertilizer products over the growing season.

This product may stain driveways, walks, patios, pools, etc. Wash or sweep off thoroughly immediately after each use.

SPREADER SETTINGS to apply at the listed pounds per 1000 square feet			
	4 lbs	5.25 lbs	8 lbs
Spreader A	X	X	X
Spreader B	X	X	X
Spreader C	X	X	X
Spreader D	X	X	X

(May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in a locked storage area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Fold empty bag and discard in trash.

GUARANTEE: This product is unconditionally guaranteed.

(Alternate wording for warranty statement:)

NOTICE: Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

(Alternate wording for warranty statement:)

WARRANTY AND DISCLAIMER: Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label. Manufacturer/seller makes no other warranty, expressed or implied, and shall not be liable for indirect or consequential damages.

(Alternate wording for warranty statement:)

WARRANTY AND DISCLAIMER: Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label. No other warranty, expressed or implied, is made concerning the merchantability or fitness for a particular purpose. Manufacturer/seller's liability shall be limited to the purchase price of the product. In no case shall manufacturer/seller be liable for consequential or incidental damages or loss.

(Alternate wording for warranty statement:)

WARRANTY AND DISCLAIMER: Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. No other warranty, expressed or implied, is made concerning the merchantability or fitness for a particular purpose. Manufacturer/seller's liability shall be limited to the purchase price of the product. In no case shall manufacturer/seller be liable for consequential or incidental damages or loss. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label. Products not accepted under these terms should be returned within 20 days in the original, unopened container for return of purchase price.