

P/M23

32802-35

1/13/89

1 of 6

JAN 13 1989

Mr. Thomas J. Schmit  
Howard Johnson's Enterprises, Inc.  
700 W. Virginia Street, #222  
Milwaukee, WI 53204-1548

Dear Mr. Schmit:

Subject: Lawn Service Crabgrass Control

EPA Registration No. 32802-35

Proteck 58

EPA Registration No. 32802-40

Your 8 Month Response To the TNEPLAN RED, Product Chemistry and Acute Toxicology Studies and Your Response To the "Preliminary Label Assessment Enclosed in Our Letter Dated November 3, 1998, Notice of Reregistration Dated November 3, 1998 and Your letter with Final Printed Labeling Dated November 11, 1998

Registration of the subject pesticide products under Section 4 of the Federal Insecticide, Fungicide and Rodenticide Act, as amended on November 3, 1998 was in error because the Reregistration Evaluation Document (RED) for the active ingredient benefin has not been issued. The purpose of this letter is to correct this error and response to your subject responses as the Agency is presently practicing. The labeling submitted in your letters dated November 11, 1998 has been reviewed and found acceptable as amendments to the subject pesticide product registrations under Section 3 of the Federal Insecticide, Fungicide and Rodenticide Act, as amended; provided that you:

1. Revise the entry restrictions on each of the labels by adding: "If the treated area is watered or irrigated following application for most effective weed control, do not enter or allow others to enter the treated area until the surface is dry."
2. Revise the "General Information" and the "When To Apply" sentences that bear the phrase "since established weeds will not be controlled." to read: "because established weeds will not be controlled."
3. Submit one (1) copy of the final printed label before releasing these products for shipment under the subject labeling.

If these conditions are not complied with, these registrations will be subject to cancellation in accordance with FIFRA, Section

2 of 6  
-2-

6(e). Your release for shipment of these products constitutes acceptance of this condition.

A stamped copy of each of the labels is enclosed for your records.

Reregistration of these product under Section 4 of the Federal Insecticide, Fungicide and Rodenticide Act, as amended, must await the issuance of a Reregistration Evaluation Document (RED) for benefin which is an active ingredient in each of these products.

Sincerely yours,

Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505C)

Enclosures (4 stamped labels)

E.Wilson:Diskette ~~Trifluralin: 11-03-98~~ RED ABC7

ACTIVE INGREDIENTS:  
 benefin: . . . . . 0.58%  
 N-butyl-N-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*p*-toluidine . . . . . 0.58%  
 trifluralin: . . . . . 0.29%  
 $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-*N,N*-dipropyl-*p*-toluidine . . . . . 0.29%  
 INERT INGREDIENTS: . . . . . 99.13%  
 TOTAL: . . . . . 100.00%

EPA Reg. No. 32802-35

EPA Est. No. 32802-WI-1 (A), 32802-MO-1 (B)

First letter of bag code shows establishment where manufactured.

Manufactured by:

WARD JOHNSON'S ENTERPRISES, INC.

700 W. Virginia Street, Milwaukee, WI 53204-1548

\*TEAM is a trademark of Dow AgroSciences LLC

## KEEP OUT OF REACH OF CHILDREN

**WARNING:** See back panel for further cautions,  
 first aid information and directions for use.

NET WEIGHT: 20 pounds (9.0 kg)

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING:** Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through skin. Do not get in eyes or clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some persons. Wear gloves or face shield, long-sleeved shirt, long pants, shoes and socks, chemical resistant gloves when handling. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing wash clothing before reuse.

### FIRST AID

**IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention.

**IF SWALLOWED:** Do not induce vomiting. Call physician or Poison Control Center. Give person a large quantity of milk, egg white, gelatin solution, or if these are not available, large quantities of water. Avoid alcohol.

Note to physician: Probable mucosal damage may contraindicate the use of gastric lavage.

**ENVIRONMENTAL HAZARDS:** This pesticide is extremely toxic to freshwater, marine and estuarine fish, and aquatic invertebrates including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Entry restrictions: Do not allow people or pets to enter the treated area until dusts have settled.

**GENERAL INFORMATION:** This product contains turf food plus a preemergence herbicide for pre-emergence control of annual weed grasses and broadleaf weeds in established turf. Apply to turf before germination of weed seeds, since established weeds will not be controlled.

### WEEDS CONTROLLED:

Annual grasses: annual bluegrass (*Poa annua*), smooth crabgrass (*Digitaria ischaemum*), large or hairy crabgrass (*Digitaria sanguinalis*), goosegrass (silver crabgrass or crowfootgrass) (*Eleusine indica*), barnyardgrass (watergrass) (*Echinochloa crus-galli*), green foxtail (*Setaria viridis*), yellow foxtail (*Setaria glauca*).

Broadleaf weeds: common chickweed (*Stellaria media*), prostrate knotweed (*Polygonum aviculare*), common lambsquarter (*Chenopodium album*), pigweed (*Amaranthus spp.*), common purslane (*Portulaca oleracea*).

Weeds suppressed: Texas panicum (*Panicum texanum*) Florida pusley (*Richardia scabra*), prostrate spurge (*Euphorbia humistrata*), spotted

WHEN TO APPLY: Successful preemergence weed control requires proper timing of the application. This product must be applied 1 to 2 weeks before the germination of weed seeds, since established weeds will not be controlled. All annual weed grasses do not germinate at the same time. For most effective weed control, apply at the proper time:

Northern areas:

For crabgrass, foxtail, barnyardgrass, etc., apply Mar. 1 through June 1

For goosegrass, which germinates later, apply May 15 through July 1

For annual bluegrass, apply July 1 through Oct. 1

Southern areas:

For crabgrass, foxtail, barnyardgrass, etc., apply Feb. 1 through June 1

For goosegrass, which germinates later, apply April 1 through July 1

For annual bluegrass, apply Oct. 1 through Feb. 1

**APPLICATION RATES:** Apply in spring applications to control spring-to-summer germinating annual weeds, or fall applications (to control fall-germinating weeds). A single application will usually provide season-long weed control. Apply this product at the rate of 4 to 8 pounds per 1000 square feet. Make a single application 1 to 2 weeks before the expected germination of weed seeds.

To control goosegrass or annual bluegrass, apply 5.25 to 8 pounds per 1000 square feet, in late spring or early summer 1 to 2 weeks prior to onset of conditions favorable to weed seed germination. Better control may be possible by using the higher rate.

Apply this product evenly over the turf, using a properly calibrated granular fertilizer spreader which will uniformly distribute the product. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur. For the most effective weed control, water or irrigate the treated areas soon after application. Cultural practices that disturb the soil, such as aerification, should be done before application.

**SPLIT APPLICATIONS FOR FULL SEASON WEED CONTROL:** A second or split application is recommended when the initial application is made well in advance of weed germination, or where an extended period of weed control is required (areas with longer growing seasons). In cool season turfgrass areas and the northern half of warm season turfgrass areas, apply 4 pounds per 1000 square feet plus a second application of 4 pounds per 1000 square feet. Make first application in spring 1 to 2 weeks prior to expected time of weed germination, and make the second application 8 to 10 weeks later.

In the southern half of warm season turfgrass areas, apply 5.25 pounds per 1000 square feet plus a second application of 5.25 pounds per 1000 square feet. Make first application in spring 1 to 2 weeks prior to expected time of weed germination, and make the second application 10 to 12 weeks later.

**RESEEDING:** This product will inhibit germination of grass seed if seeded too soon after application. When reseeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. Delay reseeding treated areas for a minimum of:

8 to 10 weeks after last application of 4 pounds/1000 square feet  
 10 to 12 weeks after last application of 5.25 pounds/1000 square feet  
 12 to 16 weeks after last application of 8 pounds/1000 square feet

This product may stain driveways, walks, patios, pools, etc. Wash or sweep off thoroughly immediately after each use.

### SPREADER SETTINGS:

The settings shown here are approximate for new equipment. Your spreader may need calibration, due to differences in terrain, walking speed, and equipment condition. Calibrate spreader to ensure correct application rate.

Spreader setting to apply at the listed rate (pounds per 1000 sq. ft.)	4 lbs.	5.25 lbs.	8 lbs.
Ames	15	17	22
EarthWay	15	17	22
Precision Red Devil	8	10	12
Cyclone SB3000 Rotary	7	7.25	8
Scotts AccuGreen Drop	3.5	4	5
Scotts EvenGreen Drop	8	9	11
Scotts EasyGreen Rotary	28	29	30
Scotts SpeedyGreen Rotary	7	8	12

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in original container in a locked storage area out of reach of children and domestic animals.

**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse bag. Fold empty bag and discard in trash.

**NOTICE:** Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

92718 (House)

4 of 6

**ACCEPTED**  
**with COMMENTS**  
**In EPA Letter Dated**

**JAN 13 1999**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

32802-35

CH

"Occupational Use" Label Version

# LAWN SERVICE CRABGRASS CONTROL

WITH TEAM® HERBICIDE FOR PRE-EMERGENCE  
CONTROL OF CRABGRASS, FOXTAIL, GOOSEGRASS,  
ANNUAL BLUEGRASS (*POA ANNUA*) AND OTHER  
ANNUAL WEED GRASSES AND BROADLEAF WEEDS  
IN ESTABLISHED LAWNS AND ORNAMENTAL TURF

ACTIVE INGREDIENTS:

benefin:  
N-butyl-N-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-p-toluidine . . . . . 0.58%  
trifluralin:  
 $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine . . . . . 0.29%  
INERT INGREDIENTS: . . . . . 99.13%  
TOTAL: . . . . . 100.00%

Contains 0.435 pounds active ingredient per 50 pound bag

EPA Reg. No. 32802-35

EPA Est. No. 32802-WI-1 (A), 32802-MO-1 (B)

First letter of bag code shows establishment where manufactured.

See tag attached to bag for fertilizer guaranteed analysis.

Manufactured by:

HOWARD JOHNSON'S ENTERPRISES, INC.

W. Virginia Street, Milwaukee, WI 53204-1548

\*TEAM is a trademark of Dow AgroSciences LLC

## KEEP OUT OF REACH OF CHILDREN

**WARNING:** See back panel for further cautions,  
first aid information and directions for use.

NET WEIGHT: 50 pounds (22.7kg)

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING:** Causes substantial but temporary eye injury.  
Harmful if swallowed or absorbed through skin. Do not get in eyes or  
clothing. Avoid contact with skin. Prolonged or frequently repeated  
skin contact may cause allergic reactions in some persons.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):** Applicators and  
other handlers must wear goggles or face shield, long-sleeved shirt,  
pants, shoes and socks, chemical resistant gloves when  
handling. Wash hands before eating, drinking, chewing gum, using  
tobacco, or using the toilet. Remove contaminated clothing and  
wash clothing before reuse. Follow manufacturer's instructions for  
cleaning/maintaining PPE. If no such instructions for washables, use  
detergent and hot water. Keep and wash PPE separately from other  
laundry.

**USER SAFETY RECOMMENDATIONS:** Users should remove  
clothing immediately if pesticide gets inside. Then wash  
thoroughly and put on clean clothing. User should remove PPE  
immediately after handling this product. Wash the outside of  
gloves before removing. As soon as possible, wash thoroughly  
and change into clean clothing.

## FIRST AID

**IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of  
water for 15 minutes. Get medical attention.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention.

**IF SWALLOWED:** Do not induce vomiting. Call physician or Poison  
Control Center. Give person a large quantity of milk, egg white,  
gelatin solution, or if these are not available, large quantities of water.  
Avoid alcohol.

**Note to Physician:** Probable mucosal damage may contraindicate the  
use of gastric lavage.

**ENVIRONMENTAL HAZARDS:** This pesticide is extremely toxic to  
freshwater, marine and estuarine fish, and aquatic invertebrates  
including shrimp and oyster. Do not apply in a manner that will  
directly expose canals, lakes, streams, ponds, marshes or estuaries  
to aerial drift. Do not apply directly to water, or to areas where  
surface water is present or to intertidal areas below the mean high  
water mark. Do not contaminate water when disposing of equipment  
washwaters.

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this  
product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod,  
or for commercial seed production, or for research purposes.

**Entry restrictions:** Do not enter or allow others to enter the treated area  
until dusts have settled.

**GENERAL INFORMATION:** This product contains turf food plus a  
preemergence herbicide for pre-emergence control of annual weed  
grasses and broadleaf weeds in established turf. Apply to turf before  
germination of weed seeds, since established weeds will not be  
controlled.

## WEEDS CONTROLLED:

Annual grasses: annual bluegrass (*Poa annua*), smooth crabgrass  
(*Digitaria ischaemum*), large or hairy crabgrass (*Digitaria sanguinalis*),  
goosegrass (silver crabgrass or crowfootgrass) (*Eleusine indica*),  
barnyardgrass (watergrass) (*Echinochloa crus-galli*), green foxtail

(*Setaria viridis*), yellow foxtail (*Setaria glauca*).

Broadleaf weeds: common chickweed (*Stellaria media*), prostrate  
knotweed (*Polygonum aviculare*), common lambsquarter  
(*Chenopodium album*), pigweed (*Amaranthus spp.*), common purslane  
(*Portulaca oleracea*).

Weeds suppressed: Texas panicum (*Panicum texanum*) Florida pusley  
(*Richardia scabra*), prostrate spurge (*Euphorbia humistrata*), spotted  
spurge (*Euphorbia maculata*), creeping spurge (*Euphorbia serpens*),  
creeping woodsorrel (*Oxalis corniculata*), woodsorrel (*Oxalis spp.*).

**WHERE TO APPLY:** This product may be applied to the following  
established turfgrass species:

Established cool season turfgrass: bentgrass fairways, Kentucky  
bluegrass, perennial ryegrass, fescues.

Established warm season turfgrass: bahia, bermuda, centipede, tall  
fescue, St. Augustine, zoysia.

**Important precautions:** Do not apply this product to golf course putting  
greens, dichondra, or newly sprigged area of bermuda, St. Augustine,  
centipede or zoysiagrass. Apply only to healthy, established turf. Do  
not apply in spring to turfgrass planted the previous fall. Will not harm  
most nearby ornamental trees, shrubs and flowers. Do not apply this  
product in a way that will contact workers or other persons, either  
directly or through drift. Only protected handlers may be in the area  
during application.

**WHEN TO APPLY:** Successful preemergence weed control requires  
proper timing of the application. This product must be applied 1 to 2  
weeks before the germination of weed seeds, since established weeds  
will not be controlled. All annual weed grasses do not germinate at the  
same time. For most effective weed control, apply at the proper time:

**Northern areas:**

For crabgrass, foxtail, barnyardgrass, etc., apply Mar. 1 through June 1

For goosegrass, which germinates later, apply May 15 through July 1

For annual bluegrass, apply July 1 through Oct. 1

**Southern areas:**

For crabgrass, foxtail, barnyardgrass, etc., apply Feb. 1 through June 1

For goosegrass, which germinates later, apply April 1 through July 1

For annual bluegrass, apply Oct. 1 through Feb. 1

**APPLICATION RATES:** Apply in spring applications to control spring-  
to-summer germinating annual weeds, or fall applications (to control  
fall-germinating weeds). A single application will usually provide  
season-long weed control. Apply this product at the rate of 4 to 8  
pounds per 1000 square feet. Make a single application 1 to 2 weeks  
before the expected germination of weed seeds.

To control goosegrass or annual bluegrass, apply 5.25 to 8 pounds per  
1000 square feet, in late spring or early summer 1 to 2 weeks prior to  
onset of conditions favorable to weed seed germination. Better control  
may be possible by using the higher rate.

Apply this product evenly over the turf, using a properly calibrated  
granular fertilizer spreader which will uniformly distribute the product.  
Avoid skips and excessive overlaps as poor weed control or turfgrass  
injury may occur. For the most effective weed control, water or irrigate  
the treated areas soon after application. Cultural practices that disturb  
the soil, such as aerification, should be done before application.

**SPLIT APPLICATIONS FOR FULL SEASON WEED CONTROL:** A  
second or split application is recommended when the initial application  
is made well in advance of weed germination, or where an extended  
period of weed control is required (areas with longer growing seasons).  
In cool season turfgrass areas and the northern half of warm season  
turfgrass areas, apply 4 pounds per 1000 square feet plus a second  
application of 4 pounds per 1000 square feet. Make first application in  
spring 1 to 2 weeks prior to expected time of weed germination, and  
make the second application 8 to 10 weeks later.

In the southern half of warm season turfgrass areas, apply 5.25 pounds  
per 1000 square feet plus a second application of 5.25 pounds per  
1000 square feet. Make first application in spring 1 to 2 weeks prior to  
expected time of weed germination, and make the second application 10  
to 12 weeks later.

**RESEEDING:** This product will inhibit germination of grass seed if  
seeded too soon after application. When reseeding, follow good  
cultural practices such as soil cultivation, irrigation and fertilization.  
Delay reseeding treated areas for a minimum of:  
8 to 10 weeks after last application of 4 pounds/1000 square feet  
10 to 12 weeks after last application of 5.25 pounds/1000 square feet  
12 to 16 weeks after last application of 8 pounds/1000 square feet

This product may stain driveways, walks, patios, pools, etc. Wash or  
sweep off thoroughly immediately after each use.

## SPREADER SETTINGS:

The settings shown here are  
approximate for new equipment  
Your spreader may need calibration,  
due to differences in terrain, walking  
speed, and equipment condition.  
Calibrate spreader to ensure correct  
application rate.

Spreader setting to apply at the listed rate (pounds per 1000 sq ft)	4 lbs	5.25 lbs	8 lbs
Spyker 24-22 Rotary	4.1	4.5	5.5
Lesco Rotary	12(F)	14(G)	18(I)
Prizelawn CB Rotary	4.5	4.75	6.0
EarthWay 2400 Rotary	18	19	20
Cyclone Pro	4.0	4.5	5.5
Scotts R&A	1	J	M
Lely Broadcast	3.0	3.4	4.2

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in original container in a locked storage area out  
of reach of children and domestic animals.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this  
product may be disposed of on site or at an approved waste disposal  
facility.

**CONTAINER DISPOSAL:** Completely empty bag into application  
equipment. Then dispose of empty bag in a sanitary landfill, or by  
incineration, or, if allowed by State and local authorities, by burning.  
If burned, stay out of smoke.

**NOTICE:** Buyer and user assumes all risks and liability of use, storage  
and/or handling of this product not in accordance with the terms of this  
label.

6076

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

JAN 13 1999

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
**32802-35**