

32802-36

106 b
PM 21

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460 NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	EPA REGISTRATION NO. 32802-36	DATE OF ISSUANCE AUG 28 1986
	TERM OF ISSUANCE	
	NAME OF PESTICIDE PRODUCT PCRB 10-C SOIL FURIGICIDE	

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Howard Johnson's Enterprises, Inc.
 Railroad Avenue
 P.O. Box 828
 Viroqua, WI 54665

160 175373
18 L

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment:
 - a. Add the phrase "EPA Registration No. 32802-36."
 - b. Under lawn, for snow mold control, revise the lower dosage rate to 5 lb/1000 sq ft.
 - c. Under ornamentals, the paragraph following the entry for calandulas, etc. appears to be directions for use on Dutch iris for control of bulb rot and stem rot (Sclerotium rolfsii). Add the intended crop and diseases for this entry.
 - d. For Easter lily, include an equivalent rate for smaller areas.
 - e. Many of the dosage rates included on the label would not appear to be appropriate for an 8 lb package. Such a small package

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL: *Henry M. Sacks* DATE: 8/28/86

would appear to be intended for use on smaller areas. Revise the dosage rates to be more appropriate for the package size.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.



Henry M. Jacoby
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (TS-767C)

Enclosures

P C N B 1 0 - G
S O I L F U N G I C I D E

FOR CONTROL OF SOIL BORNE DISEASES AS LISTED

ON LAWNS & TURF - COTTON - BEANS - CABBAGE - CAULIFLOWER
BROCCOLI - PEANUTS - POTATOES - ORNAMENTAL PLANTINGS

ACTIVE INGREDIENT:	
Pentachloronitrobenzene	10.0%
INERT INGREDIENTS:	90.0%
Total	100.0%

ACCEPTED
with COMMENTS
in EPA Letter Dated:

AUG 28 1986

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
32802-36

STATEMENTS OF PRACTICAL TREATMENT

- IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
- IF IN EYES: Flush eyes with plenty of water. Get medical attention immediately.
- IF ON SKIN: Remove contaminated clothing and wash affected area with soap & water.
- IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated. Get medical attention.


SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REF. NO 32802-__

EPA EST. NO. 32802-WI-1
-MO-1

NET WEIGHT 8 POUNDS (40 lb. bag)

HOWARD JOHNSON'S ENTERPRISES, INC.
VIROQUA, WI 54665



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed. Avoid inhaling dust. This product may cause skin irritation. Wash hands & face thoroughly after using and before eating or smoking. Wear the following protective clothing during mixing/loading operations: Gloves, Longsleeved shirt, Long Pants, Socks and Shoes. Do not contaminate feed or foodstuffs.

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. This product is toxic to fish. Do not apply directly to any body of water. Do not apply directly adjacent to potable water supplies.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not plant any root crop not registered for PCNB in rotation on PCNB treated soil.

STORAGE AND DISPOSAL


Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Store product in its original container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved wasted disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

DIRECTIONS FOR USE (CONT.)

COTTON: For the control of Damping-off, Seed and Seedling Rots -- **APPLY AS AN "IN-FURROW SOIL TREATMENT"** for Rhizoctonia solani: Apply 10 to 20 lbs. per acre as an "In-Furrow" soil application which is equivalent to 1 to 2 lbs. per 1300 linear feet of row based on a 40 inch spaced row. By using suitable application equipment apply over the seed and immediately cover with the seed at planting time. Do not allow livestock to graze or feed on cotton foliage in treated fields.

BEANS: Root and Stem Rot- Rhizoctonia solani: **APPLY AS AN "IN-FURROW SOIL TREATMENT"** - Apply 10 to 15 lbs. per 14,500 linear feet of row for Bush Beans or per 8,340 Sq.Ft. of row for Pole Beans. Apply over the seed at planting time in the furrow and immediately cover with soil utilizing appropriate application equipment. Do not feed treated vines to livestock.

CRUCIFERS - CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, AND BROCCOLI - Clubroot (Plasmodiophora brassicae): **APPLY AS A "PRE-TRANSPLANTING ROW TREATMENT"**. Apply 200 lbs. per 13,100 feet of row for Cabbage, Broccoli and Brussels Sprouts or to 10,900 linear feet of row for Cauliflower, in a 12 to 15 inch band. Apply & rototill the granules to a depth of 4 to 6 inches just before transplanting. - - - - - **TO APPLY AS A "BROADCAST TREATMENT"** Apply 300 lbs. per acre to the soil surface prior to transplanting, then disc the granules into the soil to a depth of 4 to 6 inches. Thorough mixing with the soil will give best disease control. Double cross-discing is preferred.



CAULIFLOWER: For control of Wire stem or Black Rot - *Corticium solani* -
APPLY AS A BAND ROW TREATMENT - Using 110 to 150 lbs. per 10,900 linear feet
of row in an 8 inch wide band centered on the row, immediately before
planting. For small areas use 12 lbs. per 1000 Sq.Ft. Mix granules into
soil before planting to a depth of 2 inches by rototilling or discing.

PEANUTS: For Southern Blight - *Sclerotium rolfsii* - **AS A SURFACE BAND
TREATMENT** use 100 lbs. per 12,400 linear feet of row, in an 8 to 12 inch wide
band, centered on the row at pegging time. **AS A PRE-PLANT SOIL TREATMENT:**
Use 100 lbs. per 12,400 linear feet of row, in an 8 to 12 inch band,
centered on the row. Mix to a depth of 2 to 4 inches. This should be done
simultaneously using appropriate equipment. **AS A SOIL MIX "DURING
CULTIVATION"** it may be applied in three split applications when cultivating
the crop at the rate of 33 lbs. per 12,400 Sq.Ft. per treatment by attaching
two delivery tubes per row ahead of inside cultivator sweeps (One on each
side) and adjusted so as to treat a total band of 12 inches wide, centered on
the row. Do not pile untreated soil on crown of plant. **NOTE:** Do not exceed 10
lbs. actual PCNB per acre in any one season, regardless of the application
methods used. Do not feed treated vines to livestock. Do not allow hogging
down of peanuts.

POTATOES: *Rhizoctonia solani* - **AS AN OVER-ALL BROADCAST TREATMENT** -
Apply 180 to 250 lbs. per acre over the soil surface prior to planting and
thoroughly disc to a depth of 4 to 6 inches by rototilling or "cross-discing".
AS AN "IN-FURROW" TREATMENT - Apply 100 lbs/12,400 linear feet of a 36 inch
row by thoroughly working granules to a depth of 4 to 6 inches & into the
tuber-forming area of the soil. It is best to apply when the soil is
slightly moist.

IN LAWNS AND ORNAMENTAL TURF

FOR BROWN PATCH (*RHIZOCTONIA SOLANI*)

SOUTHERN GRASSES - For St. Augustine, Bermuda, etc. and *Dichondra* apply 7½
lbs./1,000 Sq.Ft. in the fall or the spring at the first sign of infrection.
Lightly sprinkle the area to wash the granules to soil level. An additional
application in 3 to 4 weeks may be needed for continued control if heavy rain-
fall or flooding occurs after this treatment. Apply at the same rate.

NORTHERN GRASSES - For Bluegrass, Fescue, Rye, Bent, etc. apply 2 to 2 1/2
lbs. per 1,000 Sq.Ft. when disease first appears and repeat at 7 to 10 day
intervals during warm, damp weather. Lightly sprinkle the area to wash
granules to soil level. Under certain climatic conditions a temporary
dicoloration of grass may occur however this is not harmful and will
disappear in a few days.

HELMINTHOSPORIUM LEAF SPOT

FOR CONTROL OF "LEAF SPOT" ON SOUTHERN OR NORTHERN GRASSES: Apply at the rate
of 5 to 7½ lbs./1,000 Sq.Ft. Preventative applications should be made in the
spring or fall of each year. For control for existing infections, apply at
the higher rate (7½ lbs.) as soon as symptoms are first noted. After
applying, the treated area should be lightly watered to insure movement
through the grass to soil level. If treated area is subjected to unusually
heavy rainfall, or flooded, or if the disease is severe or reappears, the
areas should be retreated 3 to 4 weeks later.

SNOW MOLD and FUSARIUM PATCH (*TYPHULA* Species & *FUSARIUM NIVALE*)

FOR CONTROL ON NORTHERN GRASSES: Apply at the rate of 45 to 7 1/2 lbs. per
1,000 Sq.Ft. using a granular lawn spreader or comparable equipment.
Application should be made prior to the first snowfall. Do not graze treated
areas or feed clippings to livestock.

DIRECTIONS FOR USE (Cont.)

ORNAMENTAL PLANTS

BEDDING PLANTS: For African Violets, Begonias, Carnations, Chrysanthemums, Poinsettias, Snapdragons and similar bedding plants treat soil prior to planting with 15 lbs. granules / 1,000 Sq.Ft. for control of RHIZOCTONIA, STEM ROT. Thoroughly work into the top 2 inches of soil by rototilling.

BULBS: For Tulips, Narcissus, Hyacinth, and Bulbous Iris apply granules using 1,000 to 2,000 lbs. per acre to the soil as a pre-plant broadcast application for control of CROWN ROT and BLACK ROT. (Equivalent to 2.25 to 4.5 lbs./100 Sq.Ft.). Mix with the top 6 to 7 inches of soil with appropriate equipment. Use the higher rate in heavy organic soils and especially for the control of Black Rot.


CALENDULAS, LARKSPUR, SNAPDRAGON, SWEET PEA - For control of SCLEROTINA STEM ROT apply 1,000 lbs./acre as a Broadcast application 1 week before planting; (Equiv. to 2.25 lbs./ 100 Sq.Ft.). Mix with soil to a depth of 4 inches.

make a PRE-PLANTING BROADCAST SOIL APPLICATION of 1,000 to 1,500 lbs. per acre which is equivalent to 2.25 to 3.5 lbs. /100 Sq.Ft. Mix the granules with the soil to a depth of 2 to 3 inches. FOR A ROW "BAND TREATMENT" apply 60 lbs. per 14,500 Sq.Ft. of linear row in a 12 inch band centered on the row, at planting time. Follow this treatment with 3 additional band applications at 15 day intervals at the same rate. Use the low rate for light sandy soils.

EASTER LILY: For control of Rhizoctonia Root Rot apply AS AN OVER-ALL BROADCAST TREATMENT at the rate of 1,000 to 2,000 lbs. per acre. Use the low rate for sandy light soils. Follow this application with rototilling or double discing to mix granules with soil to a depth of 6 to 7 inches or to bulb planting depth.

GLADIOLUS: For Neck or Corm Dry Rot caused by Stromatinia or Sclerotinia Gladioli apply 1,200 lbs. per acre (equivalent to 2 3/4 lbs. per 100 Sq.Ft.) AS A BROADCAST SOIL TREATMENT. Thoroughly mix with soil to depth of 4 to 6 inches. Corms may be planted immediately.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.



P C N B 1 0 - G

S O I L F U N G I C I D E

FOR CONTROL OF SOIL BORNE DISEASES AS LISTED

ON LAWNS & TURF - COTTON BEANS - CABBAGE - CAULIFLOWER
BROCCOLI - PEANUTS - POTATOES - ORNAMENTAL PLANTINGS

ACTIVE INGREDIENT:

Pentachloronitrobenzene	10.0%
INERT INGREDIENTS:	90.0%
Total	100.0%

NOT FINISHED
In Accordance with...
Based on Draft...

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

STATEMENTS OF PRACTICAL TREATMENT

- IF SWALLOWED: Call a physician or Poison Control Center immediately. vomiting by giving victim 1 or 2 glasses of water and touching back of with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
- IF IN EYES: Flush eyes with plenty of water. Get medical attention immediately.
- IF ON SKIN: Remove contaminated clothing and wash affected area with water.
- IF INHALED: Remove victim to fresh air. Apply artificial respiration indicated. Get medical attention.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 32802-36

EPA EST. NO. 32802-

NET WEIGHT 8 POUNDS [Package size For Use On Ornamental Turf, Onl
NET WEIGHT 40 POUNDS [Package Size For All Other Product Uses.]

HOWARD JOHNSON'S ENTERPRISES, INC
VIROQUA, WI 54665

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhaling dust. This product may cause skin irritation. Wash hands & face thoroughly after using and before eating or smoking. Wear the following protective clothing during mixing/loading operations: Gloves, Longsleeved shirt, Long Pants, Socks and Shoes. Do not contaminate feed or foodstuffs.

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. This product is toxic to fish. Do not apply directly to any body of water. Do not apply directly adjacent to potable water supplies.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not plant any root crop not registered for PCNB in rotation on PCNB treated soil.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. STORAGE: Store product in its original container in a cool, dry, locked place out of reach of children. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

DIRECTIONS FOR USE (CONT.)

COTTON: For the control of Damping-off, Seed and Seedling Rots -- APPLY AS AN "IN-FURROW SOIL TREATMENT" for Rhizoctonia solani: Apply 10 to 20 lbs. per acre as an "In-Furrow" soil application which is equivalent to 1 to 2 lbs. per 1300 linear feet of row based on a 40 inch spaced row. By using suitable application equipment apply over the seed and immediately cover with the seed at planting time. Do not allow livestock to graze or feed on cotton foliage in treated fields.

BEANS: Root and Stem Rot- Rhizoctonia solani: APPLY AS AN "IN-FURROW SOIL TREATMENT" - Apply 10 to 15 lbs. per 14,500 linear feet of row for Bush Beans or per 8,340 Sq.Ft. of row for Pole Beans. Apply over the seed at planting time in the furrow and immediately cover with soil utilizing appropriate application equipment. Do not feed treated vines to livestock.

CRUCIFERS - CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, AND BROCCOLI: Clubroot (Plasmodiophora brassicae): APPLY AS A "PRE-TRANSPLANTING ROW TREATMENT". Apply 200 lbs. per 13,100 feet of row for Cabbage, Broccoli and Brussels Sprouts or to 10,900 linear feet of row for Cauliflower, in a 12 to 15 inch band. Apply & rototill the granules to a depth of 4 to 6 inches just before transplanting. - - - - TO APPLY AS A "BROADCAST TREATMENT" - Apply 300 lbs. per acre to the soil surface prior to transplanting, then disc the granules into the soil to a depth of 4 to 6 inches. Thorough mixing with the soil will give best disease control. Double cross-discing is preferred.

BEST AVAILABLE COPY

DIRECTIONS FOR USE (Cont.)

CAULIFLOWER: For control of Wire stem or BLACK ROT - CORTICIUM SOLANI -
APPLY AS A BAND ROW TREATMENT - Using 110 to 150 lbs. per 10,900 linear feet
of row in an 8 inch wide band centered on the row, immediately before
planting. For small areas use 12 lbs. per 1000 Sq.Ft. Mix granules into
soil before planting to a depth of 2 inches by rototilling or discing.

PEANUTS: For SOUTHERN BLIGHT - SCLEROTIUM ROLFII - AS A SURFACE BAND
TREATMENT use 100 lbs. per 12,400 linear feet of row, in an 8 to 12 inch wide
band, centered on the row at pegging time. AS A PRE-PLANT SOIL TREATMENT:
Use 100 lbs. per 12,400 linear feet of row, in an 8 to 12 inch band,
centered on the row. Mix to a depth of 2 to 4 inches. This should be done
simultaneously using appropriate equipment. AS A SOIL MIX "DURING
CULTIVATION" it may be applied in three split applications when cultivating
the crop at the rate of 33 lbs. per 12,400 Sq.Ft. per treatment by attaching
two delivery tubes per row ahead of inside cultivator sweeps (One on each
side) and adjusted so as to treat a total band of 12 inches wide, centered on
the row. Do not pile untreated soil on crown of plant. NOTE: Do not exceed 10
lbs. actual PCNB per acre in any one season, regardless of the application
methods used. Do not feed treated vines to livestock. Do not allow hogging
down of peanuts.

POTATOES: RHIZOCTONIA SOLANI - AS AN OVER-ALL BROADCAST TREATMENT -
Appl 180 to 250 lbs. per acre over the soil surface prior to planting and
thoroughly disc to a depth of 4 to 6 inches by rototilling or "cross-discing".
AS AN "IN-FURROW" TREATMENT - Apply 100 lbs/12,400 linear feet of a 36 inch
row by thoroughly working granules to a depth of 4 to 6 inches & into the
tuber-forming area of the soil. It is best to apply when the soil is
slightly moist.

IN LAWNS AND ORNAMENTAL TURF

FOR BROWN PATCH (RHIZOCTONIA SOLANI)

SOUTHERN GRASSES - For St. Augustine, Bermuda, etc. and Dichondra apply 7½
lbs./1,000 Sq.Ft. in the fall or the spring at the first sign of infrection.
Lightly sprinkle the area to wash the granules to soil level. An additional
application in 3 to 4 weeks may be needed for continued control if heavy rain-
fall or flooding occurs after this treatment. Apply at the same rate.

NORTHERN GRASSES - For Bluegrass, Fescue, Rye, Bent, etc. apply 2 to 2½
lbs. per 1,000 Sq.Ft. when disease first appears and repeat at 7 to 10 day
intervals during warm, damp weather. Lightly sprinkle the area to wash
granules to soil level. Under certain climatic conditions a temporary
discoloration of grass may occur however this is not harmful and will
disappear in a few days.

HELMINTHOSPORIUM LEAF SPOT

FOR CONTROL OF "LEAF SPOT" ON SOUTHERN OR NORTHERN GRASSES: Apply at the rate
of 5 to 7½ lbs./1,000 Sq.Ft. Preventative applications should be made in the
spring or fall of each year. For control for existing infections, apply at
the higher rate (7½ lbs.) as soon as symptoms are first noted. After
applying, the treated area should be lightly watered to insure movement
through the grass to soil level. If treated area is subjected to unusually
heavy rainfall, or flooded, or if the disease is severe or reappears, the
areas should be retreated 3 to 4 weeks later.

SNOW MOLD and FUSARIUM PATCH

(TYPHULA Species & FUSARIUM NIVALE)

FOR CONTROL ON NORTHERN GRASSES: Apply at the rate of 5 to 7½ lbs. per
1,000 Sq.Ft. using a granular lawn spreader or comparable equipment.
Application should be made prior to the first snowfall. Do not graze treated
areas or feed clippings to livestock.

DIRECTIONS FOR USE (Cont.)

ORNAMENTAL PLANTS

BEDDING PLANTS: For African Violets, Begonias, Carnations, Chrysanthemums, Poinsettias, Snapdragons and similar bedding plants treat soil prior to planting with 7½ lbs. granules / 500 Sq.Ft. for control of RHIZOCTONIA, STEM POT. Thoroughly work into the top 2 inches of soil by rototilling.

BULBS: For Tulips, Narcissus, Hyacinth, and Bulbous Iris apply granules using 1,000 to 2,000 lbs. per acre to the soil as a pre-plant broadcast application for control of CROWN ROT and BLACK ROT. (Equivalent to 2½ to 4½ lbs./100 Sq.Ft.). Mix with the top 6 to 7 inches of soil with appropriate equipment. Use the higher rate in heavy organic soils and especially for the control of Black Rot.

CALENDULAS, LARKSPUR, SNAPDRAGON, SWEET PEA - For control of SCLEROTINIA STEM ROT apply 1,000 lbs./acre as a Broadcast application 1 week before planting; (Equiv. to 2½ lbs./ 100 Sq.Ft.). Mix with soil to a depth of 4 inches.

DUTCH IRIS: For Bulb and Stem Rot caused by SCLEROTIUM ROLESII disease make a PRE-PLANTING BROADCAST SOIL APPLICATION of 1,000 to 1,500 lbs. per acre which is equivalent to 2½ to 3½ lbs. /100 Sq.Ft. Mix the granules with the soil to a depth of 2 to 3 inches. FOR A ROW "BAND TREATMENT" apply 60 lbs. per 14,500 Sq.Ft. of linear row in a 12 inch band centered on the row, at planting time (equivalent to 1 lb per 240 Sq.Ft. of liner row) Follow this treatment with 3 additional band applications at 15 day intervals at the same rate. Use the low rate for light sandy soils.

EASTER LILY: For control of RHIZOCTONIA ROOT ROT apply AS AN OVER-ALL BROADCAST TREATMENT at the rate of 1,000 to 2,000 lbs. per acre. (equivalent to 2½ to 4½ lbs. per 100 Sq.Ft.) Use the low rate for sandy light soils. Follow this application with rototilling or double discing to mix granules with soil to a depth of 6 to 7 inches or to bulb planting depth.

GLADIOLUS: For Neck or Corm Dry Rot caused by STROMATINIA or SCLEROTINIA GLADIOLI apply 1,200 lbs. per acre (equivalent to 2½ lbs. per 100 Sq.Ft.) AS A BROADCAST SOIL TREATMENT. Thoroughly mix with soil to depth of 4 to 6 inches. Corms may be planted immediately.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.