

Reg # 32802 - 39

PM - 19

12/1

DURSBAN .5 GRANULES INSECTICIDE

LAWN AND TURF INSECT CONTROL KILLS
SOD WEBWORMS, WHITE GRUBS, ANTS, FLEAS,
BROWN DOG TICKS, CRICKETS, CHINCH BUGS,
CUTWORMS, EARWIGS, SOWBUGS, AND ARMYWORMS
EUROPEAN CRANE FLY LARVAE
HYPERODES (TURFGRASS) WEEVIL,
IN HOME LAWNS AND ORNAMENTAL TURF


FOR CONTROL OF IMPORTED FIRE ANTS BY
DIRECT MOUND TREATMENT

FOR PREMISE TREATMENT AROUND THE HOME
FOR BROWN DOG TICKS — FLEAS — ANTS —
CRICKETS — SOWBUGS

ACTIVE INGREDIENT:

CHLORPYRIFOS: (0, 0,-diethyl 0-
(3,5,6-trichloro-2-pyridyl) phosphorothioate) 0.50%
INERT INGREDIENTS: 99.50%

DURSBAN® is the Registered Trademark of The Dow/Elanco Chemical Company.



KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
 IF IN EYES: Flush eyes with plenty of water. Get medical attention.
 IF ON SKIN: Remove contaminated clothing and wash affected area with soap and warm water. Wash clothing before re-use.
 IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated. Get medical attention.
 NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine by injection only is an antidote.

EPA Reg. No. 32802-39

EPA EST. NO. 32802-WI-1(A)
32802-MO-1(B)

The first letter of code indicates location of production facility.

NET WEIGHT 50 POUNDS**LAWNS AND ORNAMENTAL TURF**
SURFACE FEEDING INSECTS

For control of Chinch Bugs, Sod Webworms (Lawn Moth), Ants, Fleas, Armyworms, Brown Dog Ticks, Crickets, Cutworms, Earwigs, Sowbugs and Millipedes apply 4.5 lbs. to treat 1,000 Sq. Ft. of turf. (50 lb. bag treats 11,000 Sq. Ft.) Refer to Spreader Settings. Treat in the spring or when insects first appear troublesome and repeat as needed for control. Additional treatments at 30 day intervals may be needed for season long control of Sod Webworms. After application lightly sprinkle the area to move the granules into the zone of insect infestation. Do not allow children or pets on treated areas until the grass is dry.

FOR ROOT FEEDING INSECTS**WHITE GRUB LARVAE CONTROL IN THE TURF**

FOR ROOT FEEDING INSECTS such as BILLBUG LARVAE, and the "White Grub" larvae of JAPANESE BEETLE, EUROPEAN CHAFER, SOUTHERN CHAFER AND HYPERODES WEEVIL apply 9 lbs./1,000 Sq. Ft. of turf (25 lb. bag treats 2,750 Sq. Ft.). Make sure your spreader is properly calibrated to apply this product. Irrigate the turf after treatment as soon as possible to wash the insecticide into the root zone where the insects are feeding. Apply when the White Grub Larvae are young and actively feeding near the soil surface in mid July and early August. For the HYPERODES (Turfgrass) WEEVIL larvae apply in Mid-April and again in Mid-May at the same rate as above. After application do not allow children or pets on treated areas until insecticide has been washed into the soil and the grass is dry. Consult your Agricultural Extension Service for the best time to treat for "Grubs" in your area. Refer to the Spreader Settings Chart.

EUROPEAN CRANE FLY LARVAE
(Leatherjackets)

For specific control of the Crane Fly Larvae feeding in western states turf-grass apply 6.8 lbs. to cover 1,000 sq. ft. The larvae will also be controlled simultaneously when an application rate for Grubs (9 lbs./1,000 Sq. Ft.) has been made. Consult your Agricultural Extension Service for the best time to treat in your area. Keep children and pets off treated area until granules have been washed into soil and the grass is dry.

IMPORTED FIRE ANT
TO CONTROL IMPORTED FIRE ANTS BY
DIRECT MOUND APPLICATION

For control of Imported Fire Ants by individual Ant mound treatment use a total of 3 ounces of this product per mound of granules evenly in a circle around the outer edges (1 to 2 feet from the center) of mounds which measure from 3 to 12 inches in diameter. Finish the treatment then by applying 1 ounce sprinkled evenly over the center of the mound. After application of the granules, it is recommended that the treated area be gently sprinkled with a minimum of 1 quart of water per 6 inches of mound diameter. For best results, do not disturb the mound before or during the application; apply in cool weather (65-80°F), or in early morning or late evening. Apply the granules when Ants have become a nuisance and repeat as needed but not more than once every 7 days.

**HOWARD JOHNSON'S ENTERPRISES, INC.
MILWAUKEE, WI 53201**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. May be injurious to eyes and skin. Do not get on skin or in eyes. Wash thoroughly after handling. Do not wear contaminated clothing. Wash contaminated clothing before reuse. Avoid breathing dust. Keep away from food, feedstuffs, and water supplies.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Store product in its original container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not re-use empty container. Wrap container and put in trash.

PREMISE TREATMENT AROUND THE HOME

To protect the dwelling from possible insect infestations a band of granules may be placed around the building to control Ants, Brown Dog Ticks, Crickets, Earwigs, Fleas, Millipedes, Sowbugs, Springtails and Spiders. One half pound (8 ounces) of this product will treat (100 Sq. Ft.) a 1/2 foot wide band 20 ft. long around the home. This will act as a barrier against insects invading the home. Apply the protective band when insects have become troublesome or before they cause problems in the spring or early fall. Repeat as needed for control. Consult your Agricultural Extension Service for the best time to treat in your area. Keep children and pets off treated areas until grass is dry.

HOW TO APPLY

The following spreader settings are for use with new equipment. Your used spreader may need calibrating. Since proper application is based on the spreaders condition with regard to age and use, your working speed and the evenness of the terrain your spreading equipment may require adjustments.

SPREADER SETTINGS

| SPREADER | 4.5#/1,000 | 6.8#/1,000 | 9.0#/1,000 |
|-------------------|------------|------------|------------|
| PRIZE CB-85 | 6.60 | 6.0-2X | 6.6-2X |
| CBT-85 | 6.60 | 6.0-2X | 6.6-2X |
| SCOTT'S R7X | C | M-2X | 0.2X |
| LESCO B'CAST | C | M-2X | 0-2X |
| CYCLONE | 7.0 | 6.0-2X | 7.0-2X |
| SPYKER | 6.50 | 6.0-2X | 6.5-2X |
| LELY (3 MPH, 20') | 3.50 | 3.0-2X | 3.5-2X |

NOTICE: Buyer and user assumes all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.