

PMJ/32802-42 1856

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460 NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	EPA REGISTRATION NO: 32802-42	DATE OF ISSUANCE SEP 29 1987
	TERM OF ISSUANCE	
	NAME OF PESTICIDE PRODUCT Pro-Tek Lawn Fungicide	

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Howard Johnson's Enterprises, Inc.
 161 West Wisconsin Avenue, Suite 4006
 Milwaukee, WI 53203

ASG
198122/161
17

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment:
 - a. Add the phrase "EPA Registration No. 32802-42."
 - b. The proposed labeling is generally acceptable; however, we suggest that you expand the Spreader Settings Table to include settings for all the label recommended dosage rates. Also change the word "have" to "half" in the sentence below this table.
3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL	DATE
---------------------------------	------

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Lois A. Rossi
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (TS-767C)

Enclosures



18 Pounds Treats Up To
6,000 Sq. Ft. of Lawn

PRO-TEK™ LAWN FUNGICIDE

LAWN DISEASE CONTROL FOR BLUEGRASS, FESCUE,
RYEGRASS, BERMUDAGRASS, ZOYSIAGRASS, AND ST. AUGUSTINE
ALSO FOR DISEASE CONTROL ON BENTGRASS

FOR THE PREVENTION OR CURE OF DOLLAR SPOT, COPPER SPOT, POWDERY MILDEW,
RED THREAD, RUSTS, BROWN PATCH, ANTHRACNOSE, POA ANNUA
DECLINE, STRIPED SMUT, FUSARIUM BLIGHT, GRAY SNOW MOLD,
PINK SNOW MOLD, FUSARIUM PATCH AND SOUTHERN BLIGHT

ACTIVE INGREDIENT:

1-(4-chlorophenoxy)-3,3-dimethyl-1-(1H-
1,2,4-triazol-1-yl)-2-butanone.

INERT INGREDIENTS:
Total

ACCEPTED
COMMENTS
to EPA Letter Dated:

SEP 29 1987
99.5% the Federal Insecticide,
100% the Federal Fungicide and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
32802-42
Kodak Corp.

Contains BAYLETON® Turf Fungicide a Registered Trade Mark of Kodak Corp.

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Give water until vomitus is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
IF IN EYES: Flush eyes with plenty of water. Get medical attention.
IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Get medical attention. Wash clothing before re-use.
IF INHALED: Remove victim to fresh air. Administer artificial respiration if indicated.

EPA REG. NO. 32802-___

EPA EST. NO. 32802-WI-1.
32802-MO-1.

NET WEIGHT 18 POUNDS

HOWARD JOHNSON'S ENTERPRISES, INC.
RAILROAD AVENUE
VIROQUA, WI 54665

PRO-TEK™ LAWN FUNGICIDE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, **INHALED** or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapor, or dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Wash contaminated clothing with soap and hot water before re-use.

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. Do not use on crops grown for food or forage. Do not apply directly to water. Apply this product only as specified on this label. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE.** Store product in original bag in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not re-use empty bag. Wrap and put in trash.

Bayleton is absorbed rapidly and works systemically from within the plant. Rainfall or sprinkler irrigation soon after application will assure rapid systemic uptake. Control will be less effective on turf suffering from drought stress. Therefore in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices. Apply when the turf is fully established and actively growing. Applications should be applied at recommended intervals to maintain control.

DISEASES AND RATES FOR PREVENTION OR CURE

DISEASE	HOW MUCH TO USE PER 1,000 SQ.FT.		REMARKS
DOLLAR SPOT	PREVENTIVE 3 lbs.	CURATIVE -	PREVENTIVE RATE: Apply recommended rate on 30 day intervals. Protective activity of product may extend for as long as 60 days dependent upon environmental conditions.
	1.5 lbs.	-	PREVENTIVE RATE: Apply recommended rate on 14 day intervals. Protective activity of product may extend for as long as 30 days dependent upon environmental conditions.
	-	6 lbs.	CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventive schedule.

PRO-TEK™ LAWN FUNGICIDE

DISEASES AND RATES FOR PREVENTION OR CONTROL

DISEASE	HOW MUCH TO USE PER 1,000 SQ.FT.	REMARKS
COPPER SPOT POWDERY MILDEW RED THREAD (Laetisaria fuciformis)	PREVENTATIVE CURATIVE 3 lbs. 6 lbs.	PREVENTIVE RATE: Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of this product can be greater than 30 days depending on environmental conditions.
RUSTS BROWN PATCH (Rhizoctonia solani) (Suppression)		CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventive schedule and rate.
ANTHRACNOSE POA ANNUA DECLINE	6 lbs. - 3 lbs. - - 6 lbs.	PREVENTIVE RATE: Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45 days. PREVENTIVE RATE: Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 30 days. CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied on preventive schedule and rate.
STRIPED SMUT	PREVENTIVE RATE ONLY 6 lbs.	Make a single application in spring as grass growth begins.
FUSARIUM BLIGHT (Fusarium spp.)	(Except California) 6 lbs. (California Only) 6 to 12 lbs.	Apply 1st applications in mid-June or 30 days prior to time this blight normally becomes evident. Apply at 30 day intervals. For California Only apply at 6 lbs. rate on 15 day intervals or at the 12 lb. rate on 30 day intervals.
GRAY SNOW MOLD (Typhula incarnata) Except California	12 lbs.	Apply in fall 30 days prior to turf dormancy. If turf breaks dormancy during winter months a second application should be made. Do not apply over snow cover, or when turf is dormant.
PINK SNOW MOLD FUSARIUM PATCH (Fusarium nivale) SOUTHERN BLIGHT (Sclerotium rolfsii) Except California	6 to 12 lbs.	Apply before conditions favorable for infection occur. Repeat as needed at 60 to 90 day intervals. Do not apply over snow cover or when turf is dormant. Use high rate in turf with a history of severe yearly disease damage.
GRAY SNOW MOLD PINK SNOW MOLD California Only	Single Application 15-24 lbs./1,000 sq.ft.	Apply in late fall before snow cover. Do not apply over snow. Use the high rate in areas with a history of severe snow mold turf damage.

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

PRO-TEK™ LAWN FUNGICIDE

LAWN DISEASE CONTROL IS BASED ON PROPER APPLICATION

The following spreader settings are for use with new equipment. Your used spreader may need calibrating. Since proper application is based on the spreaders condition with regard to age and use, your walking speed and the evenness of the terrain, your spreader may require a slight adjustment.

If you have reason to believe that your spreader is not calibrated correctly test apply by placing the amount for 500 Sq.Ft. into your spreader and apply to your pre-measured 500 Sq.Ft. (5 ft. wide by 100 ft long)

HOW TO MAKE SURE YOUR SPREADER IS OPERATING CORRECTLY

Using a Scott Spreader For Disease Control place 1.50 lbs. granules in your spreader using No.4 1/2 setting & apply to 500 Sq.Ft. With a 20 inch spreader this will equal 3 spreader widths wide (3 passes) by 100 feet long. If you run out before you cover the pre-measure area close the spreader setting by 1/2 the number. If you have granules left over, open (increase) the spreader setting by 1/2 the number. For further help refer to your Spreader Guide Book.

POOR DISEASE CONTROL WILL RESULT IF YOUR SPREADER APPLIES LESS GRANULES THAN RECOMMENDED.... MAKE SURE TO APPLY THE TOTAL POUNDS PER SQUARE FEET AS SHOWN IN THE FOLLOWING CHART.

SPREADER SETTINGS

SPREADER	AMOUNT TO APPLY	18 POUNDS	18 POUNDS
		PER 6,000 SQ.FT.	PER 3,000 Sq.Ft.
		3.0 lbs./1,000 Sq.Ft.	6.0 lbs./1,000 Sq.Ft.
Scott _____	18 pounds	# 4 1/2	# 6
Cyclone _____	18 pounds	# 3	# 4

When using a 18 lb. bag; spreader settings will be the same as for the 36 lb. bag but it will treat only have the amount of Sq.Ft. as the 36 lb bag. These settings are based on a "One Time Over" application.

NOTICE: Buyer and user assumes all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.