



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

APR 14 1995

H.R. MCLANE
H.R. MCLANE INC.
AGENT FOR: HOWARD JOHNSON'S ENTERPRISES
7210 RED ROAD SUITE 206
MIAMI, FL. 33143

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 07/01/94 in Response to PR Notice 93-7
EPA Reg. No. 32802-42
PRO-TEK LAWN FUNGICIDE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.

1034



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

PRO-TEK® LAWN FUNGICIDE

LAWN DISEASE CONTROL FOR BLUEGRASS, FESCUE, RYEGRASS, BERMUDAGRASS, ZOYSIAGRASS, AND ST. AUGUSTINE

FOR THE PREVENTION OR CURE OF DOLLAR SPOT, COPPER SPOT, POWDERY MILDEW, RED THREAD, RUSTS, BROWN PATCH, ANTHRACNOSE, POA ANNUA DECLINE, STRIPED SMUT, FUSARIUM BLIGHT, GRAY SNOW MOLD, PINK SNOW MOLD, FUSARIUM PATCH AND SOUTHERN BLIGHT,

ACTIVE INGREDIENT:

1-(4-chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone. 0.50%

INERT INGREDIENTS: 99.50%
Total 100.00%

ACCEPTED
with COMMENTS
In EPA Letter Dated

Contains BAYLETON® Turf Fungicide a Registered Trade Mark of Mobay Corp.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENTS OF PRACTICAL TREATMENT

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

32802-42

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Give water until vomitus is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. IF IN EYES: Flush eyes with plenty of water. Get medical attention. IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Get medical attention. Wash clothing before re-use. IF INHALED: Remove victim to fresh air. Administer artificial respiration if indicated.

EPA REG. NO. 32802-42

EPA EST. NO. 32802-W1-1
32802-M0-1

NET WEIGHT 18 POUNDS

HOWARD JOHNSON'S ENTERPRISES, INC.
MILWAUKEE, WI 53201

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, INHALED or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapor, or dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Wash contaminated clothing with soap and hot water before re-use. ENVIRONMENTAL HAZARDS Do not contaminate water by cleaning of equipment or disposal of wastes. Do not use on crops grown for food or forage. Do not apply directly to water. Apply this product only as specified on this label. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store product in original bag in a cool, dry, locked place out of reach of children. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. HOUSEHOLD: Do not re-use empty bag. Wrap and put in trash.

Bayleton is absorbed rapidly and works systemically from within the plant. Rainfall or sprinkler irrigation soon after application will assure rapid systemic uptake. Control will be less effective on turf suffering from drought stress. Therefore in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices. Apply when the turf is fully established and actively growing. Applications should be applied at recommended intervals to maintain control.

BEST COPY AVAILABLE

DISEASES AND RATES FOR PREVENTION OR CURE

DISEASE	HOW MUCH TO USE PER 1,000 SQ.FT.	REMARKS
	PREVENTIVE 3 lbs.	CURATIVE - PREVENTIVE RATE: Apply recommended rate on 30 day intervals. Protective activity of product may extend for as long as 60 days dependent upon environmental conditions.
DOLLAR SPOT	1.5 lbs.	- PREVENTIVE RATE: Apply recommended rate on 14 day intervals. Protective activity of product may extend for as long as 30 days dependent upon environmental conditions.
	-	CURATIVE RATE: To control existing infections, apply them curative rate. Subsequent applications should be applied on a 6 lbs. preventive schedule.
COPPER SPOT POWDERY MILDEW RED THREAD (Laetisaria fuciformis)	PREVENTATIVE 3 lbs.	CURATIVE 6 lbs. PREVENTIVE RATE: Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of this product can be greater than 30 days depending on environmental conditions.
RUSTS BROWN PATCH (Rhizoctonia solani) (Suppression)		CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventive schedule and rate.
	6 lbs.	- PREVENTIVE RATE: Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45 days.
ANTHRACNOSE	3 lbs.	- PREVENTIVE RATE: Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 30 days.
POA ANNUIA DECLINE	-	6 lbs. CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied on preventive schedule and rate.
STRIPED SMUT	PREVENTIVE RATE ONLY 6 lbs.	Make a single application in spring as grass growth begins.
FUSARIUM BLIGHT (Fusarium spp.)	(Except California) 6 lbs. (California Only) 6 to 12 lbs.	Apply 1st applications in mid-June or 30 days prior to time this blight normally becomes evident. Apply at 30 day intervals. For California Only apply at 6 lbs. rate on 15 day intervals or at the 12 lb. rate on 30 day intervals.
GRAY SNOW MOLD (Typhula incarnata) Except California	12 lbs.	Apply in fall 30 days prior to turf dormancy. If turf breaks dormancy during winter months a second application should be made. Do not apply over snow cover, or when turf is dormant.
PINK SNOW MOLD FUSARIUM PATCH (Fusarium nivale)	6 to 12 lbs.	Apply before conditions favorable for infection occur. Repeat as needed at 60 to 90 day intervals. Do not apply over snow cover or when turf is dormant. Use high rate in turf with a history of
SOUTHERN BLIGHT (Sclerotium rolfsii) Except California		severe yearly disease damage.
GRAY SNOW MOLD PINK SNOW MOLD California Only	Single Application 15-24 lbs/1,000 sq.ft.	Apply in late fall before snow cover. Do not apply over snow. Use the high rate in areas with a history of severe snow mold turf damage.

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

BEST COPY AVAILABLE