(Front Panel)

ALL SEASON

SPRAYABLE WEED & FEED

Use only with the hose-end sprayer supplied. For use on home lawns.

This product will effectively control dandelions, chickweed, clover, ground ivy, oxalis, plantain, and over forty other common broadleaf weeds,

Triamine - a trademark of riverdale chemical company

Patent Applied For.

COVERAGE: Northern Turf - 5,000 Square Feet

Southern Turf - 7,500 Square Feet

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Other Panels For Additional Precautionary Statements

EPA REG. NO. 32802-

EPA EST. NO. 228-IL-1

NET CONTENTS 64 FLUID OUNCES

15% Urea Nitrogen

NET WEIGHT 4.5 POUNDS

Manufactured by:

HOWARD JOHNSON'S ENTERPRISES, INC. Milwaukee, WI 53201

ACCEPTED

JUN 2 3 1987

Under the Federal Insecticide, Fungliside, and Redenticide Action on amounted, for the position registered under EPA Reg. No. 3 2502-45

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

ICAUTION: Harmful if absorbed through skin. Wash thoroughly lwith soap and water after handling. IF SWALLOWED: Get medical lattention. IF IN EYES: Immediately wash eyes with blenty of lwater. IF ON SKIN: Wash with plenty of water. If irritation ldevelops, get medical attention.

ENVIRONMENTAL HAZARDS

1Do not apply directly to water or wetlands. Do not contaminatel lwater by cleaning of equipment or disposal of wastes. Do not ! lapply when weather or conditions favor drift from target areas.!

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

WHEN TO USE: Spray anytime during the growing season when weeds are actively growing, except as noted for Southern grasses. If lawn is extremely dry, water thoroughly 1 or 2 days before applying. Do not apply when temperatures are expected to exceed 85 degrees F or if rain is expected within 24 hours of application. Do not apply to new lawns until new turf has been mowed at least three times. Do not reseed until three weeks after application.

Covers 5,000 square feet of turf when applied to bahia, bermuda, bluegrass, fescue, rye, and zoysia grasses.

Covers 7,500 square feet of turf when applied to St. Augustine, centipede, and bent grasses. Apply only in the spring and fall, and only when these grasses are actively growing. Some temporary damage may occur at this time.

DO NOT exceed recommended rate. Failure to comply with this may result in damage to the lawn.

OO NOT apply to Dichondra, Lippia or Clover. Application to any of these will cause lawn damage.

NOT use this product to control weeds in or near flower and vegetable gardens, shrubs, or ornamental plants.

DO NOT mow or water the lawn for at least 24 hours after in application in order for the weed killer to penetrate the weed leaves.

Revised 6/5/87

WEEDS CONTROLLED

(::

Black Medic
Buckhorn
Buttercup
Canada Thistle
Carpet Weeds
Chickweed
Chicory
Clover
Cocklebur
Dandelion
Dog Fennel
English Daisy
Florida Pursley
Frenchweed

Heartloaf Drymary

Ground Ivv

Heal-All

Henbit

Jimson Weed Knotweed Lambsquarter Mallow Morning Glory Mustard Oxalis Pennywort Pepperweed Pia Weed Plantain Poison lvv Poison Oak Purslane Rao Weed Red Clover

Red Sorrell Sheep Sorrell Speedwell Spurge Spur Weed Stitchwort Toadflax Wild Aster Wild Carrot Wild Garlic Wild Geranium Wild Lettuce Wild Onion Wild Radish Wood Sorrell Yarrow, and Other Broadleaf Weeds

After applying this Weed & Feed product, some weeds will begin to show curling and other visible signs of herbicide activity within two or three days. In the case of some hard to kill broadleaf weeds, a second application of this product, three to four weeks after the original application, may be necessary.

HOW TO USE AND APPLY

- (1) Shake container to ensure that all materials are dissolved.
- (2) Make sure the valve on the container is in the "OFF" position.
- (3) Connect the spray nozzle to the hose end.
- (4) Remove the container cap and screw the container to the spray nozzle.
- (5) Turn on water at the tap.
- (6) Turn the valve on the nozzle to the "UN" cosition and begin a sweeping motion of the spray on the lawn. You may wish to practice this motion before you actually start applying the Weed and Feed (by placing nozzle on hose without Weed & Feed' container and spraying plain water).
- (7) Begin application at the point farthest from the hose shut- off and work backwards toward the shut-off. Overlap slightly as you spray the lawn, keeping the spray pointed slightly (, downward.

Easy to use.

No premixing necessary.

Just fasten to your hose end and follow the easy to understand instructions.

STORAGE AND DISPOSAL

ISTORAGE: Keep from freezing. To be stored in original containeri land placed in a locked area inaccessible to children. Do not 1 Istore near food or feedstuff. PSSTICIDE DISPOSAL: Product remaining in original container should be disposed Isecurely wrapping original contaner in several layers of Inewspaper and discarding in trash. CONTAINER DISPOSAL: Do not (Ireuse container. Rinse thoroughly before discarding in trash. I

NOTICE: Manufacturer warrants that this product conforms to all label guarantees and will perform as indicated when used as No claim for or uses of this product that are not on directed. the label will be honored by the manuacturer.

MARKS ON THE CONTAINER WILL GIVE APPROXIMATE SQUARE FOOT COVERAGE YOU SHOULD HAVE TREATED WHEN AT THE GIVEN REMAINING LIGUID LEVEL.

SAMPLE OF MEASURING GUIDE (Not to Scale)

NORTHERN	i Ii	JRE
1000	ទួច	EI

SOUTHERN TURE

<u>1875 SQ FT</u>

2000 SQ FI

3750 50 ET


3000 SQ FT

4525 SQ EI


4000 SQ FT

5000 SQ FI


7500 SO ET



Connect spray notifie to hose,



Remove bottle cap and screw bottle into rottle.



Turn the payod lever on top of the nozzle to the OFF position. Turn on the water at the tap. When ready to treat, turn the onlot lever to its ON notion. The Usuki Weed and Food will mix automorally at the



Slowly walk back and forth at a steady pace. Overlap slightly for uniform coverage, Maria on right side panel indicate square fool coverage.

Revised 6/5/87