DURSBAN® 100 GRANULE Insecticide
LAWN AND TURF INSECT CONTROL

KILLS SOD WEBWORMS, WHITE GRUBS, ANTS, FLEAS, BROWN DOG TICKS, CRICKETS, CHINCH BUGS, CUTWORMS, EARWIGS, SOWBUGS, AND ARMYWORMS, EUROPEAN CRANE FLY LARVAE, HYPERODES (TURFGRASS) WEEVIL, IMPORTED FIRE ANT

IN HOME LAWNS. ORNAMENTAL & RECREATIONAL TURF

Total

DURSBAN® a Registered Trademark of The DowElanco

### KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT
LOWED: Call a physician or Poison Control Center immediately. Induce vomiting by clim 1 or 2 glasses of water and touching back of throat with finger. Do not induce

vorming or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention.

IF ON SKIN: Remove contaminated clothing and wash affected area with soep & warm water.

Wash clothing before reuse IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated. Get medical

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated. Get medical attention immediately.

NOTE TO PHYSICIAN: Chlorpyrifox is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an artifote.

SEE SIDEBACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. May be injurious to eyes and skin. Do not get on skin for in eyes. Wash thoroughly after handling. Do not wear contaminated clothing. Wash contaminated clothing before reuse. Avoid breathing dust. Keep sway from food, feedstuffs, and water supplies. Flush contaminated eyes with plenty of water and get medical attention. As directed, water lawn after application and allow to dry before permitting children or pets on treated areas. treated areas.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment of disposal of equipment washwaters. Apply this product only as specified on this label.

DIRECTIONS FOR USE violation of Federal law to use this product in a manner inconsistent with its labeling. luse on turf grown for sale or other commercial use as sod, or for commercial seed ction, or for research purposes.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. <u>STORAGE</u>: Store product in its original container in a cool, dry, locked place cut of reach of children. <u>PESTICIDE DISPOSAL</u>: Wastes resulting from the use of this product may be disposed of on site or at an approved weste disposel facility. <u>CONTAINER DISPOSAL</u>: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. <u>HOUSEHOLD</u>: Do not reuse empty container. Wrap container and put in tresh.

LAWNS AND ORNAMENTAL TURF SURFACE FEEDING INSECTS
For control of Chinch Bugs, Sod Webworms (Lawn Moth), Ants, Fleas, Armyworms, Brown Dog Ticks, Crickets, Cutworms, Earwigs, Sowbugs and Millipedes apply 50 lb. to treat 22,000 sq.ft. Refer to Spreader Settings. Treat in the spring or when insects first appear troublesome. Additional treatments at 30 day intervals may be medically assisting insects.

Additional treatments at 30 day intervals may be medically assisting insects.

Manufactured By

SEP 2 0 1994

Under the Federal Insecticide. Pungleide, and Rodenticlide Act. on amended, for the pesticide replained on the pesticide PARSE, No.

#### FOR ROOT FEEDING INSECTS

For Billbug Larvae, and the "White Grub" larvae of Japanese Beetle, European Chater, Southern Chater and Hyperodes Weevil apply 50 lbs. to treat 11,000 sq.ft. of lawn. European Crane Fly larvae feeding in western states turigrass will be controlled using 50 lbs per 15,000 sq.ft with an individual application or if present, simultaneously with the same application being applied for control of "White Grubs" i.e., (50 lbs/11,000 sq.ft.). Consult your Agricultural Extension Agent for proper time

to apply.

For the <u>Hyperodes (Turfgrass) Weevil</u> apply in Mid-April and again in Mid-May using 50 lbs. per 11,000 sq.ft. Refer to the Spreader Settings

For soil insects apply uniformly over the area and follow with a thorough watering to wash the insecticide into the soil where insects are feeding. For best control apply when the "grubs" are young and actively feeding near the soil surface in mid-July and early August.

#### INSECT CONTROL AROUND THE HOME

To protect the dwelling and other buildings from possible insect infestations a band of granules may be placed around the building to control Ants, Brown Dog Ticks, Crickets, Earwigs, Fleas, Millipedes, Sowbugs, Springtails and Spiders. One half pound (8 ounces) of this product will treat (200 Sq. Ft.) a 5 foot wide band 40 feet long around the structure. This will act as a barrier against insects invading the home. Apply the protective band when insects have become troublesome or before they cause problem3 in the spring or early fall. Repeat one month later and then as needed for control but not more than once every 7 days. Consult your Agricultural Extension Service for the best time to treat in your area. your Agricultural Extension Service for the best time to treat in your area.

#### IMPORTED FIRE ANT

AS AN AID IN THE CONTROL AND REDUCTION OF FIRE ANTS AS AN AID IN THE CONTROL AND REDUCTION OF FIRE ANTS As an aid in the control and reduction of Imported Fire Ants treat individual Ant mounds measuring from 3 to 12 inches in diameter by applying 4 level tablespoonfuls (2/3 ounce) of granules evenly in a circle around the outer edges (3 to 4 feet from the center) of the mound. Finish the treatment then by applying 2 level tablespoonfuls (1/3 ounce) sprinkled evenly over the center of the mound. Best results will be obtained if mound is not disturbed after treatment thus allowing ants to come and go and to be exposed to the insecticide treated area. Repeat treatment as needed for additional control but not more than once every 7 days. 7 days.

KNOW YOUR FERTILIZER SPREADER

The following spreader settings are offered for use with new equipment. Your used spreader may need calibrating. Since proper application is based on the spreaders condition with regard to age and use, your walking speed and the evenness of the terrain, your spreading equipment may require adjustments from the following figures. If you have reason to believe that your spreader is not calibrated correctly (est apply by placing the amount for 500 sq.ft. into your spreader and apply to your premeasured 500 sq.ft. (5 ft. wide by 100 ft. long): Faulty and/or erratic application will result in poor insect control!

-			PREADER SETTINGS	
	SPREADER	AMOUNT TO	SOD WEBWORMS & OTHER SURFACE FEEDING INSECTS	VI HITE GRUBS AND
	<u> </u>	APPLY.	50 lbs /22,000 Sq.Ft.	50 lbs /11,000 Sq. Ft.
	Scott	50 pounds	2 1/4 lbs /1,000 Sq.Ft. # 4 1/2	4 1/2 lbs/1000 Sq. Ft. # 6
	Cyclone	50 pounds	# 4	#5

For control of Crane Fly Larvae use Scott Spreader setting of #5 and Cyclone Spreader Setting of #4 1/2.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label. (49-WPS.PM5 \ 07-15-94)

EPA EST. NO. 32802-WI-1; MO-1

NET WEIGHT 50 POUNDS

Milwaukee, WI 5320 HOWARD JOHNSON'S ENTERPRISES, INC. P.O. BOX 2990 - MILWAUKEE WI 53201 BEST AVAILABLE COPY

# WORKER PROTECTION STANDARD CERTIFICATION STATEMENT FOR

# DURSBAN® 100 GRANULES INSECTICIDE

## EPA REG. NO. 32802-49 HOWARD JOHNSON'S ENT. INC. MILWAUKEE, WI 53201

(Addition of a WPS Exclusionary Statement to labeling)
Certification when deviations are proposed:

I certify that the revised labeling being submitted for this product is in complete accordance with the labeling requirements of PR Notice 93-7, which reflects the requirements of EPA's labeling regulations for worker protection statements (40 CFR part 156, subpart K) except where noted. I have, in one or more places in the revised labeling, proposed changes that deviate from the exact language or location specified in the PR Notice. I have identified all such statements by highlighting them in a way that distinguishes them from any other label revisions. I have attached an explanation for each proposed deviation. I have proposed no changes to this labeling other than those required to comply with 40 CFR part 156, subpart K. I understand that EPA cannot guarantee to render its decision on revised labeling not in complete conformance with PR Notice 93-3 by April 21, 1994. I understand that it would be a violation of FIFRA if I or my supplemental registrants were to sell or distribute this product after April 21, 1994, without amended labeling complying with the requirements of 40 CFR part 156, Subpart K.

Signed:	Date _	
H. R. McLane, Inc. By H.	R. McLane, Authorized Agent	
The Acute toxicity categor	ries for this product as entered in bloc	ks 3 through
7 of the Product Workshe	eet are as follows:	•

)

Toxicity (	Category
Acute oral	Ш
Acute dermal	111
Skin irritation potentia	al III
Eye irritation potential	
Acute inhalation	IV

BEST AVAILABLE COPY