

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460	EPA REGISTRATION NO. 32802-65	DATE OF ISSUANCE 01 OCT 1992
	TERM OF ISSUANCE Conditional, Expires: January 1, 1995	
	NAME OF PESTICIDE PRODUCT Craggrass Preventer Plus with 0.20 Barricade™ preemergence herbicide	

NOTICE OF PESTICIDE:  REGISTRATION  
 REREGISTRATION  
 (Under the Federal Insecticide, Fungicide,  
 and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Howard Johnson's Enterprises, Inc.  
 P.O. Box 2990  
 Milwaukee, WI 53201

**BEST AVAILABLE COPY**

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(C) provided that you:

1. Submit nitrosamine data to fulfill the regulations requiring a discussion of the formation of impurities (Code of Federal Regulations, 40: section 156.167(b)) within one year of date of this notice.
2. Submit storage stability data for product within 16 months.
3. Add the phrase "EPA Registration No. 32802-65" to the label before you release the product for shipment.
4. Submit five (5) copies of the final printed labeling before you release the product for shipment.

and provided that Sangoz Agro, Inc., the manufacturer of the active ingredient of this product, submits the required data to continue the registration of proflamline as listed in the EPA fact sheet no. 231.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Joanne I. Miller  
 Product Manager (23)  
 Fungicide-Herbicide Branch  
 Registration Division (75-7505C)

ENCLOSURE

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

*Joanne I. Miller*

DATE

9/30/92

**BEST AVAILABLE COPY**

# Crabgrass Preventer Plus 0.20 Barricade®

## Preemergence Herbicide

### For Selective Preemergence Control of Grass and Broadleaf Weeds in Established Turf Grasses and Landscape Ornamentals

#### ACTIVE INGREDIENT:

Proflumarine (N,N-Di-n-propyl-2,4-dinitro-6-trifluoromethyl-m-phenylenediamine) - 0.20%  
 Inert Ingredients - 99.80%  
 Total 100.00%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See Side Panel for Additional Precautionary Statements  
 Barricade® is a registered trademark of Sandaz, Ltd.  
 EPA Reg. No. 32802-AL EPA Est. No. 32802-WI-1

#### PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION:** Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

**Statements of Practical Treatment**  
**If in Eyes:** Flush thoroughly with water for several minutes. Get medical attention if irritation persists.  
**on Skin:** Wash with soap and water. Rinse thoroughly. Get medical attention if irritation persists.  
**If Inhaled:** Remove victim to fresh air.

#### ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in original container away from other fertilizer, feed and foodstuffs, and separated from other pesticides.

#### PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Do not reuse empty bag. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or at an incineration facility, or as directed by State or local authorities, by burning. If burned stay out of smoke.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is a turf fertilizer and a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in turf grasses and landscapes. It controls susceptible weeds by inhibiting weed seeds germination and root development. For most effective weed control it must be incorporated by 1/2 inch of rainfall, irrigation, or shallow mechanical incorporation. Incorporation should take place as soon as possible after application and should not be delayed past 14 days from the date of application. Do not apply this product through any type of irrigation system. Do not apply heavily.

#### APPLICATION DIRECTIONS

Apply uniformly through suitable, calibrated commercial applicator.

#### ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established grass turf. The species of turf grass determines the maximum amount of material that may be applied during a year. Most effective weed control in turf grasses will be obtained when this product is activated by at least 1/2 inch of rainfall or irrigation within 14 days after application and prior to weed germination. See the map below for approximate crabgrass germination dates.

#### CRABGRASS SEED GERMINATION DATES



#### Approximate Date

- Zone 1 After May 30
- Zone 2 After May 10
- Zone 3 After April 20
- Zone 4 After March 20
- Zone 5 Jan 1 to Mar 20

Footfall (yellow-green) tend to germinate before crabgrass. Therefore, if targeting footfall, apply 3 weeks before the corresponding crabgrass germination date.

The following use precautions apply to this product in turf grasses. In turf grasses, application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to **FALL SEEDED TURF** until the following spring and only after the second mowing. Do not apply to **SPRING SEEDED TURF** or to newly set sod until the following year. Application to turf stressed by drought, low fertility, or poor drainage may result in turf injury. Do not apply this product to trees or green, or to areas where dichondra, colonial bermudagrass, velvet bentgrass or annual bluegrass (Poa annua) are desirable species.

Maximum Annual Rates  
 This product should not be applied at a maximum annual rate of 750 lbs. per acre

#### TURF SPECIES AND RATES

For season long control this product may be applied as a single application, or if needed followed by a second application for increased length of control. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rate when the listed range for longer control. Longer control is desirable in areas where crabgrass germinates between January 1 and April 20. (See Crabgrass Seed Germination Date Map.)

Howard Johnson's Enterprises, Inc.  
 P.O. Box 2990 - Milwaukee, WI 53201 U.S.A.

#### TURF SPECIES AND RATES (Cont.)

Select maximum use rate based on turf grass species. All applications must be made prior to germination of the target weed seeds. The rates listed in the table below are for weed control of approximately 2 - 6 months.

#### Turf Species and Rates Table

Turf Species	Minimum & Maximum Rate Lbs / Acre/ Calendar Year
Bermudagrass	160 to 750 lbs.
Centipedegrass	"
Seashore Paspalum	"
All Fescue	"
Zoysia	"
Perennial Bluegrass	160 to 500 lbs.
Perennial Ryegrass	"
Creeping Red Fescue	160 to 370 lbs.
St. Augustinegrass	"
Creeping Bentgrass min 1/2"	160 to 330 lbs.
Bermudagrass; newly sprigged or plugged <sup>(1)</sup>	160 to 280 lbs.

<sup>(1)</sup> Suppression only of Foxtail, Goosegrass & Rescuegrass due to reduced product rates. Do not apply more than 375 lbs. of product per application. Do not make more than 2 applications per calendar year. Do not apply more than 750 lbs. of product per calendar year. If making two applications, wait at least 60 days after the first application before making a second application. Use higher rates of this product to achieve higher levels of fertility and weed control for each turf type.

#### WEEDS CONTROLLED

When used at the recommended rates (See Turf Species and Rates Table) this product will control:

- |                             |                  |
|-----------------------------|------------------|
| Annual Bluegrass            | Knotted          |
| Common Chickweed (1)        | Oxalis           |
| Crabgrass - (Large, Smooth) | Prostrate Spurge |
| Foxtail - (Yellow, Green)   | Rescuegrass (3)  |
| Goosegrass (2)              | Sheepspurge (1)  |
| Henbit                      |                  |

(1) Fall application. (2) For best control of goosegrass apply 375 lbs. / acre. Sixty days later make a second application with 125 to 375 lbs. / acre of product. (see information in Turf Species and Rates section above). (3) Suppression Only.

#### OVERSEEDING/RESEEDING

This product will inhibit germination of turf species if overseeded / reseeded too soon after application. The interval between application and overseeding/reseeding is dependent on the application rate, soil temperature, and the soil moisture. (cont. on next panel)

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

#### GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	32%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	4%
Soluble Potash (K <sub>2</sub> O)	6%
Ingredients: Urea formaldehyde, Urea, Sulfur Coated Urea, Ammoniated Phosphate, Murate of Potash	

#### SPREADER SETTINGS

Spreader	Apply 200 lbs. per 1,000 sq. ft.	Apply 300 lbs. per 1,000 sq. ft.	Apply 400 lbs. per 1,000 sq. ft.	Apply 500 lbs. per 1,000 sq. ft.
Cyclone B-1	28	30	34	41
Ohio Drop	35	40	425	575
Scotts Drop	35	55	625	70

#### Net Weight 20 Pounds

#### Overseeding/Reseeding (Cont.)

A shorter interval is possible if growing conditions are warm and wet, while a longer interval is needed if conditions are cool and dry. Use the following Overseeding/Reseeding Intervals Table as a guide when choosing the appropriate overseeding/reseeding interval for each situation. The table below lists the Overseeding/Reseeding Intervals (Months) Based Upon Seeding Method and Predominant Soil Conditions for the Season.

Lbs./Acre of product	Broadcast Seeding		Drill Seeding	
	wet & warm	cool & dry	wet & warm	cool & dry
160 to 250	4 mos.	6 mos.	3 mos.	4 mos.
251 to 500	6 mos.	8 mos.	6 mos.	8 mos.
501 to 750	12 mos.	18 mos.	9 mos.	12 mos.

#### LANDSCAPE ORNAMENTALS

This product will not harm most established ornamental trees, shrubs and flowers when applied in accordance with the directions on the label. However, it may harm some ornamental plants in the landscape. The following species have shown tolerance to this product.

- |                     |                   |
|---------------------|-------------------|
| Acer palmatum       | Japanese Maple    |
| Acer rubrum         | Norway Maple      |
| Actinidia chinensis | Kiwi              |
| Agave attenuata     | Agave             |
| Agave schottlandii  | Yucca             |
| Agave vivipara      | Yucca             |
| Albizia julibrissin | Silk Tree         |
| Berberis aquifolium | Japanese Barberry |
| Berberis thunbergii | Mountain Barberry |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |
| Berberis zeylanica  | Japanese Barberry |
| Berberis thunbergii | Warty Barberry    |
| Berberis vulgaris   | Japanese Barberry |
| Berberis zeylanica  | Warty Barberry    |
| Berberis thunbergii | Japanese Barberry |
| Berberis vulgaris   | Warty Barberry    |

3 73

ACCEPTED  
with COMMENTS  
EPA Letter 10/1/02

NOV 1 1902

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, this pesticide registered under EPA Reg. No.

32803-65

32803-65