

PM 23 32802-65 10-22-97 PS 186

Accept. Stamp on p. 676

OCT 22 1997

Mr. Thomas J. Schmit  
Howard Johnson's Enterprises, Inc.  
720 W. Virginia St., 50. 222  
Milwaukee, WI 53204

Dear Mr. Schmit:

Subject: Crabgrass Preventer Plus with 4.20 Barricade®  
Preemergence Herbicide  
EPA Registration No. 32802-65  
Crabgrass Preventer Plus with 0.37 Barricade®  
Preemergence Herbicide  
EPA Registration No. 32802-66  
Crabgrass Preventer Plus with 0.55 Barricade®  
Preemergence Herbicide  
EPA Registration No. 32802-67  
Crabgrass Preventer Plus with 0.25 Barricade®  
Preemergence Herbicide  
EPA Registration No. 32802-70

Application and Letter Dated June 24, 1997, Request to Amend Labeling: Updating Label to Match that Accepted for Novartis® Barricade® P Herbicide and Revising Size of Retail Container and Type of Containers

Your request to amend the label of the subject pesticide product registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, to reflect the subject amendments has been reviewed and is acceptable under the Act, provided that you:

1. Correct the address shown on all labels to reflect your address of record with this Agency.
2. Correct the statement on the front panel that reads: "BARRICADE® is a Registered Trademark of Sandoz, Inc." to read "BARRICADE® is a Registered Trademark of Novartis Crop Protection, Inc."

266

3. On labels for the 0.37% and 0.58% active ingredient:

Under "Directions for Use" section correct the statement that reads: "Most effective weed control will be ..... by at least 0.5 inch of rainfall or irrigation, or with shallow (12 inches) incorporation, prior to weed seed germination and within 14 days following application." to read: "Most..... shallow (1 to 2 inches) .....application."

Certainly, you do not want 12 inches of shallow incorporation.

4. Correct footnote 1, for the "Weed Controlled" section to reflect the correct dosages. In case of the 0.37% product the dosage should be 175 to 405 lbs/A.

5. On label for the 0.37% and 0.58% active ingredient:

In the list of "Tolerant Ornamental Species" Geranium cinereum ... Cranesbill was omitted or was blurred on the proposed labeling. Either add this species to the list or omit it; if you wish to delete it, resubmit labeling reflecting your choice.

- 6 Submit one (1) copy of your final printed labeling for each of these products before you release it for shipment under the proposed label, reflecting the above comments.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Stamped copies of labels are enclosed for your records.

Sincerely yours,

Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505C)

SYMBOL	Enclosures	CONCURRENCES				
SURNAME	Wilson	Diskette	prodimine	100p	10-22-97	
DATE						

3046

# CRABGRASS PREVENTER PLUS WITH .20 BARRICADE® PREEMERGENCE HERBICIDE

FOR SELECTIVE PREEMERGENCE CONTROL OF GRASS AND BROADLEAF WEEDS IN LAWNS AND ORNAMENTAL TURF (excluding golf course putting greens), LANDSCAPE ORNAMENTALS, ESTABLISHED PERENNIALS AND WILDFLOWER PLANTINGS

### ACTIVE INGREDIENT:

Prodiamine: [N<sup>3</sup>,N<sup>3</sup>-Di-*n*-propyl-2,4-dinitro-6-(trifluoromethyl)-*m*-phenylenediamine] .....0.20%  
INERT INGREDIENTS: ..... 99.80%  
TOTAL: ..... 100.00%

EPA Reg. No. 32802-65  
EPA Est. No. 32802-WI-1 (A), 32802-MO-1 (B)

First letter of bag code shows establishment where manufactured.  
See tag attached to bag for fertilizer guaranteed analysis.  
Manufactured by:

**HOWARD JOHNSON'S ENTERPRISES, INC.**  
700 W. Virginia Street, Milwaukee, WI 53204-1548  
BARRICADE® is a registered trademark of Novartis Crop Protection, Inc.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION:** See back panel for further precautions and statement of practical treatment.


**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION:** Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.  
**STATEMENT OF PRACTICAL TREATMENT**  
**IF IN EYES:** Flush thoroughly with water for several minutes. Get medical attention if irritation persists.  
**IF ON SKIN:** Wash with soap and water. Rinse thoroughly. Get medical attention if irritation persists.  
**IF INHALED:** Remove person to fresh air.  
**ENVIRONMENTAL HAZARDS:** This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.  
*(Alternate wording for WPS exclusionary statement, for use when the product labeling contains only turf uses (ie when label is split to exclude the directions for use on ornamental sites:)*  
Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds germination and root development. When used according to directions, it will control: barnyardgrass; bluegrass, annual (*Poa annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass<sup>1</sup>; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; Kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass<sup>3</sup>; shepard's purse<sup>2</sup>; signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

<sup>1</sup> In many areas a single application of 7.6 to 17.2 pounds/1000 square feet (330 to 750 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 7.6 to 11.5 pounds / 1000 square feet (330 to 500 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).  
<sup>2</sup> For this weed, apply in late summer, fall or winter prior to germination.  
<sup>3</sup> Suppression only.

**WHEN TO APPLY TO ESTABLISHED TURF:** This product must be applied prior to germination of weed seeds, as it will not control established weeds.



THIS MAP SHOWS APPROXIMATE CRABGRASS GERMINATION DATES

- After May 30
- After May 10
- After April 30
- After March 20
- January 1 to March 20

**APPLICATION DIRECTIONS:** Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days after application.

**USE PRECAUTIONS:** The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil). Do not apply to newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

**RATES OF APPLICATION:** This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed.

*(Label for EPA Reg. No 32802-65, PAGE 1 OF 4)*  
91747

4 of 6

Turf Species	APPLICATION RATES (by Turf Grass Species)	
	Minimum Rate	Maximum Annual Rate <sup>1</sup>
Bahiagrass, Bermudagrass <sup>2</sup> Centipede grass, Zoysiagrass Seasonal Paspalum St. Augustinegrass Tall Fescue (including turf-type)	4 lbs / 1000 sq ft (174 lbs / acre)	17.2 lbs / 1000 sq ft per year (750 lbs / acre per year)
Buffalograss Kentucky Bluegrass Perennial Ryegrass	4 lbs / 1000 sq ft (174 lbs / acre)	11.5 lbs / 1000 sq ft per year (500 lbs / acre per year)
Creeping Red Fescue	4 lbs / 1000 sq ft (174 lbs / acre)	8.5 lbs / 1000 sq ft per year (370 lbs / acre per year)
Creeping Bentgrass	4 lbs / 1000 sq ft (174 lbs / acre)	7.6 lbs / 1000 sq ft per year (330 lbs / acre per year)

<sup>1</sup>These are the maximum rates per calendar year.  
<sup>2</sup>May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 6 pounds/1000 square feet (260 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

**WHEN TO OVERSEED AFTER APPLICATION:** This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/ reseeding results.

USE RATE	MONTHS BEFORE OVERSEEDING		
	North	Transition	South
4 pounds / 1000 sq ft (174 pounds / acre)	4	4	4
7.6 pounds / 1000 sq ft (330 pounds / acre)	5	4	4
8.5 pounds / 1000 sq ft (370 pounds / acre)	5	5	5
9.2 pounds / 1000 sq ft (400 pounds / acre)	—	6	6
11.5 pounds / 1000 sq ft (500 pounds / acre)	—	7	7
13 pounds / 1000 sq ft (570 pounds / acre)	—	—	9
14.9 pounds / 1000 sq ft (650 pounds / acre)	—	—	10
17.2 pounds / 1000 sq ft (750 pounds / acre)	—	—	12

SPREADER SETTINGS	to apply at the listed pounds per 1000 square feet					
	4 lbs	6 lbs	7.6 lbs	8.5 lbs	11.5 lbs	17.2 lbs
Spreader A	X	X	X	X	X	X

(May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product)

**LANDSCAPE ORNAMENTALS** (including established perennials and wildflower plantings): This product may be applied for residual preemergence weed control in ornamentals. Apply at the rate of 7.6 to 17.2 pounds/1000 square feet (330 to 750 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 17.2 pounds/1000 square feet (750 pounds/acre) per year.

**APPLICATION TIMING AND INFORMATION:** This product may be applied to newly transplanted and established ornamentals a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch)

mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

**TOLERANT ORNAMENTAL SPECIES:** This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

SCIENTIFIC NAME	COMMON NAME
Abies spp.	Fir species** (Balsam, Frasier, Noble, etc.)
Abelia grandiflora	Abelia: Sherwood
Acer palmatum	Japanese Maple
A. platanoides	Norway Maple
Achillea sp.	Yarrow: King Edward
Actinidia chinensis	Kiwi*
Agapanthus africanus	Lily-of-the-Nile (African Lily)
Agapanthus orientalis	
Akebia quintata	Five-Leaf or Chocolate Vine
Allium cernuum	Lady's Leek, Nodding Onion
Anemonehydrida	Japanese Anemone
Aquilegia sp.	Aquilegia: Red and Gold
Arctostaphylos densiflora	Vine Hill Manzanita
Arctotheca calendula	Cape Weed
Artemisia sp.	Wormwood; Silver Mound, Castle
Aster sp.	Aster: Bonnie Blue, Purple Dome
Aster X Frikartii	
Athrium Filix-femina	Lady Fern; Fern Lady
Aucuba japonica	Japanese Aucuba
Begonia sp.	Fibrous Begonia: Hardy Grandis
Berberis gladwynensis	Barberry
B. Julianae	Wintergreen Barberry
B. mentorensis	Mentor Barberry
B. Thunbergii	Japanese Barberry
B. verruculosa	Warty Barberry
Bergenia cordifolia	
Boltonia asteroides	Snowbank
Buddleia davidii	Butterfly-Bush (Dwarf Blue); Royal Red
Buxus microphylla	Japanese Boxwood
Callistemon viminalis	Weeping Bottlebrush
Calluna vulgaris	Scotch Heather
Campanula carpatica	Tussock Bellflower, (White Clips)
Campis X Tagliabuana	Trumpet Flower, Madame Galen
Carpobrotus edulis	Hottentot Fig (Ice Plant)
Cassia artemisoides	Feathery Cassia
Ceanothus rigidus	Wild Lilac
Ceratostigma plumbagonoides	
Chamaecyparis pisifera	False Cypress
Cleyera japonica	Cleyera
Citrus spp.	Citrus species*
Coreopsis sp.	Coreopsis (Calliopsis) Early Sunrise, Moonbeam
Cornus florida	Flowering Dogwood
C. stolonifera	American Dogwood
Cortaderia selloana	Pampas Grass
Cotoneaster apiculatus	Cranberry Cotoneaster
C. buxifolius	Cotoneaster
C. dammeri	Bearberry Cotoneaster
C. microphyllus	Rockspray Cotoneaster
Cretaeagus spp.	Hawthorne
Crococsmia sp.	Lucifer
Cupressus sempervirens	Italian Cypress
Delosperma alba	White Trailing Ice Plant
Delosperma sp.	Cooperi Pink
Delphinium sp.	Larkspur: Blue Elf
Dianthus deltoides	Dianthus; Maiden Pinks "Zing"
D. gratianopolitanus	Cheddar Pink
Dodonea viscosa	Hop Bush

Echinacea purpurea ..... Coneflower, Purple; Magnus  
 Elaeagnus pungens ..... Silverberry  
 Euonymus fortunei ..... Wintercreeper  
 E. japonica ..... Evergreen Euonymus  
 E. kiautschovica ..... Spreading Euonymus  
 Fatsia japonica ..... Japanese Aralia  
 Forsythia intermedia ..... Border Forsythia  
 Forsythia suspensa ..... Weeping Forsythia  
 F. viridissima ..... Greenstem Forsythia  
 Gaillardia sp. .... Gaillardia, Blanket Flower: "Goblin"  
 Gardenia jasminoides ..... Gardenia, Cape-Jasmine  
 Gaura sp.  
 Gentiana dahurica ..... Gentian  
 Geranium cinereum ..... Cranesbill  
 Gladiolus spp. .... Gladiolus species\*\*  
 Gypsophila repens ..... Baby's Breath  
 Hedera helix ..... English Ivy  
 Helianthemum sp. .... Sunrose  
 Hemerocallis sp. .... Daylily: Axtec Gold, Stella de Oro, Tender Love  
 Hibiscus ..... Rose of Sharon\*\*  
 Hibiscus sp. .... Mallow: Disco Belle White  
 Hibiscus rosa-sinensis ..... Chinese Hibiscus  
 Hosta plantaginea ..... Hosta, Plantain Lily (Fragrant)  
 H. Sieboldiana ..... Hosta, "Searsucker"  
 Houttuynia cordata var. variegata  
 Hydrangea macrophylla ..... Bigleaf Hydrangea  
 Ilex Bemyi ..... Holly  
 Ilex yuta ..... Chinese Holly  
 I. crenata ..... Japanese Holly  
 I. opaca ..... American Holly  
 I. vomitoria ..... Yaupon Holly, Schillings  
 Inula ensifolia ..... Sword-Leaved Iris; Jodlesong  
 Iris ensata ..... Siberian Iris; Cabernet  
 Iris sibirica ..... Siberian Iris; Cabernet  
 Iris spp. .... Iris species\*\*  
 Jasminium nudiflorum ..... Winter Jasmine  
 Juniperus chinensis ..... Chinese Juniper; Nick's Compact, Parsonii  
 J. conferta ..... Shore Juniper; Blue Pacific  
 J. savinica ..... Parsoni  
 J. horizontalis ..... Creeping Juniper  
 Juglans sp. .... Walnut\*  
 Justicia brandegeana ..... Shrimp Plant  
 Lagerstromia indica ..... Crape Myrtle  
 Lagerstromia indica X Fauriei Crape Myrtle; Tuscarora  
 Lantana montevidensis ..... Weeping Lantana  
 Lavender sp. .... Lavender; Munstead  
 Leonopodium alpinum ..... Edelweiss  
 Ligustrum amurense ..... Amur Privet  
 Ligustrum sinense ..... Chinese Privet; Variegata  
 Ligustrum japonicum ..... Japanese Privet  
 Ligustrum lucidum ..... Glossy Privet (Waxleaf)  
 Lillium sp. .... Lily; Jazz  
 Liriope muscari ..... Big Blue Lily; Silver Mound, Evergreen Giant  
 Liriope muscari var. variegata Liriope, Variegated  
 Liriope spicata ..... Liriope, Creeping  
 Lobelia Cardinalis ..... Cardinal Flower, Indian Pink  
 Lonicera japonica ..... Japanese Honeysuckle  
 L. tatarica ..... Tatarian Honeysuckle  
 Lcrapetalum chinense ..... Burgundy  
 Lythrum sp. .... Loosestrife; Modern Pink  
 Magnolia spp. .... Magnolia species\*\*  
 Maziephora luteola ..... Ice Plant  
 Malus sp. .... Crabapple\*  
 Miscanthus sinensis ..... Yaku Jima, Silberfeder  
 Nandina domestica ..... Heavenly Bamboo  
 Narcissus spp. .... Narcissus species\*\*  
 Nerium oleander ..... Oleander  
 Olea europaea ..... Olive \*  
 Ophiopogon japonicus ..... Mondo Grass  
 Osmanthus heterophyllus ..... Osmanthus (False Holly): Gulf Tide  
 Osteospermum fruticosum ..... Trailing African Daisy  
 Oxydendrum arboreum ..... Sourwood  
 Paeonia suffruticosa ..... Tree Peony  
 Pennisetum setaceum ..... Fountain Grass (Dwarf)  
 Perovskia atriplicifolia  
 Persea americana ..... Avocado\*  
 Photinia Fraseri ..... Fraser's Photinia (Redtip)  
 Phytolagia Virginiana ..... Dragonhead, False; Vivid  
 Picea spp. .... Spruce species\*\* (Colorado Blue, Norway, etc.)  
 Pieris japonica ..... Lily-of-the-Valley Bush

Pinus brutia ..... Calabrian Pine  
 P. canariensis ..... Canary Island Pine  
 P. elliottii ..... Slash Pine  
 P. halepensis ..... Aleppo Pine  
 P. nigra ..... Austrian Pine  
 P. palustris ..... Longleaf Pine  
 P. radiata ..... Monterey Pine  
 P. strobus ..... Eastern White Pine  
 P. sylvestris ..... Scotch Pine  
 P. taeda ..... Loblolly Pine  
 P. Thunbergiana ..... Japanese Black Pine  
 P. virginiana ..... Virginia Pine  
 Pistachio sp. .... Pistachio\*  
 Pittosporum rhombifolium ..... Queensland Pittosporum  
 P. tobira ..... Japanese Pittosporum  
 Podocarpus macrophyllus ..... Japanese Yew  
 Prunus laurocerasus ..... English Laurel  
 Prunus sp. .... Almond, Apricot, Nectarine, Peach, Plum and Prune\*  
 Pseudotsuga menziesii ..... Douglas Fir\*\*  
 Pyracantha coccinea ..... Firethorn (Scarlet)  
 P. fortuneana ..... Firethorn  
 P. koidzumii ..... Firethorn  
 Pyrus sp. .... Bradford Pear sp.  
 Quercus shumardii ..... Oak, Shumard's Red  
 Quercus spp. .... Oak species  
 Raphiolepis indica ..... Indian Hawthorne  
 Raphiolepis umbellata ..... Yedda Hawthorne  
 Rhododendron ..... Coral Bells, 'Formosa,' 'Hino-crimson,' 'PJM' (including Azalea) ..... 'Delaware Valley White,' 'Roseum Elegans,' 'Flame Creeper,' 'Girard Crimson,' 'Wakeiebisu,' 'White Gumbo,' 'George L. Tabor'  
 Rosa banksiae ..... Lady Bank's Rose  
 Rosmarinus officinalis ..... Rosemary\*  
 Rudbeckia sp. .... Black-Eyed Susan; Goldstrum  
 Santolina virens ..... Santolina  
 Saxifraga sp. .... Saxifrage; Purple Dome  
 Scabiosa sp. .... Pincushion Flower  
 Sedum album ..... Stonecrop  
 Sedum cauticola ..... Stonecrop; Lidakense  
 Sedum dasyphyllum ..... Stonecrop  
 Sedum spurium ..... Stonecrop; Dragon's Blood  
 Spiraea bumalda ..... Spirea; Anthony Watercr  
 Syzygium paniculatum ..... Australian Brushcherry  
 Taxus cuspidata ..... Japanese Yew  
 T. media ..... Yew  
 Teucrium sp ..... Germander  
 Thalictrum dipterocarpum ..... Meadow Rue  
 Thuja occidentalis ..... American Arborvitae  
 Trachelospermum asiaticum ..... Star Jasmine  
 Tsuga canadensis ..... Canada Hemlock  
 Tulipa spp. .... Tulip species  
 Veronicac sp. .... Veronica, Speedwell; Sunny Border  
 Viburnum japonicum ..... Japanese Viburnum  
 V. odoratissimum ..... Sweet Viburnum  
 V. plicatum ..... Japanese Snowball  
 V. rigidum ..... Canary Island Viburnum  
 V. suspensum ..... Arrowwood Viburnum  
 V. tinus ..... Laurustinus  
 V. trilobium ..... Cranberry Bush  
 V. wrightii ..... Leatherleaf Viburnum  
 Vinca major ..... Greater Periwinkle, Vinca  
 Vinca minor ..... Common Periwinkle, Vinca  
 Vitis sp. .... Grape\*  
 Weigela florida ..... Old Fashioned Weigela  
 Yucca aloifolia ..... Spanish Bayonet, Yucca  
 Y. filamentos ..... Adam's Needle, Yucca  
 \*Do not use on food producing trees, vines or plants.  
 \*\*Not for use on container grown plants.

6046

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in original container in a locked storage area out of reach of children and domestic animals.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

*(Alternate wording for household use:)*

**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse bag. Fold empty bag and discard in trash.

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

OCT 22 1997

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
32802-65

**GUARANTEE:** This product is unconditionally guaranteed.

*(Alternate wording for warranty statement:)*

**GREEN GUARANTEED:** Your satisfaction is guaranteed when this product is used in accordance with instructions. If product performance is less than satisfactory, seller will refund the full purchase price with evidence of purchase.

*(Alternate wording for warranty statement:)*

**NOTICE:** Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

*(Alternate wording for warranty statement:)*

**WARRANTY AND DISCLAIMER:** Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

*(Alternate wording for warranty statement:)*

**WARRANTY AND DISCLAIMER:** Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. No other warranty, expressed or implied, is made concerning the merchantability or fitness for a particular purpose. Manufacturer/seller's liability shall be limited to the purchase price of the product. In no case shall manufacturer/seller be liable for consequential or incidental damages or loss. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

*(Alternate wording for warranty statement:)*

**WARRANTY AND DISCLAIMER:** Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. No other warranty, expressed or implied, is made concerning the merchantability or fitness for a particular purpose. Manufacturer/seller's liability shall be limited to the purchase price of the product. In no case shall manufacturer/seller be liable for consequential or incidental damages or loss. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label. Products not accepted under these terms should be returned within 20 days in the original, unopened container for return of purchase price.

NET WEIGHT: 16 lbs (or 18, 20, 25, 30, 36, 40, 50 or 60 lbs)

(Label for EPA Reg. No 32802-65, PAGE 4 OF 4)

91747