

1/15/2002

1/5



## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

Ms. Kresti A. Lyddon  
Director of Regulatory Affairs  
Howard Johnson's Enterprises, Inc.  
700 W. Virginia St., Suite 222  
Milwaukee, WI 53204-1548

JAN 15 2002

Dear Ms. Lyddon:

Subject: Crabgrass Preventer Plus with .37 Barricade®  
Preemergence Herbicide  
EPA registration No. 32802-66  
Application and Letter Received December 12, 2001,  
Request To Amend Registration by Revising the First  
Aid Statements and the Disposal Statements on the  
Labeling in Compliance with PR Notices 2001-1 and  
2001-6

The proposed subject amendments to the registration of "Crabgrass Preventer Plus with .37 Barricade® Preemergence Herbicide" have been reviewed and found acceptable under the Federal Insecticide, Fungicide and Rodenticide Act, (FIFRA) as amended, provided that you:

1. With any of the alternate wording for warranty statements you must add the commonly used language: "Read the entire "Directions for Use" and "Warranty Statements" before using this product. If terms are not acceptable, return the unopened product container at once to the seller." (You may specify a reasonable time limitation, if "at once" is not preferred.). The above requirement is to assure that you have complied with UCC Articles (Sections 2-719 and 2-316) that apply to the claims for limitation of remedy in the warranty statements.
2. Submit one (1) copy of your final printed labeling prior to your shipment of the subject pesticide product under the enclosed stamped label.


If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

-2-

A stamped copy of the proposed label is enclosed for your records.

Please note that the amendments allowed do not involve exclusive use data, therefore a letter of authorization was not required.

Sincerely yours,

  
for Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505C)

Enclosure

INGREDIENTS: 100.00%

EPA Reg. No. 32802-66

EPA Est. No. 32802-WI-1 (A), 32802-MO-1 (B)

First letter of bag code shows establishment where manufactured.

See tag attached to bag for fertilizer guaranteed analysis.

Manufactured by:

HOWARD JOHNSON'S ENTERPRISES, INC.

700 W. Virginia Street, Milwaukee, WI 53204-1548

BARRICADE® is a registered trademark of Syngenta Crop Protection

## KEEP OUT OF REACH OF CHILDREN

**CAUTION:** See back panel for further precautions and First Aid information.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

**ENVIRONMENTAL HAZARDS:** This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

### First Aid

If in eyes	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
If on skin or clothing	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
If swallowed	<ul style="list-style-type: none"><li>• Call a poison control center or physician immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or physician.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
If inhaled	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>

### Hot line number:

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.


established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds germination and root development. When used according to directions, it will control: banyardgrass; bluegrass, annual (*Poa annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass<sup>1</sup>; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass<sup>2</sup>; shepherd's purse<sup>2</sup>; signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

<sup>1</sup> In many areas a single application of 4 to 9.3 pounds/1000 square feet (175 to 405 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 4 to 6.2 pounds/1000 square feet (175 to 270 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

<sup>2</sup> For this weed, apply in late summer, fall or winter prior to germination.

<sup>3</sup> Suppression only.

**WHEN TO APPLY TO ESTABLISHED TURF:** This product must be applied prior to germination of weed seeds, as it will not control established weeds.



THIS MAP SHOWS  
APPROXIMATE CRABGRASS  
GERMINATION DATES

After May 30  
After May 10  
After April 30  
After March 20  
January 1 to March 20

**APPLICATION DIRECTIONS:** Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days after application.

**USE PRECAUTIONS:** The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil). Do not apply to newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

JAN 15 2002

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

32802-66

(Label for EPA Reg. No 32802-66, PAGE 1 OF 3)

03241

**RATES OF APPLICATION:** This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)		
Turf Species	Minimum Rate	Maximum Annual Rate <sup>1</sup>
Bahia grass, Bermudagrass <sup>2</sup> Centipedegrass, Zoysiagrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type)	4 lbs / 1000 sq ft (175 lbs / acre)	9.3 lbs / 1000 sq ft per year (405 lbs / acre per year)
Buffalograss Kentucky Bluegrass Perennial Ryegrass	4 lbs / 1000 sq ft (175 lbs / acre)	6.2 lbs / 1000 sq ft per year (270 lbs / acre per year)
Creeping Red Fescue	4 lbs / 1000 sq ft (175 lbs / acre)	4.6 lbs / 1000 sq ft per year (202 lbs / acre per year)
Creeping Bentgrass	4 lbs / 1000 sq ft (175 lbs / acre)	4.0 lbs / 1000 sq ft per year (175 lbs / acre per year)

<sup>1</sup>These are the maximum rates per calendar year.

<sup>2</sup>May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 3.3 pounds/1000 square feet (145 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

**WHEN TO OVERSEED AFTER APPLICATION:** This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/ reseeding results.

USE RATE	MONTHS BEFORE OVERSEEDING		
	North	Transition	South
3 pounds / 1000 sq ft (135 pounds / acre)	4	4	4
4 pounds / 1000 sq ft (175 pounds / acre)	5	4	4
4.6 pounds / 1000 sq ft (202 pounds / acre)	6	5	5
4.9 pounds / 1000 sq ft (215 pounds / acre)	---	6	6
6.2 pounds / 1000 sq ft (270 pounds / acre)	---	7	7
6.2 pounds / 1000 sq ft (308 pounds / acre)	---	---	9
9.3 pounds / 1000 sq ft (350 pounds / acre)	---	---	10
9.3 pounds / 1000 sq ft (405 pounds / acre)	---	---	12

**LANDSCAPE ORNAMENTALS** (including established perennials and wildflower plantings): This product may be applied for residual preemergence weed control in ornamentals. Apply at the rate of 4 to 9.3 pounds/1000 square feet (175 to 405 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 9.3 pounds/1000 square feet (405 pounds/acre) per year.

SPREADER SETTINGS to apply at the listed pounds per 1000 square feet						
	3 lbs	4 lbs	4.6 lbs	6.2 lbs	8 lbs	9.3 lbs
Spreader A	X	X	X	X	X	X

(May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product.)

**APPLICATION TIMING AND INFORMATION:** This product may be applied to newly transplanted and established ornamentals a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

**TOLERANT ORNAMENTAL SPECIES:** This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

#### SCIENTIFIC NAME

#### COMMON NAME

Abies spp.	Fir species** (Balsam, Fraser, Noble, etc.)
Abelia grandiflora	Abelia: Sherwood
Acer palmatum	Japanese Maple
A. platanoides	Norway Maple
Achillea spp.	Yarrow: King Edward
Actinidia chinensis	Kiwi*
Agapanthus africanus	Lily-of-the-Nile (African Lily)
Agapanthus orientalis	
Akebia quintata	Five-Leaf or Chocolate Vine
Allium cernuum	Lady's Leek, Nodding Onion
Anemone hybrida	Japanese Anemone
Aquilegia spp.	Aquilegia: Red and Gold
Arctostaphylos densiflora	Vine Hill Manzanita
Arctotheca calendula	Cape Weed
Artemesia spp.	Wormwood; Silver Mound, Castle
Aster spp.	Aster: Bonnie Blue, Purple Dome
Aster X Frikartii	
Athrium Filiz-femina	Lady Fern; Fern Lady
Aucuba japonica	Japanese Aucuba
Begonia spp.	Fibrous Begonia: Hardy Grandis
Berberis gladwynensis	Barberry
B. Julianae	Wintergreen Barberry
B. mentorensis	Mentor Barberry
B. Thunbergii	Japanese Barberry
B. verruculosa	Warty Barberry
Bergenia cordifolia	
Boltonia asteroides	Snowbank
Buddleia davidii	Butterfly-Bush (Dwarf Blue); Royal Red
Buxus microphylla	Japanese Boxwood
Callistemon viminalis	Weeping Bottlebrush
Calluna vulgaris	Scotch Heather
Campanula carpatia	Tussock Bellflower, (White Clips)
Campis X Tagliabuana	Trumpet Flower, Madame Galen
Carpobrotus edulis	Hottentot Fig (Ice Plant)
Cassia artemisoides	Feathery Cassia
Ceanothus rigidus	Wild Lilac
Ceratostigma plumbagonoides	
Chamaecyparis pisifera	False Cypress
Cleyera japonica	Cleyera
Citrus spp.	Citrus species*
Coreopsis spp.	Coreopsis (Calliopsis) Early Sunrise, Moonbeam
Cornus florida	Flowering Dogwood
C. stolonifera	American Dogwood
Cortaderia selioana	Pampas Grass
Cotoneaster apiculatus	Cranberry Cotoneaster
C. buxifolius	Cotoneaster
C. dammeri	Bearberry Cotoneaster
C. microphyllus	Rockspray Cotoneaster
Cretaeus spp.	Hawthorne
Crocsmia spp.	Lucifer
Cupressus sempervirens	Italian Cypress
Delosperma alba	White Trailing Ice Plant
Delosperma spp.	Cooperi Pink
Delphinium spp.	Larkspur: Blue Elf
Dianthus deltoides	Dianthus; Maiden Pinks "Zing"
D. gratianopolitanus	Cheddar Pink
Dodonea viscosa	Hop Bush
Echinacea purpurea	Coneflower, Purple; Magnus
Elaeagnus pungens	Silverberry
Euonymus fortunei	Wintercreeper
E. japonica	Evergreen Euonymus
E. kiautschovica	Spreading Euonymus
Fatsia japonica	Japanese Aralia
Forsythia intermedia	Border Forsythia
Forsythia suspensa	Weeping Forsythia
F. viridissima	Greenstem Forsythia
Gaillardia spp.	Gaillardia, Blanket Flower: "Goblin"
Gardenia jasminoides	Gardenia, Cape-Jasmine
Gaura spp.	
Gentiana dahurica	Gentian
Geranium cinereum	Cranesbill
Gladiolus spp.	Gladiolus species**
Gypsophila repens	Baby's Breath
Hedera helix	English Ivy
Helianthemum spp.	Sunrose
Hemerocallis spp.	Daylily: Aztec Gold, Stella de Oro, Tender Love
Hibiscus	Rose of Sharon**
Hibiscus spp.	Mallow: Disco Belle White
Hibiscus rosa-sinensis	Chinese Hibiscus
Hosta plantaginea	Hosta, Plantain Lily (Fragrant)
H. Sieboldiana	Hosta, "Seersucker"
Houttuynia cordata var. variegata	
Hydrangea macrophylla	Bigleaf Hydrangea
Ilex Bernyi	Holly
Ilex cornuta	Chinese Holly
I. crenata	Japanese Holly
I. opaca	American Holly
I. vomitoria	Yaupon Holly, Schillings
Inula ensifolia	
Iris ensata	Sword-Leaved Iris; Jodlesong
Iris siberica	Siberian Iris; Cabernet
Iris spp.	Iris species**
Jasminum nudiflorum	Winter Jasmine
Juniperus chinensis	Chinese Juniper; Nick's Compact, Parsonii
J. conferta	Shore Juniper; Blue Pacific
J. davurica	Parsoni
J. horizontalis	Creeping Juniper
Juglans spp.	Walnut*
Justicia brandegeana	Shrimp Plant
Lagerstromia indica	Crape Myrtle
Lagerstromia indica X Fauriei	Crape Myrtle; Tuscarora

- Lantana montevidensis . . . . .Weeping Lantana
- Lavender spp. . . . .Lavender; Munstead
- Leonopodium alpinum . . . . .Edelweiss
- Ligustrum amurense . . . . .Amur Privet
- Ligustrum sinense . . . . .Chinese Privet; Variegata
- L. japonicum . . . . .Japanese Privet
- L. lucidum . . . . .Glossy Privet (Waxleaf)
- Lilium spp. . . . .Lily; Jazz
- Liriope muscari . . . . .Big Blue Lily; Silver Mound, Evergreen Giant
- Liriope muscari var. variegata Liriope, Variegated
- Liriope spicata . . . . .Liriope, Creeping
- Lobelia Cardinalis . . . . .Cardinal Flower, Indian Pink
- Lonicera japonica . . . . .Japanese Honeysuckle
- L. tatarica . . . . .Tatarian Honeysuckle
- Lorapetalum chinense . . . . .Burgundy
- Lythrum spp. . . . .Loosestrife; Modern Pink
- Magnolia spp. . . . .Magnolia species\*\*
- Maleophora luteola . . . . .Ice Plant
- Malus spp. . . . .Crabapple\*
- Miscanthus sinensis . . . . .Yaku Jima, Silberfeder
- Nandina domestica . . . . .Heavenly Bamboo
- Narcissus spp. . . . .Narcissus species\*\*
- Nerium oleander . . . . .Oleander
- Olea europaea . . . . .Olive \*
- Ophiopogon japonicus . . . . .Mondo Grass
- nanthus heterophyllus . . . . .Osmanthus (False Holly); Gulf Tide
- ospermum fruticosum . . . . .Trailing African Daisy
- Oxydendrum arboreum . . . . .Sourwood
- Paeonia suffruticosa . . . . .Tree Peony
- Pennisetum setaceum . . . . .Fountain Grass (Dwarf)
- Perovskia atriplicifolia
- Persea americana . . . . .Avocado\*
- Photinia Fraseri . . . . .Fraser's Photinia (Redtip)
- Physotegia Virginiana . . . . .Dragonhead, False; Vivid
- Picea spp. . . . .Spruce species\*\* (Colorado Blue, Norway, etc.)
- Pieris japonica . . . . .Lily-of-the-Valley Bush
- Pinus brutia . . . . .Calabrian Pine
- P. canariensis . . . . .Canary Island Pine
- P. elliptii . . . . .Slash Pine
- P. halepensis . . . . .Aleppo Pine
- P. nigra . . . . .Austrian Pine
- P. palustris . . . . .Longleaf Pine
- P. radiata . . . . .Monterey Pine
- P. strobus . . . . .Eastern White Pine
- P. sylvestris . . . . .Scotch Pine
- P. taeda . . . . .Loblolly Pine
- P. Thunbergiana . . . . .Japanese Black Pine
- rginiana . . . . .Virginia Pine
- achio spp. . . . .Pistachio\*
- Pittosporum rhombifolium . . . . .Queensland Pittosporum
- P. tobira . . . . .Japanese Pittosporum
- Podocarpus macrophyllus . . . . .Japanese Yew
- Prunus laurocerasus . . . . .English Laurel
- Prunus spp. . . . .Almond, Apricot, Nectarine, Peach, Plum and Prune\*
- Pseudotsuga menziesii . . . . .Douglas Fir\*\*
- Pyracantha coccinea . . . . .Firethorn (Scarlet)
- P. fortuneana . . . . .Firethorn
- P. koidzumii . . . . .Firethorn
- Pyrus spp. . . . .Bradford Pear spp.
- Quercus Shumardii . . . . .Oak, Shumard's Red
- Quercus spp. . . . .Oak species
- Raphiolepis indica . . . . .Indian Hawthorne
- Raphiolepis umbellata . . . . .Yedda Hawthorne
- Rhododendron . . . . .Coral Bells, 'Formosa,' 'Hino-crimson,' 'PJM' (including Azalea) . . . . .Delaware Valley White, 'Roseum Elegans,' 'Flame Creeper,' 'Girard Crimson,' 'Wakeiebisu,' 'White Gumbo,' 'George L. Tabor'
- Rosa banksiae . . . . .Lady Bank's Rose
- Rosmarinus officinalis . . . . .Rosemary\*
- Rudbeckia spp. . . . .Black-Eyed Susan; Goldsturm
- Santolina virens . . . . .Santolina
- Saxifraga spp. . . . .Saxifrage; Purple Dome
- Scabiosa spp. . . . .Pincushion Flower
- Sedum album . . . . .Stonecrop
- Sedum cauticola . . . . .Stonecrop; Lidakense
- Sedum dasyphyllum . . . . .Stonecrop
- Sedum spurium . . . . .Stonecrop; Dragon's Blood
- Spiraea bumalda . . . . .Spiraea; Anthony Waterer
- Syzygium paniculatum . . . . .Australian Brushcherry
- Taxus cuspidata . . . . .Japanese Yew
- T. media . . . . .Yew
- Teucrium spp. . . . .Germander
- Thalictrum dipterocarpum . . . . .Meadow Rue
- Thuja occidentalis . . . . .American Arborvitae
- Trachelospermum asiaticum . . . . .Star Jasmine
- Tsuga canadensis . . . . .Canada Hemlock
- Tulipa spp. . . . .Tulip species
- Veronica spp. . . . .Veronica, Speedwell; Sunny Border
- Viburnum japonicum . . . . .Japanese Viburnum
- V. odoratissimum . . . . .Sweet Viburnum
- V. plicatum . . . . .Japanese Snowball
- V. rigidum . . . . .Canary Island Viburnum
- V. suspensum . . . . .Arrowwood Viburnum
- V. tinus . . . . .Laurustinus
- V. trilobium . . . . .Cranberry Bush
- V. wrightii . . . . .Leatherleaf Viburnum
- Vinca major . . . . .Greater Periwinkle, Vinca
- Vinca minor . . . . .Common Periwinkle, Vinca
- Vitis spp. . . . .Grape\*
- Weigela florida . . . . .Old Fashioned Weigela
- Yucca aloifolia . . . . .Spanish Bayonet, Yucca
- Y. filametosa . . . . .Adam's Needle, Yucca

\*Do not use on food producing trees, vines or plants.  
\*\*Not for use on container grown plants.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.  
**STORAGE:** Store in original container in a locked storage area out of reach of children and domestic animals.  
**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.  
**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.  
  
(Alternate wording for household use:)  
**IF EMPTY:** Do not reuse this container. Place in trash or offer for recycling if available.  
**IF PARTLY FILLED:** Call your local solid waste agency for disposal instructions. Never place any unused product down any indoor or outdoor drain or toilet.

**GUARANTEE:** This product is unconditionally guaranteed.  
(Alternate wording for warranty statement:)  
**GREEN GUARANTEED:** Your satisfaction is guaranteed when this product is used in accordance with instructions. If product performance is less than satisfactory, seller will refund the full purchase price with evidence of purchase.  
(Alternate wording for warranty statement:)  
**NOTICE:** Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label.  
(Alternate wording for warranty statement:)  
**WARRANTY AND DISCLAIMER:** Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label.  
(Alternate wording for warranty statement:)  
**WARRANTY AND DISCLAIMER:** Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. No other warranty, expressed or implied, is made concerning the merchantability or fitness for a particular purpose. Manufacturer/seller's liability shall be limited to the purchase price of the product. In no case shall manufacturer/seller be liable for consequential or incidental damages or loss. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label.  
(Alternate wording for warranty statement:)  
**WARRANTY AND DISCLAIMER:** Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. No other warranty, expressed or implied, is made concerning the merchantability or fitness for a particular purpose. Manufacturer/seller's liability shall be limited to the purchase price of the product. In no case shall manufacturer/seller be liable for consequential or incidental damages or loss. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label. Products not accepted under these terms should be returned within 20 days in the original, unopened container for return of purchase price.

(Optional wording required by some states in order to effect fertilizer registration:)  
When soil testing or professional advice indicates the need for additional plant nutrients, make additional applications of fertilizer products over the growing season.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.wa.gov/agr/>

(Optional wording for use with fertilizer inert ingredients that cause staining:)  
This product may stain driveways, walks, patios, pools, etc. Wash or sweep off thoroughly immediately after each use.