

32802-66

4/24/2009

1/7



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

APR 24 2009

Ms. Janet P. Baguhn
Compliance Manager
Howard Johnson's Enterprises Inc.
700 W. Virginia Street #222
Milwaukee, WI 53201

RE: Notification to revise Environmental Hazard Statement PRN 2008-1 and other label changes per
PRN 98-10
EPA Registration Number: 32802-66
Date of Submission: February 24, 2009

Dear Ms. Baguhn:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 and PRN 2008-1, dated February 24, 2009, for the product, Crabgrass Preventer Plus with .37 Prodiamine Preemergence Herbicide. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10, PRN 2008-1 and finds that the actions requested fall within the scope of PRN 98-10 and PRN 2008-1. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs



3/7

February 24, 2009

Document Processing Desk (NOTIF)
Ms. Joanne Miller (PM 23)
Office of Pesticide Programs (7405P)
U.S. Environmental Protection Agency
Room S-4900 One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Product: Crabgrass Preventer Plus With .37 Prodiamine Preemergence Herbicide
EPA Reg. No. 32802-66

Registrant: Howard Johnson's Enterprises, Inc.
700 W. Virginia Street
Milwaukee, WI 53204-1548

Action: Notification of revision of Environmental Hazard Statement as stated
in PR Notice 2008-1 and change to the minimum application rate
from 4.0 pounds per 1000 sq ft (175 lbs/acre) to 3.2 pounds per 1000
sq ft (139 lbs/acre)

Enclosed is a copy of the proposed labeling for the above-mentioned product. This
label reflects the change to the Environmental Hazard statement and the minimum
application rate..


Additionally, one copy of the Application for Pesticide and one copy of the
Certification of Notification of Compliance.

Thank you for your assistance. Please contact me if you have any questions: 414-276-
1505 ext 104, or janet.baguhn@hjfertilizer.com.

Sincerely,

A handwritten signature in black ink, appearing to read 'Janet P. Baguhn', written over a circular stamp.

Janet P. Baguhn
Compliance Manager





Date: February 24, 2009

Product: Crabgrass Preventer Plus With .37 Prodiamine Preemergence Herbicide
EPA Reg. No. 32802-66


Registrant: Howard Johnson's Enterprises, Inc.
700 W. Virginia Street
Milwaukee, WI 53204-1548

Action: Notification of revision of Environmental Hazard Statement as stated in PR Notice 2008-1 and change to the minimum application rate from 4.0 pounds per 1000 sq ft (175 lbs/acre) to 3.2 pounds per 1000 sq ft (139 lbs/acre)

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152-46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and CFR 152-46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

A handwritten signature in black ink, appearing to read 'Janet P. Baguhn', written over a horizontal line.

Janet P. Baguhn
Compliance Manager



**CRABGRASS PREVENTER
PLUS WITH .37 PRODIAMINE
PREEMERGENCE HERBICIDE**
FOR SELECTIVE PREEMERGENCE CONTROL OF GRASS AND
BROADLEAF WEEDS IN LAWNS AND ORNAMENTAL TURF
(excluding golf course putting greens),
LANDSCAPE ORNAMENTALS, ESTABLISHED PERENNIALS
AND WILDFLOWER PLANTINGS

**STOPS MANY UGLY WEED GRASSES BEFORE THEY GROW
PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL & MORE**

ACTIVE INGREDIENT:
Prodiamine: [N³,N³-Di-n-propyl-2,4-dinitro-6-
(trifluoromethyl)-m-phenylenediamine] 0.37%
INERT INGREDIENTS: 99.63%
TOTAL: **100.00%**

NOTIFICATION

EPA Reg. No. 32802-66
EPA Est. No. 32802-WI-1 (A), 32802-~~APR 24~~ 2009
First letter of bag code shows establishment where manufactured.
See tag attached to bag for fertilizer guaranteed analysis.
Manufactured by:
HOWARD JOHNSON'S ENTERPRISES, INC.
700 W. Virginia Street, Milwaukee, WI 53204-1548

KEEP OUT OF REACH OF CHILDREN
CAUTION: See back panel for further precautions
and First Aid information.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Causes moderate eye injury (irritation).
Harmful if inhaled or absorbed through the skin. Avoid contact
with skin, eyes, or clothing. Avoid breathing dust. Wash
thoroughly with soap and water after handling. Prolonged or
repeated skin contact with product may cause allergic
reactions in some individuals.

ENVIRONMENTAL HAZARDS: To protect the environment,
do not allow pesticide to enter or run off into storm drains,
drainage ditches, gutters or surface waters. Applying this
product in calm weather when rain is not predicted for the next
24 hours will help to ensure that wind or rain does not blow or
wash pesticide off the treatment area. Sweeping any product
that lands on a driveway, sidewalk, or street, back onto the
treated area of the lawn or garden will help to prevent run off to
water bodies or drainage systems.

First Aid	
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> Call a poison control center or physician immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or physician. Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Hot line number:
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

(Alternate wording for WPS exclusionary statement, for use when the product labeling contains only turf uses (ie when label is split to exclude the directions for use on ornamental sites:))


Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds germination and root development. When used according to directions, it will control: barnyardgrass; bluegrass, annual (*Poa annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass¹; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass²; shepherd's purse²; signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

¹ In many areas a single application of 3.2 to 9.3 pounds/1000 square feet (139 to 405 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 3.2 to 6.2 pounds/1000 square feet (139 to 270 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).


² For this weed, apply in late summer, fall or winter prior to germination.
³ Suppression only.

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seeds, as it will not control established weeds.



APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days after application.

USE PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil). Do not apply to newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.



16/7

RATES OF APPLICATION: This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

Turf Species	APPLICATION RATES (by Turf Grass Species)	
	Minimum Rate	Maximum Annual Rate*
Bahiagrass, Bermudagrass* Centipede grass, Zoysiagrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type)	3.2 lbs / 1000 sq ft (139 lbs / acre)	9.3 lbs / 1000 sq ft per year (405 lbs / acre per year)
Buffalograss Kentucky Bluegrass Perennial Ryegrass	3.2 lbs / 1000 sq ft (139 lbs / acre)	6.2 lbs / 1000 sq ft per year (270 lbs / acre per year)
Creeping Red Fescue	3.2 lbs / 1000 sq ft (139 lbs / acre)	4.6 lbs / 1000 sq ft per year (202 lbs / acre per year)
Creeping Bentgrass	3.2 lbs / 1000 sq ft (139 lbs / acre)	4.0 lbs / 1000 sq ft per year (175 lbs / acre per year)

*These are the maximum rates per calendar year.
 *May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 3.3 pounds/1000 square feet (145 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

WHEN TO OVERSEED AFTER APPLICATION: This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/ reseeded results.

USE RATE	MONTHS BEFORE OVERSEEDING		
	North	Transition	South
3 pounds / 1000 sq ft (135 pounds / acre)	4	4	4
4 pounds / 1000 sq ft (175 pounds / acre)	5	4	4
4.6 pounds / 1000 sq ft (202 pounds / acre)	6	5	5
4.9 pounds / 1000 sq ft (215 pounds / acre)	---	6	6
6.2 pounds / 1000 sq ft (270 pounds / acre)	---	7	7
7 pounds / 1000 sq ft (308 pounds / acre)	---	---	9
8 pounds / 1000 sq ft (350 pounds / acre)	---	---	10
9.3 pounds / 1000 sq ft (405 pounds / acre)	---	---	12

LANDSCAPE ORNAMENTALS (including established perennials and wildflower plantings): This product may be applied for residual preemergence weed control in ornamentals. Apply at the rate of 3.2 to 9.3 pounds/1000 square feet (139 to 405 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 9.3 pounds/1000 square feet (405 pounds/acre) per year.

SPREADER SETTINGS	to apply at the listed pounds per 1000 square feet					
	3 lbs	4 lbs	4.6 lbs	6.2 lbs	8 lbs	9.3 lbs
Spreader A	X	X	X	X	X	X

(May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product.)

APPLICATION TIMING AND INFORMATION: This product may be applied to newly transplanted and established ornamentals a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

TOLERANT ORNAMENTAL SPECIES: This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

NET WEIGHT: 16 lbs (or 18, 20, 25, 30, 36, 40, 50 or 60 lbs)

SCIENTIFIC NAME	COMMON NAME
Abies spp.	Fir species** (Balsam, Fraser, Noble, etc.)
Abelia grandiflora	Abelia; Sherwood
Acer palmatum	Japanese Maple
A. platanoides	Norway Maple
Achillea spp.	Yarrow; King Edward
Actinidia chinensis	Kiwi*
Agapanthus africanus	Lily-of-the-Nile (African Lily)
Agapanthus orientalis	
Akebia quintata	Five-Leaf or Chocolate Vine
Allium cernuum	Lady's Leek, Nodding Onion
Anemone hybrida	Japanese Anemone
Aquilegia spp.	Aquilegia; Red and Gold
Arctostaphylos densiflora	Vine Hill Manzanita
Arctotheca calendula	Cape Weed
Artemisia spp.	Wormwood; Silver Mound, Castle
Aster spp.	Aster; Bonnie Blue, Purple Dome
Aster X Frikartii	
Athrium Filix-femina	Lady Fern; Fern Lady
Aucuba japonica	Japanese Aucuba
Begonia spp.	Fibrous Begonia; Hardy Grandis
Berberis gladywensis	Barberry
B. Julianae	Wintergreen Barberry
B. mentorensis	Mentor Barberry
B. Thunbergii	Japanese Barberry
B. verruculosa	Warty Barberry
Bergenia cordifolia	
Boltonia asteroides	Snowbank
Buddleia davidii	Butterfly-Bush (Dwarf Blue); Royal Red
Buxus microphylla	Japanese Boxwood
Callistemon viminalis	Weeping Bottlebrush
Calluna vulgaris	Scotch Heather
Campanula carpatica	Tussock Bellflower, (White Clips)
Campis X Tagliabuana	Trumpet Flower, Madame Galen
Carpobrotus edulis	Hottentot Fig (Ice Plant)
Cassia artemisoides	Feathery Cassia
Ceanothus rigidus	Wild Lilac
Ceratostigma plumbagonoides	
Chamaecyparis pisifera	False Cypress
Cleyera japonica	Cleyera
Citrus spp.	Citrus species*
Coreopsis spp.	Coreopsis (Calliopsis) Early Sunrise, Moonbeam
Cornus florida	Flowering Dogwood
C. stolonifera	American Dogwood
Cortaderia selloana	Pampas Grass
Cotoneaster apiculatus	Cranberry Cotoneaster
C. buxifolius	Cotoneaster
C. dammeri	Bearberry Cotoneaster
C. microphyllus	Rockspray Cotoneaster
Cretaegus spp.	Hawthorne
Crococsmia spp.	Lucifer
Cupressus sempervirens	Italian Cypress
Delosperma alba	White Trailing Ice Plant
Delosperma spp.	Cooperi Pink
Delphinium spp.	Larkspur; Blue Elf
Dianthus deltoides	Dianthus; Maiden Pinks "Zing"
D. gratianopolitanus	Cheddar Pink
Dodonea viscosa	Hop Bush
Echinacea purpurea	Coneflower, Purple; Magnus
Elaeagnus pungens	Silverberry
Euonymus fortunei	Wintercreeper
E. japonica	Evergreen Euonymus
E. kiautschovica	Spreading Euonymus
Fatsia japonica	Japanese Aralia
Forsythia intermedia	Border Forsythia
Forsythia suspensa	Weeping Forsythia
F. viridissima	Greenstem Forsythia
Gaillardia spp.	Gaillardia, Blanket Flower: "Goblin"
Gardenia jasminoides	Gardenia, Cape-Jasmine
Gaura spp.	
Gentiana dahurica	Gentian
Geranium cinereum	Cranesbill
Gladiolus spp.	Gladiolus species**
Gypsophila repens	Baby's Breath
Hedera helix	English Ivy
Hellanthemum spp.	Sunrose
Hemerocallis spp.	Daylily; Aztec Gold, Stella de Oro, Tender Love
Hibiscus	Rose of Sharon**
Hibiscus spp.	Mallow; Disco Belle White
Hibiscus rosa-sinensis	Chinese Hibiscus
Hosta plantaginea	Hosta, Plantain Lily (Fragrant)
H. Sieboldiana	Hosta, "Seersucker"
Houttuynia cordata var. variegata	
Hydrangea macrophylla	Bigleaf Hydrangea
Ilex Beryll	Holly
Ilex cornuta	Chinese Holly
I. crenata	Japanese Holly
I. opaca	American Holly
I. vomitoria	Yaupon Holly, Schillings
Inula ensifolia	
Iris ensata	Sword-Leaved Iris; Jodlesong
Iris siberica	Siberian Iris; Cabernet*
Iris spp.	Iris species**
Jasminum nudiflorum	Winter Jasmine
Juniperus chinensis	Chinese Juniper; Nick's Compact, Parsonii
J. conferta	Shore Juniper; Blue Pacific*
J. davurica	Parson
J. horizontalis	Creeping Juniper
Juglans spp.	Walnut*
Justicia brandegeana	Shrimp Plant*
Lagerstromia indica	Crape Myrtle
Lagerstromia indica X Fauriei	Crape Myrtle; Tuscarora

Lantana montevidensis	Weeping Lantana
Lavender spp.	Lavender; Munstead
Leonopodium alpinum	Edelweiss
Ligustrum amurense	Amur Privet
Ligustrum sinense	Chinese Privet; Variegata
L. japonicum	Japanese Privet
L. lucidum	Glossy Privet (Waxleaf)
Lilium spp.	Lily; Jazz
Liriope muscari	Big Blue Lily; Silver Mound, Evergreen Giant
Liriope muscari var. variegata	Liriope, Variegated
Liriope spicata	Liriope, Creeping
Lobelia Cardinalis	Cardinal Flower, Indian Pink
Lonicera japonica	Japanese Honeysuckle
L. tatarica	Tatarian Honeysuckle
Lorapetalum chinense	Burgundy
Lythrum spp.	Loosestrife; Modern Pink
Magnolia spp.	Magnolia species**
Maleophora luteola	Ice Plant
Malus spp.	Crabapple*
Miscanthus sinensis	Yaku Jima, Silberfeder
Nandina domestica	Heavenly Bamboo
Narcissus spp.	Narcissus species**
Nerium oleander	Oleander
Olea europaea	Olive *
Ophiopogon japonicus	Mondo Grass
Osmanthus heterophyllus	Osmanthus (False Holly); Gulf Tide
Osteospermum fruticosum	Trailing African Daisy
Oxydendrum arboreum	Sourwood
Paeonia suffruticosa	Tree Peony
Pennisetum setaceum	Fountain Grass (Dwarf)
Perovskia atriplicifolia	
Persea americana	Avocado*
Photinia Fraseri	Fraser's Photinia (Redtip)
Physotegia Virginiana	Dragonhead, False; Vivid
Picea spp.	Spruce species** (Colorado Blue, Norway, etc.)
Pieris japonica	Lily-of-the-Valley Bush
Pinus brutia	Calabrian Pine
P. canariensis	Canary Island Pine
P. elliottii	Slash Pine
P. halepensis	Aleppo Pine
P. nigra	Austrian Pine
P. palustris	Longleaf Pine
P. radiata	Monterey Pine
P. strobus	Eastern White Pine
P. sylvestris	Scotch Pine
P. taeda	Loblolly Pine
P. thunbergiana	Japanese Black Pine
P. virginiana	Virginia Pine
Pistachio spp.	Pistachio*
Pittosporum rhombifolium	Queensland Pittosporum
P. tobira	Japanese Pittosporum
Podocarpus macrophyllus	Japanese Yew
Prunus laurocerasus	English Laurel
Prunus spp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
Pseudotsuga menziesii	Douglas Fir**
Pyracantha coccinea	Firethorn (Scarlet)
P. fortuneana	Firethorn
P. koidzumii	Firethorn
Pyrus spp.	Bradford Pear spp.
Quercus shumardii	Oak, Shumard's Red
Quercus spp.	Oak species
Raphiolepis indica	Indian Hawthorne
Raphiolepis umbellata	Yedda Hawthorne
Rhododendron	'Coral Bells,' 'Formosa,' 'Hino-crimson,' 'PJM'
(including Azalea)	'Delaware Valley White,' 'Roseum Elegans,'
	'Flame Creeper,' 'Girard Crimson,' 'Wakeiebisu,'
	'White Gumbo,' 'George L. Tabor'
Rosa banksiae	Lady Bank's Rose
Rosmarinus officinalis	Rosemary*
Rudbeckia spp.	Black-Eyed Susan; Goldsturm
Santolina virens	Santolina
Saxifraga spp.	Saxifrage; Purple Dome
Scabiosa spp.	Pincushion Flower
Sedum album	Stoncrop
Sedum cauticola	Stoncrop; Lidakense
Sedum dasyphyllum	Stoncrop
Sedum spurium	Stoncrop; Dragon's Blood
Spiraea bumalda	Spirea; Anthony Waterer
Syzygium paniculatum	Australian Brushcherry
Taxus cuspidata	Japanese Yew
T. media	Yew
Teucrium spp.	Germander
Thalictrum dipterocarpum	Meadow Rue
Thuja occidentalis	American Arborvitae
Trachelospermum asiaticum	Star Jasmine
Tsuga canadensis	Canada Hemlock
Tulipa spp.	Tulip species
Veronica spp.	Veronica, Speedwell; Sunny Border
Viburnum japonicum	Japanese Viburnum
V. odoratissimum	Sweet Viburnum
V. plicatum	Japanese Snowball
V. rigidum	Canary Island Viburnum
V. suspensum	Arrowood Viburnum
V. tinus	Laurustinus
V. trilobium	Cranberry Bush
V. wrightii	Leatherleaf Viburnum
Vinca major	Greater Periwinkle, Vinca
Vinca minor	Common Periwinkle, Vinca
Vitis spp.	Grape*
Weigela florida	Old Fashioned Weigela
Yucca aloifolia	Spanish Bayonet, Yucca
Y. filamentosa	Adam's Needle, Yucca

*Do not use on food producing trees, vines or plants.

**Not for use on container grown plants.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in original container in a locked storage area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

(Alternate wording for household use:)

IF EMPTY: Do not reuse this container. Place in trash or offer for recycling if available.

IF PARTLY FILLED: Call your local solid waste agency for disposal instructions. Never place any unused product down any indoor or outdoor drain or toilet.

GUARANTEE: This product is unconditionally guaranteed.

(Alternate wording for warranty statement:)

GREEN GUARANTEED: Your satisfaction is guaranteed when this product is used in accordance with instructions. If product performance is less than satisfactory, seller will refund the full purchase price with evidence of purchase.

(Alternate wording for warranty statement:)

NOTICE: Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

(Alternate wording for warranty statement:)

WARRANTY AND DISCLAIMER: Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

(Alternate wording for warranty statement:)

WARRANTY AND DISCLAIMER: Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. No other warranty, expressed or implied, is made concerning the merchantability or fitness for a particular purpose. Manufacturer/seller's liability shall be limited to the purchase price of the product. In no case shall manufacturer/seller be liable for consequential or incidental damages or loss. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

(Alternate wording for warranty statement:)

WARRANTY AND DISCLAIMER: Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. No other warranty, expressed or implied, is made concerning the merchantability or fitness for a particular purpose. Manufacturer/seller's liability shall be limited to the purchase price of the product. In no case shall manufacturer/seller be liable for consequential or incidental damages or loss. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label. Products not accepted under these terms should be returned within 20 days in the original, unopened container for return of purchase price.

(Optional wording required by some states in order to effect fertilizer registration:)

When soil testing or professional advice indicates the need for additional plant nutrients, make additional applications of fertilizer products over the growing season.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.wa.gov/agr/>

(Optional wording for use with fertilizer inert ingredients that cause staining:)

This product may stain driveways, walks, patios, pools, etc. Wash or sweep off thoroughly immediately after each use.

(Optional marketing statement when BARRICADE® is formulated in formula:)

Contains .37 BARRICADE® PREEMERGENCE HERBICIDE BARRICADE® is a registered trademark of Syngenta Crop Protection

(Label for EPA Reg. No 32802-66, PAGE 3 OF 3)

00322A