

#### UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Ms. Janet P. Baguhn Compliance Manager Howard Johnson's Enterprises, Inc. 700 W. Virginia Street, Suite 222 Milwaukee, WI 53204

FEB - 1 2010

Dear Ms. Baguhn:

Subject: Crabgrass Preventer Plus With .58 Barricade®
Preemergence Herbicide
EPA Registration No. 32802-67
Application and Letter Dated October 30, 2009,
Request To Amend Registration of Subject Pesticide
Product To Allow Alternate Environmental Hazards
Statements To Accommodate Need for Watering-In
Granules After Application; and Your Letter Dated
January 20, 2010 Resubmission of Revised Labeling
Based on Agency's Review, E-Mailed January 13, 2010

Your request to amend the registration of "Crabgrass Preventer Plus With .58 Barricade® Preemergence Herbicide" as described in the subject application and letters has been reviewed and found acceptable under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, provided that you:

- Delete from the chemical expression from the ingredient Statement or replace it with the CAS Index name for Prodiamine, i.e. 2,4-dinitro-N<sup>3</sup>,N<sup>3</sup>-dipropyl-6-(trifluoromethyl)-1,3-benzenediamine
- 2. Delete the date "10302009", or replace it with the date of this letter.
- 3. Submit one copy of the final printed labeling prior to shipping this product under this amended labeling.

2/6

If these conditions are not complied too, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product under the revised final printer label constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Sincerely yours,

Kathryn V. Montaque

Product Manager (23)

Herbicide Branch

Registration Division (7505P)

Enclosure

# CRABGRASS PREVENTER PLUS WITH .58 BARRICADE® PREEMERGENCE HERBICIDE

FOR SELECTIVE PREEMERGENCE CONTROL OF GRASS AND BROADLEAF WEEDS IN LAWNS AND ORNAMENTAL TURF (excluding golf course putting greens),

LANDSCAPE ORNAMENTALS, ESTABLISHED PÉRENNIALS AND WILDFLOWER PLANTINGS

STOPS MANY UGLY WEED GRASSES BEFORE THEY GROW PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL & MORE

ACTIVE INGREDIENT:
Prodiamine: [N',N'-Di-<sub>D</sub>-propyl-2,4-dinitro-6(trifluoromethyl)-m-phenylenediamine](CAS 29091-21-2) ...0.58%
INERT INGREDIENTS: 99.42%
TOTAL: 100.00%

EPA Reg. No. 32802-67

EPA Est. No. 32802-WI-1 (A), 32802-MO-1 (B)

First letter of bag code shows establishment where manufactured. See tag attached to bag for fertilizer guaranteed analysis. Manufactured by:

HOWARD JOHNSON'S ENTERPRISES, INC. 700 W. Virginia Street, Milwaukee, WI 53204-1548

BARRICADE® is a registered trademark of Syngenta Crop Protection

Read the entire "Directions for Use" and "Warranty Statements" before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

NET WEIGHT: 16 lbs (or 18, 20, 25, 30, 36, 40, 50 lbs)

## KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for further precautions and First Aid information.

## PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

**ENVIRONMENTAL HAZARDS:** This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

# (Alternate wording for home owner use:) ENVIRONMENTAL HAZARDS:

To protect the environment, do not allow this product to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather will help to ensure that wind does not blow this product off the treated area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

| Guaranteed Analysis                              |  |
|--|--|
| Total Nitrogen (N)<br>Available Phosphate (P2O5) | XX.XX%<br>XX.XX%   |
| Soluble Potash (K2O)                             | XX.XX%   |
| Derived from:                                    |  |
|  | ACCEPTED   |
| Chlorine (CI), not more than XX.XX%              | with COMMENTS<br>In EPA Letter Dated:  |
|  | FEB - 1 2010<br>Under the Federal II :: cide,<br>Funnicide and Review de Aci |
|  | as amended, for the puricide<br>registered under EPA Reg. No.                |

32802-67

| First Aid              |  |  |  |
|------------------------|--|--|--|
| If in eyes             | <ul> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>  |  |  |
| If on skin or clothing | <ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>  |  |  |
| If swallowed           | <ul> <li>Call a poison control center or physician immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or physician.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul> |  |  |
| If inhaled             | Move person to fresh air.     If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.     Call a poison control center or doctor for further treatment advice.  Hot line number:   |  |  |

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.


(Alternate wording for WPS exclusionary statement, for use when the product labeling contains <u>only</u> turf uses (ie when label is split to <u>exclude</u> the directions for use on ornamental sites:)

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seed germination and root development. When used according to directions, it will control: barnyardgrass; bluegrass, annual (Poa annua); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass'; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass'; shepherd's purse'; signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

In many areas a single application of 2.5 to 6 pounds/1000 square feet (112 to 259 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 2.5 to 3.9 pounds / 1000 square feet (112 to 173 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label). 
<sup>2</sup> For this weed, apply in late summer, fall or winter prior to germination.

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seed, as it will not control established weeds.



**APPLICATION DIRECTIONS:** Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days after application.

USE PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil). Do not apply to newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (Poa annua) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

RATES OF APPLICATION: This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

| MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)  |   |  |  |  |  |
|--|---|--|--|--|--|
| Turf Species   | Minimum Rate                              | Maximum Annual Rate <sup>1</sup>                           |  |  |  |
| Bahiagrass, Bermudagrass'<br>Centipedegrass, Zoysiagrass<br>Seashore Paspalum<br>St. Augustinegrass<br>Tall Fescue (including turf-type) | 4 lbs / 1000 sq ft<br>(175 lbs / acre)    | 6 lbs / 1000 sq ft per year<br>(259 lbs / acre per year)   |  |  |  |
| Buffalograss<br>Kentucky Bluegrass<br>Perennial Ryegrass   | 2 lbs / 1000 sq ft<br>(87.5 lbs / acre)   | 4 lbs / 1000 sq ft per year<br>(173 lbs / acre per year)   |  |  |  |
| Creeping Red Fescue  | 1.5 lbs / 1000 sq ft<br>(65.3 lbs / acre) | 2.9 lbs / 1000 sq ft per year<br>(130 lbs / acre per year) |  |  |  |
| Creeping Bentgrass   | 1.5 lbs / 1000 sq ft<br>(65.3 lbs / acre) | 2.5 lbs / 1000 sq ft per year<br>(112 lbs / acre per year) |  |  |  |

'These are the maximum rates per calendar year.

<sup>a</sup>May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 2 pounds/1000 square feet (87.5 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

WHEN TO OVERSEED AFTER APPLICATION: This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/ reseding results.

| USE RATE MONTHS BEFORE OVE                   |       | RSEEDING   |       |
|--|-------|------------|-------|
|  | North | Transition | South |
| 1.9 pounds / 1000 sq ft (86 pounds / acre)   | 4     | 4          | 4     |
| 2.5 pounds / 1000 sq ft (112 pounds / acré)  | 5     | 4          | 4     |
| 2.9 pounds / 1000 sq ft (130 pounds / acre)  | 6     | 5          | 5     |
| 3.16 pounds / 1000 sq ft (138 pounds / acre) |       | 6          | 6     |
| 4 pounds / 1000 sq ft (173 pounds / acre)    |       | 7          | 7     |
| 4.5 pounds / 1000 sq ft (197 pounds / acre)  |       |            | 9     |
| 5.14 pounds / 1000 sq ft (224 pounds / acre) |       |            | 10    |
| 5.94 pounds / 1000 sq ft (259 pounds / acre) | l     | (          | 12    |

| SPREADER SETTINGS to apply at the listed pounds per 1000 square feet |         |         |          |   |       |       |
|--|---------|---------|----------|---|-------|-------|
|  | 1.9 lbs | 2.5 lbs | 3.16 lbs |   | 5 lbs | 6 lbs |
| Spreader A   | х       | X       | х        | х | Х     | х     |

(May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product )

LANDSCAPE ORNAMENTALS (including established perennials and wildflower plantings): This product may be applied for residual preemergence weed control in ornamentals. Apply at the rate of 2.5 to 5.94 pounds/1000 square feet (112 to 259 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 5.94 pounds/1000 square feet (259 pounds/acre) per year.

APPLICATION TIMING AND INFORMATION: This product may be applied to newly transplanted and established ornamentals a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

TOLERANT ORNAMENTAL SPECIES: This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.



| SCIENTIFIC NAME  | COMMON NAME  | Ligustrum amurense                        | Amur Privet                                      |
|--|--|---|--|
| Abies spp  | .Fir species** (Balsam, Frasier, Noble, etc.)                        | Ligustrum sinense                         |  |
| Abelia grandiflora   | .Abelia: Sherwood  | L. japonicum                              | Japanese Privet                                  |
| Acer palmatum  |  | L. lucidum                                |  |
| A. platanoides   | Norway Maple   | Lillium spp                               |  |
| Actinidia chinensis  |  | Liriope muscari var. variegata            | .Big Blue Lily; Silver Mound, Evergreen Giant    |
| Agapanthus africanus   |  | Liriope spicata                           | Liriope, Creeping                                |
| Agapanthus orientalis  | ,                              | Lobelia Cardinalis                        | .Cardinal Flower, Indian Pink                    |
| Akebia quintata  | .Five-Leaf or Chocolate Vine   | Lonicera japonica                         |  |
|  | .Lady's Leek, Nodding Onion  | L. tatarica                               |  |
| Anemone hydrida  |  | Lorapetalum chinense                      |  |
| Arctostaphylos densiflora  | Vine Hill Manzanita  | Lythrum spp                               |  |
| Arctotheca calendula   | .Cape Weed   | Maleophora luteola                        |  |
|  | .Wormwood; Silver Mound, Castle                                      | Malus spp                                 | .Crabapple*                                      |
|  | .Aster: Bonnie Blue, Purple Dome                                     | Miscanthus sinensis                       |  |
| Aster X Frikartii  |  | Nandina domestica                         |  |
| Athrium Filiz-femina Aucuba japonica                                   |  | Narcissus spp                             |  |
|  | Fibrous Begonia: Hardy Grandis                                       | Olea europaea                             |  |
| Berberis gladwynensis  |  | Ophiopogon japonicus                      |  |
| B. Julianae  |  |   | .Osmanthus (False Holly): Gulf Tide              |
| B. mentorensis   |  | Osteospermum fruticosum .                 | Trailing African Daisy                           |
| B. Thunbergii  |  | Oxydendrum arboreum                       | .Sourwood  |
| Bergenia cordifolia  | .warty barberry  | Paeonia suffruticosa Pennisetum setaceumi |  |
| Boltonia asteroides  | Snowbank   | Perovskia atriplicifolia                  | .i odnam drass (Dwarr)                           |
|  | .Butterfly-Bush (Dwarf Blue); Royal Red                              | Persea americana                          |  |
| Buxus microphylla  |  | Photinia Fraseri                          | .Frasier's Photinia (Redtip)                     |
| Callistemon viminalis  |  | Physotegia Virginiana                     | .Dragonhead, False; Vivid                        |
| Calluna vulgaris   |  |   | .Spruce species** (Colorado Blue, Norway, etc.)  |
|  | .Tussock Beliflower, (White Clips) .Trumpet Flower, Madame Galen     | Pieris japonica                           | .Calabrian Pine                                  |
| Carpobrotus edulis   |  | P. canariensis                            |  |
| Cassia artemisoides  | .Feathery Cassia   | P. elliottii                              | .Slash Pine                                      |
| Ceanothus rigidus  |  | P. halepensis                             |  |
| Ceratostigma plumbagonoide   |  | P. nigra                                  |  |
| Chamaecyparis pisifera Cleyera japonica                                |  | P. palustrus                              | Longleat Pine Monterey Pine                      |
| Citrus spp   |  | P. strobus                                |  |
|  | .Coreopsis (Calliopsis) Early Sunrise, Moonbeam                      | P. sylvestris                             |  |
| Cornus florida   | .Flowering Dogwood   | P. taeda                                  | .Loblolly Pine                                   |
| C. stolonifera   |  | P. Thunbergiana                           | Japanese Black Pine                              |
| Cortaderia selloana  |  | P. virginiana                             |  |
| Cotoneaster apiculatus C. buxifolius                                   |  | Pistachio spp                             |  |
| C. dammeri   |  | P. tobira                                 |  |
| C. microphyllus  | .Rockspray Cotoneaster   | Podocarpus macrophyllus                   |  |
| Cretaegus spp  |  | Prunus laurocerasus                       |  |
| Crocosmia spp  |  | Prunus spp                                | Almond, Apricot, Nectarine, Peach, Plum and      |
| Cupressus sempervirens Delosperma alba                                 |  | Pseudotsuga menziesii                     | Prune*   |
| Delosperma spp   | .Cooperi Pink  | Pyracantha coccinea                       |  |
| Delphinium spp   | .Larkspur: Blue Elf  | P. fortuneana                             |  |
|  | .Dianthus; Maiden Pinks "Zing"                                       | P. koidzumii                              |  |
| D. gratianopolitanus   |  | Pyrus spp                                 |  |
| Dodonea viscosa Echinacea purpurea                                     | Constant Purple: Magnus  | Quercus Shumardii                         | Oak species                                      |
| Elaeagnus pungens  | Cilventer  | Raphiolepsis indica                       |  |
| Euonymus fortunei  |  | Raphiolepsis umbellata                    |  |
| E. japonica  |  |   | .'Coral Bells,' 'Formosa,' 'Hino-crimson,' 'PJM' |
| E. kiautschovica   |  | (including Azalea)                        | . 'Delaware Valley White', 'Roseum Elegans,'     |
| Fatsia japonica  |  |   | 'Flame Creeper,' 'Girard Crimson,' Wakeiebisu,   |
| Forsythia intermedia Forsythia suspensa                                |  | Rosa banksiae                             | 'White Gumbo,' 'George L. Tabor'                 |
| F. viridissima   |  | Rosmarinus officinalis                    |  |
|  | .Gaillardia, Blanket Flower: "Goblin"                                |   | .Black-Eved Susan; Goldsturm                     |
| Gardenia jasminoides   |  | Santolina virens                          |  |
| Gaura spp.   | <b></b>  | Saxifraga spp                             |  |
| Gentiana dahurica  |  | Scabiosa spp                              |  |
| Geranium cinereum  |  | Sedum cauticola                           |  |
| Gypsophilia repens   |  | Sedum dasyphyllum                         |  |
| Hedera helix   | .English Ivy   | Sedum spurium                             | .Stonecrop; Dragon's Blood                       |
| Hellanthemum spp   |  | Spiraea bumalda                           | .Spirea: Anthony Waterer                         |
|  | .Daylily: Aztec Gold, Stella de Oro, Tender Love                     | Syzygium paniculatum                      |  |
| Hibiscus   |  | Taxus cuspidata                           |  |
| Hibiscus rosa-sinensis   |  | Teucrium spp                              |  |
| Hosta plantaginea  | Hosta Plantain Lily (Fragrant)                                       | Thalictrum diptercarpum                   |  |
| H. Sieboldiana   | .Hosta, "Seersucker"   | Thuja occidentalis                        | .American Arborvitae                             |
| Houttuynia cordata var. varie  | gata   | Trachelospermum asiatum .                 |  |
| Hydrangea macrophylla  |  | Tsuga canadensis                          |  |
| Ilex Bernyi  |  | Tulipa spp                                | .Veronica, Speedwell; Sunny Border               |
| I. crenata   |  | Viburnum japonicum                        |  |
| I. opaca   |  | V. odoratissimum                          |  |
| I. vomitoria   |  | V. plicatum                               | .Japanese Snowball                               |
| Inula ensifolia  | Occasional to the state of the                                       | V. rigidum                                |  |
|  | .Sword-Leaved Iris; Jodlesong  | V. suspensum                              |  |
| Iris siberica  |  | V. tinus                                  |  |
| Jasminium nudiflorum   |  | V. wrightii                               |  |
|  | .Chinese Juniper; Nick's Compact, Parsonnii                          | Vinca major                               |  |
| J. conferta  | Shore Juniper; Blue Pacific  | Vinca minor                               | .Common Periwinkle, Vinca                        |
| J. davurica  |  | Vitis spp                                 |  |
| J. horizontalis  |  | Weigela florida                           |  |
| Jugians spp  |  | Yucca aloifolia                           |  |
| vasiona vialiutytalia  | Crops Muste  | *Do not use on food producir              |  |
| Lagerstromia indica  |  |   |  |
| Lagerstromia indica Lagerstromia indica X Faurie                       | Crape Myrtle; Tuscarora  | **Not for use on container gr             |  |
| Lagerstromia indica Lagerstromia indica X Faurie Lantana montevidensis | Crape Myrtle; Tuscarora<br>.Weeping Lantana                          | **Not for use on container gr             | own plants.                                      |
| Lagerstromia indica Lagerstromia indica X Faurie                       | i Crape Mýrtle; Tuscarora<br>.Weeping Lantana<br>.Lavender; Munstead | **Not for use on container gr             |  |

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in original container in a locked storage area out of reach of children and domestic animals.

(wording for Commercial use label:)

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

<u>CONTAINER DISPOSAL</u>: Completely empty bag into application equipment. Nonrefillable container: Do not reuse or refill this container. Offer for recycling, if available. If not available then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

(Alternate wording for home owner use:)

IF EMPTY: Nonrefillable container: Do not reuse or refill this container. Offer for recycling, if available. IF PARTLY FILLED: Call your local solid waste agency for disposal instructions. Never place any unused product down any indoor or outdoor drain or toilet.

Warranty

The directions for use of this product are believed to be adequate and must be followed carefully, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result due to such factors as weather conditions. presence or absence of other materials, or the manner of use or application, all of which are beyond the control of Howard Johnson's Enterprises, Inc the manufacturer, or the seller. To the extent consistent with applicable law, the products sold to you are furnished "as is" by Howard Johnson's Enterprises, Inc. The manufacturer and the seller are subject only to the manufacturer's warranties, if any, which appear on the label of the product sold to you. Except as warranted by this label Howard Johnson's Enterprises, Inc, the manufacturer, or the seller makes no warranties, guarantees, or representations of any kind to the buyer or the user, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold or use of the product, including, but not limited to, merchantability, fitness for a particular purpose or use, or eligibility of the product for any particular trade usage. To the extent consistent with applicable law, Buyer's or user's exclusive remedy, and Howard Johnson's Enterprises, Inc the manufacturer's or the seller's total liability shall be limited to damages not exceeding the cost of the product. No agent or employee of Howard Johnson's Enterprises, Inc, or the seller is authorized to amend the terms of this warranty disclaimer or the product's label or to make a presentation or recommendation different from or inconsistent with the label of this product. To the extent consistent with applicable law, Howard Johnson's Enterprises, Inc, the manufacturer, or the seller shall not be liable for consequential, special, or indirect damages resulting from the use, handling, application, storage, or disposal of this product or for damages in the nature of penalties, and the buyer and the user waive any right that they may have to such damages. purchase price with evidence of purchase.

(Optional wording required by some states in order to effect fertilizer registration:)

When soil testing or professional advice indicates the need for additional plant nutrients, make additional applications of fertilizer products over the growing season.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.wa.gov/agr/

(Optional wording for use with fertilizer inert ingredients that cause staining:)

This product may stain driveways, walks, patios, pools, etc. Wash or sweep off thoroughly immediately after each use.

(Optional marketing statements:)

STOPS MANY WEED GRASSES BEFORE THEY GROW

3-5 month Weed Grass Control

### **Optional Fertilizer Marketing Claims**

Contains slow release Nitrogen for extended greening

Contains slow release Nitrogen for extended feeding

Nitrogen for a lush green lawn

Phosphate improves root growth

Potassium Vital for growth & drought resistance

Super high Iron! for lush deep green color

Increases efficiency of natural soil elements including nitrogen, phosphate and potassium

with controlled release nitrogen

covers up to XXXXX square feet (per application)

Phosphate free fertilizer