

11/30/2012



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Wendy Danowski
Howard Johnson's Enterprises, Inc.
9675 S. 60th Street
Franklin, WI 53132

NOV 30 2012

Subject: Notification; Per PR-Notice 98-10
Dithiopyr 0.086 Plus
EPA Reg. No. 32802-68
Date Submitted: November 16, 2012

Proposed Alternate Brand Names:

Dyna Green Crabgrass Preventer with 0.086% Dithiopyr with Nutri-Life
Howard Johnson's Prevents Crabgrass Plus with 0.086% Dimension
Howard Johnson's Prevents Crabgrass Plus with 0.086% Dithiopyr
Pennington Seed Prevents Crabgrass Plus with 0.086% Dimension
Pennington Seed Prevents Crabgrass Plus with 0.086% Dithiopyr
Terre Prevents Crabgrass Plus with 0.086% Dimension

Dear Ms. Danowski:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated November 16, 2012 for the product referenced above. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions regarding this letter, please contact me at (703) 306-0415 or davis.kable@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Kable Bo Davis", written over a horizontal line.

Kable Bo Davis
Product Manager 25
Herbicide Branch
Registration Division (7505P)

(A)

EPA

United States Environmental Protection Agency
Office of Pesticide Programs (H7505C)
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide:

Section I

1. Company/Product Number Howard Johnson's Ent./32802-68	2. EPA Product Manager Kathryn Montague	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Howard Johnson's Ent / Dithiopyr 0.086 Plus	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Howard Johnson's Enterprises, Inc 9675 S 60th Street, Franklin, WI 53132 <input checked="" type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRT Section 3©(3) (b)(l), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section II

<input type="checkbox"/> Amendment - Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION NOV 30 2012 c c c c c c
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - explain below.	

Explanation: Use additional page(s) if necessary. (For Section I and Section II.) Notification per PR 98-10
Submission of 6 new Alternate Brand Names Per 98-10; Dyna Green Crabgrass Preventer Plus with 0.086% Dithiopyr; with Nutri-Life; Howard Johnson's Prevents Crabgrass Plus with 0.086% Dithiopyr; Howard Johnson's Prevents Crabgrass Plus with 0.086% Dimension; Pennington Seed Prevents Crabgrass Plus with 0.086% Dimension; Pennington Seed Prevents Crabgrass Plus with 0.086% Dithiopyr; Terre Prevents Crabgrass Plus with 0.086% Dimension
 This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. Since this is a notification, we expect no PRIA fee is required. For further information, please contact Wendy Danowski: E-mail: wdanowski@central.com; Telephone: (414) 304-3590

Section III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No.	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
*Certification must be submitted.		If "Yes," Unit Package wgt.	No. per container	If "Yes," Package wgt.	No. per container
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) of Retail Container 10# - 50#		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner In Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		Other (paper in clear poly envelope)			

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Wendy Danowski	Title Regulatory Specialist	Telephone No. (Include Area Code) (414) 304-3590
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Regulatory Specialist	
4. Typed Name Wendy Danowski	5. Date 11-16-2012	



November 16, 2012

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

ATTENTION: K. Montague, PM23

SUBJECT: Per PRN 98-10 NOTIFICATION OF ALTERNATE BRAND NAME
For DITHIOPYR 0.086% Plus
EPA Reg. No. 32802-68

•••••
•••••
•••••

Dear Ms. Montague,

•••••
•••••
•••••
•••••
•••••

Enclosed is a completed EPA form 8570-1 and 1 copy of revised labeling for the above product. The label cover sheet details the change. The alternate brand names are:

- Dyna Green Crabgrass Preventer Plus with 0.086% Dithiopyr with Nutri-Life
- Howard Johnson's Prevents Crabgrass Plus with 0.086% Dimension
- Howard Johnson's Prevents Crabgrass Plus with 0.086% Dithiopyr
- Pennington Seed Prevents Crabgrass Plus with 0.086% Dimension
- Pennington Seed Prevents Crabgrass Plus with 0.086% Dithiopyr
- Terre Prevents Crabgrass Plus with 0.086% Dimension

•••••
•••••
•••••
•••••

If you have any questions or need further assistance please contact me directly at 414-304-3590. Thank you for your assistance.

Sincerely,
HOWARD JOHNSON'S ENT., INC.

Wendy Danowski
Regulatory Specialist
w.danowski@hjfertilizer.com

Enclosures: Notification of Additional Brand Name
Copy of Product Label
Certification

Crabgrass Preventer with **Nutri Life**
15-0-5
with 0.086% DITHIOPYR

GUARANTEED ANALYSIS

Total Nitrogen (N)	15.00%
15% Urea Nitrogen	
Soluble Potash (K ₂ O)	5.00%
Boran (B)	0.05%
Copper (Cu)	0.13%
0.03% Water Soluble Copper (Cu)	
Iron (Fe)	0.25%
0.03% Water Soluble Iron (Fe)	
Manganese (Mn)	0.13%
0.03% Water Soluble Manganese (Mn)	
Molybdenum (Mo)	0.0013%
Zinc (Zn)	0.13%
0.03% Water Soluble Zinc (Zn)	

Derived From: COPPER OXIDE, COPPER SULFATE, IRON OXIDE, IRON SULFATE, MANGANESE OXIDE, MANGANESE SULFATE, SODIUM BORATE, MURIATE OF POTASH, UREA

Chlorine (Cl), not more than **NOV 30 2002**
 F315
FG61798
Net weight 18 lbs (8.20 kg)
One bag covers up to 5,400 square feet

PREVENTS CRABGRASS
 SELECTIVE HERBICIDE PLUS FERTILIZER FOR PREEMERGENCE CONTROL OF LISTED ANNUAL GRASSES AND BROADLEAF WEEDS IN ESTABLISHED LAWNS & ORNAMENTAL TURF ACTIVE INGREDIENT:

Dithiopyr: 3,5-pyridinedicarbonyloxy acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,2-dimethyl ester 0.086%
 S-Soluble Potash 99.914%
 INERT INGREDIENTS 100.000%
 TOTAL

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
 CAUTION: Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or repeated skin contact may cause allergic reaction in some individuals.

ENVIRONMENTAL HAZARDS: To protect the environment, do not allow this product to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather will help to ensure that wind does not blow this product off the treated area. Sweeping any product that lands on a driveway, sidewalk or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

STORAGE AND DISPOSAL
STORAGE: Store in original container in a locked storage area out of reach of children and domestic animals.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container.
IF EMPTY: Place in trash or offer for recycling if available.
IF PARTIALLY FILLED: Call your local solid waste agency for disposal instructions. Never place any unused product down any indoor or outdoor drain.

Manufactured by
 Howard Johnson's Enterprises, Inc.
 9875 South 60th Street
 Franklin, WI 53132

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapco.org/metals.htm>
 Dyna Green® with fertilizer to help green up turf.



DIRECTIONS FOR USE: This is a violation of Federal law to use this product in a manner inconsistent with its labeling. In the State of New York: For use by professional turf applicators only, at no more than 13.3 lbs per 1000 square feet per year (0.5 pounds AI per acre per year). Not for use in Nassau or Suffolk Counties.

PRODUCT INFORMATION: This product contains turf fertilizer plus a selective herbicide for control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turf. Apply this product only to seeded, sodded, or sprigged turf that is well established. Turf must develop a good root system and uniform stand, and have received at least two mowings following seeding, sodding or sprigging before making the first application of this product. Use of this product on turf that is not well established, or weakened by weather, pests or other stress, may increase the chance of turf injury.

WHERE TO USE: This product may be used on the following established turfgrasses: Creeping Bentgrass; Aproxis palustris; Kentucky Bluegrass (Poa pratensis); Fine Fescue; (Festuca rubra); Tall Fescue; (Festuca arundinacea); Perennial Ryegrass (Lolium perenne); Bahiagrass (Paspalum notatum); Bermudagrass (Cynodon dactylon); Buffalo grass; (Bouteloua dactyloides); Carpetgrass (Axonopus compressus); Centipedegrass (Eremochloa ophioides); Kikuyugrass (Pennisetum clandestinum); St. Augustinegrass (Stenotaphrum secundatum); Zoysisgrass (Zoysia japonica). Turfgrass injury may result from use of this product on certain varieties of creeping bentgrass, such as Cohasset, Carmen, Seaside and Washington. Not all varieties of creeping bentgrass have been tested. Do not apply this product to Colonial Bentgrass (Agrostis tenuis) varieties. Turfgrass injury may result from use of this product on the following Fescue varieties which are sensitive to this product: Allgate, Banner, Beauty, Bigart, CF-2, HF-93, Highlight, Nolo, Koker, Jameson, Melanta, Maly, Pennlawn, Tamara, Talland, Walcott and Waldine. Not all varieties of fescue have been tested.

SPRING APPLICATIONS: For single applications in the spring or early summer, use the appropriate rate from Application Rates (table depending on user's location, mowing height, pre- or post-emergence use). The duration of residual weed control is directly related to the local rate applied, but varies depending on weather, weed pressure and location. Use Program 1 for pre-emergence control on areas where turfs are high (turf level) for 3 to 5 months of crabgrass control. Use Program 2 for pre-emergence control where turfs are low (turf level) for 3 to 5 months of crabgrass control. This product may be applied as a single application or as a split application for crabgrass control in the spring, summer or fall.

APPLICATION RATES for Crabgrass Control

REGION	PROGRAM 1 PREEMERGENCE USE (high cut turfgrass)	PROGRAM 2 PREEMERGENCE USE (low cut turfgrass)	POST-EMERGENCE USE 6.67 lbs / 1000 sq ft
North	3.33 lbs / 1000 sq ft (145 lbs / acre)	4.89 lbs / 1000 sq ft (213 lbs / acre)	9.77 lbs / 1000 sq ft (425 lbs / acre)
Transition	4.89 lbs / 1000 sq ft (213 lbs / acre)	6.67 lbs / 1000 sq ft (290 lbs / acre)	13.3 lbs / 1000 sq ft (580 lbs / acre)
South	6.67 lbs / 1000 sq ft (290 lbs / acre)	9.77 lbs / 1000 sq ft (425 lbs / acre)	13.3 lbs / 1000 sq ft (580 lbs / acre)
Coastal South	9.77 lbs / 1000 sq ft (425 lbs / acre)	13.3 lbs / 1000 sq ft (580 lbs / acre)	16.67 lbs / 1000 sq ft (733 lbs / acre)

FALL APPLICATION: This product can also be applied in late summer or early fall (late August through November) at the Program 2 rate for crabgrass control through the early part of the next spring. Fall applications must be followed by an appropriate spring application to provide season-long control.

USE PRECAUTIONS: Do not apply more than 13.3 pounds per 1000 square feet (0.5 pounds AI per acre) per application. Do not apply more than 13.3 pounds AI per acre per year. (In New York, do not apply more than 13.3 lbs per 1000 square feet per year.)

TIPS FOR IMPROVED CONTROL: For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. Irrigation or rainfall within 7 days after application may improve weed control. Use of split (half rate) applications spaced 6-8 weeks apart may provide improved weed control.

RESEEDING, OVERSEEDING, OR SPRIGGING: Reseeding, overseeding, or sprigging of treated areas within 4 months after application will help to prevent crabgrass establishment. Follow good cultural practices such as soil cultivation, irrigation and fertilization. For best results, use "soil seeders" for good seed-to-soil contact.

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS: This product will also control or suppress the following weeds when applied prior to their emergence: Barnyardgrass, Bluegrass (annual, roughstalk), Crowfootgrass, Foxtail (yellow & green), Goosegrass, Kikuyugrass, Lespedeza (common), Oxalis (buttercup; creeping & yellow woodsoorrel), Purslane (common), Ryegrass (annual & perennial), Smartweed, Speedwell (corn), Spurge (prostrate and spotted), Foxtail is also controlled at the program 2 rates.

FALL APPLICATIONS: Used as directed in late summer or early fall, for crabgrass control through the early part of the next spring, this product will also control or suppress the following weeds when applied prior to their emergence: Bluegrass (annual; roughstalk), Buttercrunch, Chickweed, Geranium (Carolina), Henbit, Parsley-plant, Pineappleweed, Shepherd's purse.

SPREADER SETTINGS to apply at the listed pounds per 1000 sq. ft.

Settings shown - refer to equipment manual for more information.	3.33 lbs	4.89 lbs	6.67 lbs	9.77 lbs
Times Drop or Rotary	14	17	19	23.5
Can/May Drop or Rotary	6	10	11	23.5
Reaction Red Drop Rotary	5	7	8	open
Republic EZ Rotary	5.5	5	5	8
Cydone SB3000 Rotary	6.75	7.25	7.75	8.5
Scotts Easygreen EG-3 (rotary)	27	28	29	31
Scotts Speedy Green (rotary)	6	8	11	16

Dithiopyr 0.086 Retail PPI (10-9-2012)



Howard Johnson's

PREVENTS CRABGRASS PLUS with 0.086% DITHIOPYR

SELECTIVE HERBICIDE PLUS FERTILIZER FOR PREEMERGENCE CONTROL OF LISTED ANNUAL GRASSES AND BROADLEAF WEEDS IN ESTABLISHED LAWNS & ORNAMENTAL TURF

ACTIVE INGREDIENT:

- Dithiopyr, 3,5-pyridinedicarboxylic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-5-
- (trifluoromethyl)-S,S-dimethyl ester 0.086%
- INERT INGREDIENTS: 99.914%
- TOTAL: 100.000%

EPA Reg. No. 32802-68
 EPA Est. No. 59144-GA-1(A), 32802-MO-1(B), 37351-CO-1(C),
 8378-IN-1 (D), 9404-FL-1 (E)
 First letter of bag code shows establishment where manufactured.

Read the entire "Directions for Use" and "Warranty Statement" before using this product. If terms are not acceptable, this product may be returned for a full refund at once to the seller.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
 Irritant if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact may cause irritation. Avoid contact with water. Do not contaminate water. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

First Aid	<ul style="list-style-type: none"> Hold eye open and flush slowly and gently with water for 15-20 minutes. Remove contaminated clothing. If retained, after the first 15 minutes, trim contaminated material. Call a poison control center or doctor for treatment advice. Get person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if trained. Call a poison control center or doctor for further treatment advice. Take off contaminated clothing. Flush skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Get person to hospital or physician immediately for treatment. Do not induce vomiting unless told to do so by the poison control center. Do not give anything by mouth to an unconscious person. Have this number: 1-800-852-2728.
If in eyes	<ul style="list-style-type: none"> Hold eye open and flush slowly and gently with water for 15-20 minutes. Remove contaminated clothing. If retained, after the first 15 minutes, trim contaminated material. Call a poison control center or doctor for treatment advice. Get person to hospital or physician immediately for treatment. Do not induce vomiting unless told to do so by the poison control center. Do not give anything by mouth to an unconscious person. Have this number: 1-800-852-2728.
If inhaled	<ul style="list-style-type: none"> Get person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if trained. Call a poison control center or doctor for further treatment advice. Take off contaminated clothing. Flush skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Get person to hospital or physician immediately for treatment. Do not induce vomiting unless told to do so by the poison control center. Do not give anything by mouth to an unconscious person. Have this number: 1-800-852-2728.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Flush skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Get person to hospital or physician immediately for treatment. Do not induce vomiting unless told to do so by the poison control center. Do not give anything by mouth to an unconscious person. Have this number: 1-800-852-2728.
If on hands	<ul style="list-style-type: none"> Wash hands thoroughly with soap and water. Call a poison control center or doctor for treatment advice. Get person to hospital or physician immediately for treatment. Do not induce vomiting unless told to do so by the poison control center. Do not give anything by mouth to an unconscious person. Have this number: 1-800-852-2728.

STORAGE AND DISPOSAL
 Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store in original container in a locked storage area out of reach of children and domestic animals.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.
 Information regarding the contents and levels of metals in this product is available on the internet at <http://www.epa.gov/pesticides>.
 Manufactured for: Howard Johnson's Enterprises, Inc., 9675 S 60th Street, Franklin, WI 53122

GUARANTEED ANALYSIS

NOTIFICATION

NOV 30 2012

Net Weight: 50 lbs (22.70 kg)

DIRECTIONS FOR USE: Use this herbicide on established turfgrass in areas where crabgrass is a major problem, at a rate more than 13.3 lbs per 1000 square feet per year (0.5 pounds AI per acre per year). Do not use in Nassau or Suffolk Counties.

PREEMERGENCE CONTROL: This product contains turf fertilizer plus a selective herbicide for control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turf. Apply this product only to seeded, sodded, or sprigged turf that is well established. Turf must develop a good root system before application of this product. Use of this product on turf that is not well established, or weakened by weather, pests or other stress, may increase the chance of turf injury. This product provides pre-emergence control of crabgrass (large, smooth, and southern species) when applied prior to emergence of crabgrass from the ground. In addition, this product provides "early post-emergence" control during the early stages of growth after crabgrass emerges from the ground. This early post-emergence crabgrass control is limited, and post-emergence applications are only effective if applied before crabgrass has displayed its fifth leaf or first tiller which generally corresponds to turf when year can be seen in crabgrass plants (see lawn). The benefit of this post-emergence activity is that it gives the user a 2 to 4 week grace period for application of crabgrass control.

WHERE TO USE: This product may be used on the following established turfgrasses: Creeping Bentgrass (*Agrostis palustris*), Kentucky Bluegrass (*Poa pratensis*), Fine Fescue* (*Festuca rubra*), Tall Fescue (*Festuca arvensis*), Perennial Ryegrass (*Lolium perenne*), Annual Ryegrass (*Lolium multiflorum*), Bermudagrass (*Cynodon dactylon*), Buffalograss** (*Bouteloua curtipendula*), Carpetgrass (*Axonopus compressus*), Centipedegrass (*Eremochloa ophiuroides*), Kikuyograss (*Pennisetum clandestinum*), St. Augustinegrass (*Strombotriophum scandentum*), Zoysiagrass (*Zoysia japonica*), and Zoysia tenuifolia (*Zoysia tenuifolia*). Do not apply this product on certain varieties of creeping bentgrass, such as: Contender, Carmen, Versatile and Washington. Not all varieties of creeping bentgrass have been tested. Do not apply this product to Coastal Bentgrass (*Agrostis tenuis*) varieties.

REGION	PREEMERGENCE USE (high cut turfgrasses)	PROGRAM 1 (low cut turfgrasses)	PROGRAM 2 (low cut turfgrasses)
North	3.33 lbs / 1000 sq ft (0.145 lbs AI / acre)	4.83 lbs / 1000 sq ft (0.20 lbs AI / acre)	9.77 lbs / 1000 sq ft (0.42 lbs AI / acre)
Transition	4.83 lbs / 1000 sq ft (0.20 lbs AI / acre)	6.33 lbs / 1000 sq ft (0.27 lbs AI / acre)	13.3 lbs / 1000 sq ft (0.58 lbs AI / acre)
South	6.33 lbs / 1000 sq ft (0.27 lbs AI / acre)	9.77 lbs / 1000 sq ft (0.42 lbs AI / acre)	13.3 lbs / 1000 sq ft (0.58 lbs AI / acre)
Coastal South	9.77 lbs / 1000 sq ft (0.42 lbs AI / acre)	13.3 lbs / 1000 sq ft (0.58 lbs AI / acre)	16.83 lbs / 1000 sq ft (0.74 lbs AI / acre)

North: All other areas.
 Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, and coastal areas of CT, NY and RI.
 South: AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, and TX.
 Coastal South: FL, southern coastal areas of AL, GA, LA, MS, NC, SC, TX.

*Apply to areas where crabgrass is a major problem.
 **Apply to areas where crabgrass is a major problem.

SPRING APPLICATIONS: For single applications in the spring or early summer, use the appropriate rate from "Application Rates" table (depending on user's location, mowing height, pre- or post-emergence use). The duration of residual weed control is directly related to the total rate applied, but varies depending on soil conditions. This product may be applied to areas where turf is cut high (from lawns) for 3 to 6 months of crabgrass control.

FALL APPLICATIONS: This product may be applied in the fall (late August through November) at the "Program 2" rates for crabgrass control. This product may also be applied in the fall (late August through November) at the "Program 2" rates for crabgrass control. Do not apply more than 13.3 pounds per 1000 square feet (0.5 pounds AI per acre) per application. Do not apply more than 4.0 pounds per 1000 square feet (0.17 pounds AI per acre) per year. (In New York, do not apply more than 13.3 lbs per 1000 square feet per year.)

USE PRECAUTIONS: Do not apply more than 13.3 pounds per 1000 square feet (0.5 pounds AI per acre) per application. Do not apply more than 4.0 pounds per 1000 square feet (0.17 pounds AI per acre) per year. (In New York, do not apply more than 13.3 lbs per 1000 square feet per year.)

CULTURAL PRACTICES: Such as aeration and de-thatching should be completed, and the turf allowed to recover before application of this product. Do not use clippings from treated turf for animal feed or for mulching around vegetables or fruit trees. Do not allow domestic animals to feed or forage on treated turf. Keep dogs and cats off of treated areas until the turf has recovered.

POST-EMERGENCE CONTROL: Do not use on golf course putting greens. Do not use on lawns where mowing and delay mowing again for a few days after the application. Irrigation or rainfall within 7 days after application may improve weed control. Use of split (fall rate) applications spaced 6-8 weeks apart may provide improved weed control.

RESEEDING, OVERSEEDING, DR SPRIGGING: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When seeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. For best results, apply this product to areas where turf is cut high (from lawns) for 3 to 6 months of crabgrass control.

CONTROL OF OTHER GRASSES AND BROADLEAF WEEDS: This product is not intended for control of other grasses or broadleaf weeds.

SPRING APPLICATIONS: Used as directed for crabgrass control in spring, this product will also control or suppress the following weeds when applied prior to their emergence: Bentsgrass, Bluegrass (annual, roughstalk), Crowfootgrass, Foxtail (yellow & green), Goosegrass, Knotgrass, Lespedeza (common), Oxalis (bulbous, creeping & yellow woodsoot), Purslane (common), Ryegrass (annual & perennial), Smutgrass, Speedwell (slow), Spurge (prostrate and spined).

FALL APPLICATIONS: Used as directed in late summer or early fall, for crabgrass control through the early part of the next spring, this product will also control or suppress the following weeds when applied prior to their emergence: Bluegrass (annual, roughstalk), Buttercrunch, Chickweed, Geranium (Carolina), Henbit, Parsley-piert, Pigeonweed, Shepherds purse.

SPRINKLING: Do not apply to areas where turf is cut high (from lawns) for 3 to 6 months of crabgrass control.

SETTINGS: Settings shown are approximate for new equipment. Your spreader may need calibration, due to differences in terrain, walking speed and equipment condition. Calibrate your spreader to ensure proper coverage. Clean & lubricate spreader after use.

SAFETY: Do not use on golf course putting greens. Do not use on lawns where mowing and delay mowing again for a few days after the application. Irrigation or rainfall within 7 days after application may improve weed control. Use of split (fall rate) applications spaced 6-8 weeks apart may provide improved weed control.

RESEEDING, OVERSEEDING, DR SPRIGGING: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When seeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. For best results, apply this product to areas where turf is cut high (from lawns) for 3 to 6 months of crabgrass control.

CONTROL OF OTHER GRASSES AND BROADLEAF WEEDS: This product is not intended for control of other grasses or broadleaf weeds.

SPRING APPLICATIONS: Used as directed for crabgrass control in spring, this product will also control or suppress the following weeds when applied prior to their emergence: Bentsgrass, Bluegrass (annual, roughstalk), Crowfootgrass, Foxtail (yellow & green), Goosegrass, Knotgrass, Lespedeza (common), Oxalis (bulbous, creeping & yellow woodsoot), Purslane (common), Ryegrass (annual & perennial), Smutgrass, Speedwell (slow), Spurge (prostrate and spined).

FALL APPLICATIONS: Used as directed in late summer or early fall, for crabgrass control through the early part of the next spring, this product will also control or suppress the following weeds when applied prior to their emergence: Bluegrass (annual, roughstalk), Buttercrunch, Chickweed, Geranium (Carolina), Henbit, Parsley-piert, Pigeonweed, Shepherds purse.

SPRINKLING: Do not apply to areas where turf is cut high (from lawns) for 3 to 6 months of crabgrass control.

SETTINGS: Settings shown are approximate for new equipment. Your spreader may need calibration, due to differences in terrain, walking speed and equipment condition. Calibrate your spreader to ensure proper coverage. Clean & lubricate spreader after use.

SAFETY: Do not use on golf course putting greens. Do not use on lawns where mowing and delay mowing again for a few days after the application. Irrigation or rainfall within 7 days after application may improve weed control. Use of split (fall rate) applications spaced 6-8 weeks apart may provide improved weed control.

RESEEDING, OVERSEEDING, DR SPRIGGING: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When seeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. For best results, apply this product to areas where turf is cut high (from lawns) for 3 to 6 months of crabgrass control.

CONTROL OF OTHER GRASSES AND BROADLEAF WEEDS: This product is not intended for control of other grasses or broadleaf weeds.

SPRING APPLICATIONS: Used as directed for crabgrass control in spring, this product will also control or suppress the following weeds when applied prior to their emergence: Bentsgrass, Bluegrass (annual, roughstalk), Crowfootgrass, Foxtail (yellow & green), Goosegrass, Knotgrass, Lespedeza (common), Oxalis (bulbous, creeping & yellow woodsoot), Purslane (common), Ryegrass (annual & perennial), Smutgrass, Speedwell (slow), Spurge (prostrate and spined).

FALL APPLICATIONS: Used as directed in late summer or early fall, for crabgrass control through the early part of the next spring, this product will also control or suppress the following weeds when applied prior to their emergence: Bluegrass (annual, roughstalk), Buttercrunch, Chickweed, Geranium (Carolina), Henbit, Parsley-piert, Pigeonweed, Shepherds purse.

SPRINKLING: Do not apply to areas where turf is cut high (from lawns) for 3 to 6 months of crabgrass control.

SETTINGS: Settings shown are approximate for new equipment. Your spreader may need calibration, due to differences in terrain, walking speed and equipment condition. Calibrate your spreader to ensure proper coverage. Clean & lubricate spreader after use.

SAFETY: Do not use on golf course putting greens. Do not use on lawns where mowing and delay mowing again for a few days after the application. Irrigation or rainfall within 7 days after application may improve weed control. Use of split (fall rate) applications spaced 6-8 weeks apart may provide improved weed control.

RESEEDING, OVERSEEDING, DR SPRIGGING: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When seeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. For best results, apply this product to areas where turf is cut high (from lawns) for 3 to 6 months of crabgrass control.

CONTROL OF OTHER GRASSES AND BROADLEAF WEEDS: This product is not intended for control of other grasses or broadleaf weeds.

SPRING APPLICATIONS: Used as directed for crabgrass control in spring, this product will also control or suppress the following weeds when applied prior to their emergence: Bentsgrass, Bluegrass (annual, roughstalk), Crowfootgrass, Foxtail (yellow & green), Goosegrass, Knotgrass, Lespedeza (common), Oxalis (bulbous, creeping & yellow woodsoot), Purslane (common), Ryegrass (annual & perennial), Smutgrass, Speedwell (slow), Spurge (prostrate and spined).

FALL APPLICATIONS: Used as directed in late summer or early fall, for crabgrass control through the early part of the next spring, this product will also control or suppress the following weeds when applied prior to their emergence: Bluegrass (annual, roughstalk), Buttercrunch, Chickweed, Geranium (Carolina), Henbit, Parsley-piert, Pigeonweed, Shepherds purse.

SPRINKLING: Do not apply to areas where turf is cut high (from lawns) for 3 to 6 months of crabgrass control.

SETTINGS: Settings shown are approximate for new equipment. Your spreader may need calibration, due to differences in terrain, walking speed and equipment condition. Calibrate your spreader to ensure proper coverage. Clean & lubricate spreader after use.

SAFETY: Do not use on golf course putting greens. Do not use on lawns where mowing and delay mowing again for a few days after the application. Irrigation or rainfall within 7 days after application may improve weed control. Use of split (fall rate) applications spaced 6-8 weeks apart may provide improved weed control.

RESEEDING, OVERSEEDING, DR SPRIGGING: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When seeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. For best results, apply this product to areas where turf is cut high (from lawns) for 3 to 6 months of crabgrass control.

CONTROL OF OTHER GRASSES AND BROADLEAF WEEDS: This product is not intended for control of other grasses or broadleaf weeds.

SPRING APPLICATIONS: Used as directed for crabgrass control in spring, this product will also control or suppress the following weeds when applied prior to their emergence: Bentsgrass, Bluegrass (annual, roughstalk), Crowfootgrass, Foxtail (yellow & green), Goosegrass, Knotgrass, Lespedeza (common), Oxalis (bulbous, creeping & yellow woodsoot), Purslane (common), Ryegrass (annual & perennial), Smutgrass, Speedwell (slow), Spurge (prostrate and spined).

FALL APPLICATIONS: Used as directed in late summer or early fall, for crabgrass control through the early part of the next spring, this product will also control or suppress the following weeds when applied prior to their emergence: Bluegrass (annual, roughstalk), Buttercrunch, Chickweed, Geranium (Carolina), Henbit, Parsley-piert, Pigeonweed, Shepherds purse.

SPRINKLING: Do not apply to areas where turf is cut high (from lawns) for 3 to 6 months of crabgrass control.

SETTINGS: Settings shown are approximate for new equipment. Your spreader may need calibration, due to differences in terrain, walking speed and equipment condition. Calibrate your spreader to ensure proper coverage. Clean & lubricate spreader after use.

SAFETY: Do not use on golf course putting greens. Do not use on lawns where mowing and delay mowing again for a few days after the application. Irrigation or rainfall within 7 days after application may improve weed control. Use of split (fall rate) applications spaced 6-8 weeks apart may provide improved weed control.

RESEEDING, OVERSEEDING, DR SPRIGGING: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When seeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. For best results, apply this product to areas where turf is cut high (from lawns) for 3 to 6 months of crabgrass control.

CONTROL OF OTHER GRASSES AND BROADLEAF WEEDS: This product is not intended for control of other grasses or broadleaf weeds.

SPRING APPLICATIONS: Used as directed for crabgrass control in spring, this product will also control or suppress the following weeds when applied prior to their emergence: Bentsgrass, Bluegrass (annual, roughstalk), Crowfootgrass, Foxtail (yellow & green), Goosegrass, Knotgrass, Lespedeza (common), Oxalis (bulbous, creeping & yellow woodsoot), Purslane (common), Ryegrass (annual & perennial), Smutgrass, Speedwell (slow), Spurge (prostrate and spined).

FALL APPLICATIONS: Used as directed in late summer or early fall, for crabgrass control through the early part of the next spring, this product will also control or suppress the following weeds when applied prior to their emergence: Bluegrass (annual, roughstalk), Buttercrunch, Chickweed, Geranium (Carolina), Henbit, Parsley-piert, Pigeonweed, Shepherds purse.

SPRINKLING: Do not apply to areas where turf is cut high (from lawns) for 3 to 6 months of crabgrass control.

SETTINGS: Settings shown are approximate for new equipment. Your spreader may need calibration, due to differences in terrain, walking speed and equipment condition. Calibrate your spreader to ensure proper coverage. Clean & lubricate spreader after use.

SAFETY: Do not use on golf course putting greens. Do not use on lawns where mowing and delay mowing again for a few days after the application. Irrigation or rainfall within 7 days after application may improve weed control. Use of split (fall rate) applications spaced 6-8 weeks apart may provide improved weed control.

RESEEDING, OVERSEEDING, DR SPRIGGING: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When seeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. For best results, apply this product to areas where turf is cut high (from lawns) for 3 to 6 months of crabgrass control.

CONTROL OF OTHER GRASSES AND BROADLEAF WEEDS: This product is not intended for control of other grasses or broadleaf weeds.



PREVENTS CRABGRASS PLUS with 0.086% DITHIOPYR

SELECTIVE HERBICIDE PLUS FERTILIZER FOR PREEMERGENCE CONTROL OF LISTED ANNUAL GRASSES AND BROADLEAF WEEDS IN ESTABLISHED LAWNS & ORNAMENTAL TURF

ACTIVE INGREDIENT: Dithiopyr: 3.5-pyridinedicarboethoic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6- (trifluoromethyl)-S-S-dimethyl ester ... 0.086% INERT INGREDIENTS: ... 99.914% TOTAL: ... 100.000%

EPA Reg. No. 32802-68 EPA Est. No. 59144-GA-1(A), 32802-MO-1(B), 37351-CO-1(C), 8378-IN-1 (D), 9404-FL-1 (E)

Read the entire "Directions for Use" and "Warranty Statement" before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: Irritant. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift or runoff may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

Table with safety instructions: If in eyes, If on skin, If on clothing, If on hands, If on shoes, If on face, If on mouth.

GUARANTEED ANALYSIS

Directions for Use: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use in Nassau or Suffolk Counties. PRODUCT INFORMATION: This product contains turf fertilizer plus a selective herbicide for control of crabgrass and other annual grasses and broadleaf weeds...

NOTIFICATION

NOV 30 2012

Net Weight 50 lbs (22.70 kg) APPLICATION RATES FOR Crabgrass Control: TABLE with columns for REGION, PROGRAM 1, PROGRAM 2, and PREEMERGENCE USE.

SPRING APPLICATIONS: For single applications in the spring or early summer, use the appropriate rate from "Application Rates" table (depending on user's location, mowing height, pre- or post-emergence use). The duration of residual weed control is directly related to the total rate applied...

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store in original container in a locked storage area out of reach of children and domestic animals. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Information regarding the contents and level of metals: Information on the Internet at http://www.aafco.org/metals.htm

Manufactured for: Howard Johnson's Enterprises, Inc. 9875 S 60th Street, Franklin, WI 53132

WHERE TO USE: This product may be used on the following established turfgrasses: Creeping Bentgrass (Agrostis perennis), Kentucky Bluegrass (Poa pratensis), Fine Fescue** (Festuca arvensis), Tall Fescue (Festuca arvensis), Perennial Ryegrass (Lolium perenne), Annual Ryegrass (Lolium multiflorum), Bermudagrass (Cynodon dactylon), Buffalograss** (Buchloe dactyloides), Carpetgrass (Axonopus compressus), Centipedegrass (Eriochloa ophiuroides), Kikuyugrass (Pennisetum clandestinum), St. Augustinegrass (Stenotaphrum secundatum), Zoysia (Zoysia tenuifolia).

Use of this product on turf that is not well established, or weakened by weather, pests or other stress, may increase the chance of crabgrass from the ground. This product provides pre-emergence control of crabgrass (large, smooth, and southern species) when applied prior to emergence of crabgrass from the ground. In addition, this product provides "early post-emergence" control during the early stages of growth after crabgrass emerges from the ground. The early post-emergence control is limited, and post-emergence applications are only effective if applied before crabgrass has displayed its fifth leaf or first tiller. The benefit of this post-emergence activity is that it gives the user a 2 to 4 week longer period of time to make applications for effective control of crabgrass.

WHERE TO USE: This product may be used on the following established turfgrasses: Creeping Bentgrass (Agrostis perennis), Kentucky Bluegrass (Poa pratensis), Fine Fescue** (Festuca arvensis), Tall Fescue (Festuca arvensis), Perennial Ryegrass (Lolium perenne), Annual Ryegrass (Lolium multiflorum), Bermudagrass (Cynodon dactylon), Buffalograss** (Buchloe dactyloides), Carpetgrass (Axonopus compressus), Centipedegrass (Eriochloa ophiuroides), Kikuyugrass (Pennisetum clandestinum), St. Augustinegrass (Stenotaphrum secundatum), Zoysia (Zoysia tenuifolia).

Use of this product on turf that is not well established, or weakened by weather, pests or other stress, may increase the chance of crabgrass from the ground. This product provides pre-emergence control of crabgrass (large, smooth, and southern species) when applied prior to emergence of crabgrass from the ground. In addition, this product provides "early post-emergence" control during the early stages of growth after crabgrass emerges from the ground. The early post-emergence control is limited, and post-emergence applications are only effective if applied before crabgrass has displayed its fifth leaf or first tiller. The benefit of this post-emergence activity is that it gives the user a 2 to 4 week longer period of time to make applications for effective control of crabgrass.

SPRING APPLICATIONS: For single applications in the spring or early summer, use the appropriate rate from "Application Rates" table (depending on user's location, mowing height, pre- or post-emergence use). The duration of residual weed control is directly related to the total rate applied...

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store in original container in a locked storage area out of reach of children and domestic animals. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Information regarding the contents and level of metals: Information on the Internet at http://www.aafco.org/metals.htm

Manufactured for: Howard Johnson's Enterprises, Inc. 9875 S 60th Street, Franklin, WI 53132



20-0-5

PREVENTS CRABGRASS PLUS with 0.086% DIMENSION®

SELECTIVE HERBICIDE PLUS FERTILIZER FOR PREEMERGENCE CONTROL OF LISTED ANNUAL GRASSES AND BROADLEAF WEEDS IN ESTABLISHED LAWNS & ORNAMENTAL TURF

ACTIVE INGREDIENT:

- Dithiopyr, 3,5-pyridinedicarbonyl acid, 2-(difluoro...)

NET WEIGHT 40 lbs (18.14 kg)

GUARANTEED ANALYSIS: Total Nitrogen (N) ... 18.83% Urea Nitrogen ... Soluble Potash (K2O) ...

Derived From Muriate of Potash, Polymer Coated Urea, Urea.

*5.00% Solvoly Available Nitrogen From Polymer Coated Urea

Chlorine(Cl), not more than 6.00%

FGP60469 F315

EPA Reg. No. 32802-68

EPA Est. No. 59144-GA-1(A), 32802-MO-1(B), 37351-CO-1 (C)

First sale of this case above label date when manufactured.

DIMENSION® is a trademark of Dow AgroSciences

Read the entire "Directions for Use" and "Warning Statement" before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. In the State of New York: For use by professional turf applicators only, at no more than 13.3 lbs per 1000 square feet per year (0.5 pounds AI per acre per year). Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

For use on established lawns and ornamental turf. Apply this product only to seeded, sodded, or sodged turf that is well established. Turf must develop a good root system and uniform stand, and have received at least two mowings following seeding, sodding or sodding before making the first application of this product. Use of this product on turf that is not well established, or weakened by weather, pests or other stress, may increase the chance of turf injury.

This product provides pre-emergence control of crabgrass (large, smooth, and southern species) when applied prior to emergence of crabgrass from the ground. In addition, this product provides pre-emergence control of crabgrass (large, smooth, and southern species) when applied prior to emergence of crabgrass from the ground.

Table with 4 columns: REGION, APPLICATION RATES for Crabgrass Control, PREEMERGENCE USE (high turf), PREEMERGENCE USE (low turf), POST-EMERGENCE USE. Rows include North, Transition, South, Coastal South.

SPRING APPLICATIONS: For single applications in the spring or early summer, use the appropriate rate from "Application Rates" table (depending on user's location, mowing height, pre- or post-emergence use). The duration of residual weed control is directly related to the total rate applied, but varies depending on turfgrass species and soil conditions.

FALL APPLICATIONS: Used as directed in late summer or early fall, for crabgrass control through the early part of the next spring. This product will also control or suppress the following weeds when applied prior to their emergence: Bluegrass (annual, roughstalk), Bilberry, Chickweed, Geranium (Carolina), Henbit, Parsley-plant, Pineappleweed, Shepherd's purse.

Table with 2 columns: Spreader setting to apply at the listed rate (lb per 1000 sq ft) and Spreader. Rows include Accu-Pro 2000, Laco 021820, Stylax 76-22, PSB Bigfoot, Lely HR, Vicron 200.

The directions for use of this product are believed to be adequate and must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, inefficacy, or other unintended consequences may result due to such factors as weather conditions, presence or absence of other materials, soil conditions, etc.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

NOTIFICATION

NOV 30 2012

Dimension 0.086 Professional PP (06-15-2010A)10-35-12

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: Harmful if inhaled. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store in original container in a locked storage area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Manufactured by: Howard Johnson's Enterprises, Inc. 8675 S 60th Street, Franklin, WI 53132

Information regarding the contents and levels of metals in this product is available on the internet at http://www.waspico.org/metal.htm

Table with 2 columns: First Aid and First Aid. Rows include If in eye, If inhaled, If on skin or clothing, If swallowed, and Use this product only.