



### UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

Ms. Janet P. Baguhn Howard Johnson's Enterprises, Inc. 700 West Virginia Street #222 Milwaukee, WI 53204

DEC - 9 2010

Subject:

Label Amendment

Request Alternate Brand Name: "Howard Johnson's Prevents Crabgrass Plus with

0.107% Dithiopyr"

Product Name: Dithiopyr 0.107 Plus

EPA Reg. No. 32802-71

Application Dated: October 5, 2010

Dear Ms. Baguhn,

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable, provided you make the following change:

1. The "GENERAL INFORMATION" header must be replaced with "PRODUCT INFORMATION".

A stamped copy of your label is enclosed for your records. This label supersedes all previously accepted labels. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after eighteen (18) months from the date of this letter or the next printing of the label, whichever occurs first, must bear the new revised label. If this condition is not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions, please call me directly at (703) 305-1243.

Sincerely,

Kathryn V. Montague, Product Manager 23

Herbicide Branch

Registration Division (7505P)



# **PLUS with 0.107%** PREVENTS CRABGRASS DITHIOPYR

PREEMERGENCE CONTROL OF LISTED ANNUAL GRASSES AND BROADLEAF WEEDS IN ESTABLISHED LAWNS & ORNAMENTAL TURF CONTAINS A COMPLETE LAWN FOOD ACTIVE INGREDIENT. SELECTIVE HERBICIDE PLUS FERTILIZER FOR

Dithiopyr. 3,5-pyridinedicarbothiolic acid, 2-(diffuo romethyr)-4(2-methylpropyr)-6-(irifuoromethyr)-5, S-dimethyl ester 0.107%.

INERT INGREDIENTS: 99.883%.
TOTAL.

EPA Reg. No. 32802-71 EPA Est. No. 32802-MO-1 (B) First letter of bag code shows establish Read the entire "Directions for Use" and "Warranty Statement" before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

## with COMMENTS n EPA Letter Dated: Dec-9 2010 ACCEPTED

Fungicide, and Rodemicide Act registered under EPA Reg. No. Under the Federal In sicide, as amended, for the pesticide

33803-71

(high cut turigrass)

2.68 lbs / 1000 sq ft (117 lbs / acre 0.125 lbs Al / acre) 3.93 lbs / 1000 sq ft (171 lbs / acre 0.183 lbs Al / acre)

DRECTIONS FOR USE: It is a violation of Federal law to use this product in a marrier inconsistent with its labeling.
In the State of New York, For use by prosticional transfer and the state of the sta

iagrass (Zoysia japonica). emergence control of crabgases (large, smooth, and southern species) when applied prior to emergence of crabgases from the ground-voicides "any large control of crabgases pains" and a species of the crabgases are required to the crabgases strong the crabgases and explored. This sent post-hords to lime when you can first see the ordinary beautiful to the control of the crabgases has displayed its fifth heat of first lime hands to lime when you can first see the ordinary as the crabgases hand in the laws.) The benefit of this post-emergence activity is that it gives the period off time when you can first see the ordinary in the lawn.) The benefit of this post-emergence activity is that it gives the period off time when you can first see the ordinary in the lawn.) The benefit of this post-emergence activity is that it gives the period off time when you can first see that the control of crabgases. The certified periods see full control or such and the control of crabgases are control or charges and the control of crabgases and the control of crabgases (Liciam persons). Beliefs that the control or ophituoides), durishing your distributions (Eremochioa ophituoides).

sotis tenuis) varieties.

(into to this product. Atlanta, Banner, Beauty, Bilgant, CF-2, and Waldins. Not all varieties of fescue have been tested the fungrass is fully green and has established new roots. APPLICATION RATES for Crab

SPRING APPLICATIONS. For single applications in the spring or early summer, use the apportions the from Application Rates\* table (depending on user's location, moning bright, the ropode-morphic place in location, moning bright), the ropode-morphic place is considered to reside a possible of the properties of the state applied. But values depending on wester, were pressure and location, the Program 1' for preemergence control on assess where tart is call their location. Use Program and obtains a possible or the program should not be used for preemergence control. This program should not be used for professionary and and a seal of the state of the provious year. This program provides 4 to 6 months of preemergence control where tart is cut to wild profession provides 4 to 6 months of preemergence control of crabigosas. This program provides 4 to 6 months of preemergence control of crabigosas. This product may be posterinegence control of crabigosas. This product may be supplied as a single application or as a disk posteriation for capital date August through Novemberl at the Program 2' rate for crabigosas control in the spring, attention or control through the and program 2' rate for crabigosas control in the spring, attention or control through the and program 2' rate for crabigosas control through the and program 2' rate for crabigosas control through the service of the program 2' rate for crabigosas control through the service of the program 2' rate for crabigosas control through the service of the crabigosas desiring applications must be followed by an appropriate spring application must be selected. PREEMERGENCE USE (low cut turigrass) POSTEMERGENCE USE 5.37 lbs / 1000 sq ft (234 lbs / acre 0.25 lbs Al/acre) 7.85 lbs / 1000 sq ft (342 lbs / acre 0.366 lb Al / acre) 10.74 lbs / 1000 sq ft (468 lbs / acre 0.50 lbs / acre)

7.85 lbs / 1000 sq ft (342 lbs / acre 0.366 lb Al / acre) 5.37 lbs / 1000 sq ft (234 lbs / acre 0.25 lbs Al / acre)

<u>USE PRECAUTIONS</u>: Do not apply more than 10.74 pounds get 1000 aquaine idea (15, pounds 4) per acres per application. Do not apply more than 32.2 pounds per 1000 square feet (1,5 pounds Al per acre) per year. (In New York, do not apply more than 10.74 bis per 1000 aquare (in Keet per year.)

if per year.) The product of this product. Do not use recover before application of this product. The per domestic animals to feed or forage on treated furf. Keep ntr. AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN and TX statal South: HI, FL, southern coastal areas of AL, GA, LA, MS, NC, SC, TX y require split or sequential applications for full control.

Official particles such as wifficultion and dubtaching about be completed, and the furt allowed to recover before application of this product. Do not use options to stand the further assimilated of formationing around regardless or full treat. On not allow demostic animals to feed or forage on treated turk. Keep proper and passed turk for an interest of this sential. On not use on golf course purifying great minimals to feed or forage on treated turk. Keep proper and the sential countries are as until a sent to be a new to a sent the sent of the sential countries are treated are as until a sport with any product whitm it we days after the application are produced minimal applications appreciately as a feet by a spiritual product minimal to a single figure applications appreciately as the treated are as the spiritual product within the polycement.

pproved veed cale. OR SPRIGGING. Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may like. <u>OVERSEEDING, OR SPRIGGING, OR SPRIGGING</u> or destinated infiguresses. When seeding, follow good cultural practices such as coil cultivation, imgation and fertilization. For best

BROADLEAF WEEDS:

SPRIGAPLICATIONS. Used as directed for craigrass control in spring, this product will also control or suppress the following needs when applied prior to their emergeners. Barnyadayass, Blaquase formula from their crand (yellow & green). To coapeas, Klaupagas, Especial common), Assall (pullerops, respiral & yellow woodsorral), Pratient (common), Regards (smula & perennial), Smulgass, Specified (com), Spurge (prostate and vial is also controlled at the program 2 rates.

APPLICATIONS. Used as directed in late summer or early fall, for cratigoras control through the early part of the rext spring, this product will also control announces the following veeds when applied prior to their emergence: Bluegrass (arrual, troughtails), Bittercess, Chickweed, Gerantum (Carolina), Herbib.

STORAGE AND DISPOSAL

spreader setting to apply at the listed rate (lbs 0.25 Closed Valency and entire the product are believed to be adequate and must be followed carefully, it is impossible to eliminate all ricks inherently associated with the use of interference, and the scale of the product and an adequate and must be followed carefully in the scale of the

information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm

# HAZARDS TO HUMANS AND DOMESTIC ANIMALS PRECAUTIONARY STATEMENTS

KEEP OUT OF REACH OF CHILDREN

CAUTION: Causes eye intrition. Harmful if inhafed, Avoid contact with eyes or clothing, Avoid breathing dust. Weah thoroughly with soap and water after handling. Remove contaminated clothing and weah before reuse. Protonged or frequently repeated sith contact may cause altergic neadron in some intrivious. ENVENDIBLETICE\_LASARDS. This peation is found in both and highly toxic to other aquatic organisms including opsters and shiring. Use with care ENVENDIBLETICE\_LASARDS. This peation is found in any body of water. Dint or mort may adversely after squalic organisms in adjacent against sites, Do not apply when waters and peating and the protonger or to interrube ansatz below the mann high water mark. Do not apply when weather conditions favor duff from treated areas. Do not containments water when disposing of equipment waterwater.

inated clothing. ediately with plenty of water for 15-20 First Aid on and rinse slowly and gently with water son control center or physician immediately ing, call 911 or an ambulance, contact lenses, if present, after the first 5 , then continue rinsing eye. oison control center or doctor for treatment solson control center or doctor for treatment finhaled

of reach of children and domestic animals.

PESITICED DISPOSAL. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal health

CONTAINER DISPOSAL. Nonrelliable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling the available or dispose of empty bag in a sanitary landfill or by incineration of, if allowed by State and local authorities, by burning. If burned, stay out of smoke. Do not contaminate water, food or feed by storage or disposal. STORAGE: Store in original container in a locked storage area out Manufactured by: Howard Johnson's Enterprises, Inc. 700 W. Virginia St., No. 222 Milwaukee, W 53204 control center or physician.

Do not give anything by mouth to an unconscious person.

Hot line number: Have the product container or control center or doctor, or go pesticide product (including the pesticide incidents), call the Network at 1-800-858-7378.