

PLEASE NOTE

**This image contains more than one label
approved for this product on this date.**

Please read instructions on reverse before completing form. Form Approved. OMB No. 2070-0060. Approval expires 2-28-



United States
Environmental Protection Agency
Washington, DC 20460

Registration
Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 32802-73	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Turf Food with Crabgrass Preventer	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Howard Johnson's Enterprises, Inc. 700 West Virginia Street #222, Milwaukee, WI 53204 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

NOTIFICATION Section - II

<input type="checkbox"/> Amendment - Explain below.	JUN 26 2006	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____		<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.		<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Additional Brand Name: Gertens Turf Special Crabgrass Preventer

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container:	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
				<input checked="" type="checkbox"/> Paper	<input type="checkbox"/> Other (Specify) _____
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 20		5. Location of Label Directions E 1	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Janet P. Baguhn	Title Compliance Manager	Telephone No. (Include Area Code) 414-276-1505
2. Signature 		3. Title Compliance Manager
4. Typed Name Janet P. Baguhn		5. Date June 5, 2006
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)

SIG



NOTIFICATION

JUN 26 2006


Date: June 5, 2006

Product: Turf Food With Crabgrass Preventer
EPS Reg. No. 32802-73

Registrant: Howard Johnson's Enterprises, Inc.
700 W. Virginia Street
Milwaukee, WI 53204-1548

Action: Notification of Additional Brand Name

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152-46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and CFR 152-46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.



 Janet P. Baguhn
 Compliance Manager



June 6, 2006

Document Processing Desk (NOTIF)
Ms. Joanne I. Miller (PM 23)
Office of Pesticide Programs (7405P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501


Dear Ms. Miller:

Re: Notification of Additional Brand Name
Turf Food with Crabgrass Preventer
EPA Reg. No. 32802-73

Enclosed please find three applications for Notification of Additional Brand Name to the above-referenced pesticide registration.

Thank you for your assistance. Please call me if you have any questions:
414-276-1505, ext. 104.

Sincerely,



Janet P. Baguhn

Enclosures: Notifications of Additional Brand Name (EPA Form 8570-1)
Copy of Product Labels
Certifications

NOTIFICATION
JUN 26 2006

Crabgrass



Gertens
Turf Special
www.gertens.com

Preventer and
lawn fertilizer for
better crabgrass
control. Contains
Dimension®!

24·0·12




P H O S P H A T E F R E E

EPA Reg. No. 82802-73
EPA Est. No. 82802-WI-1 (A),
82802-MO-1 (B)
First letter of code on bottom tape
indicates the manufacturing plant.

ACTIVE INGREDIENTS:
Dithiopyr: 3,5-pyridinedicarboxylic acid,
2-(4-fluoromethyl)-4-(2-methylpropyl)-6-
(trifluoromethyl)-S,S-dimethyl ester 0.06%
INERT INGREDIENTS: 99.94%
Total 100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION: See back panel for further cautions,
first aid information and directions for use.

Covers 5,000 sq. ft. • Net Weight: 20 lbs. (9.09 kg)



Guaranteed Analysis		
24	0	12
a) Nitrogen (N).....		24%
24.00% Urea Nitrogen*		
soluble Potash (K ₂ O).....		12%
Total Sulfur (S).....		2.08%
2.08% Free Sulfur.....		
Total Iron (Fe).....		1.00%
10% Water Soluble Iron (Fe)		
Derived from: Muriate of Potash, Polymer Coated-Sulfur Coated Urea, Ferric Oxide, Ferrous Sulfate.		
* 4.80% Slowly available Urea Nitrogen from Polymer Coated-Sulfur Coated Urea.		
Chlorine (Cl), not more than 5%		

Special Care

What

Where

When

How

Precautionary Statements

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and highly toxic to other aquatic organisms including crayfish and shrimp. Use with care when applying to turf areas adjacent to any body of water. Use of this product may adversely affect aquatic organisms in adjacent aquatic areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or physician immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or physician. Do not give anything by mouth to an unconscious person.

Not for use: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product, including health concerns, medical emergencies, or pesticide incidents, call the National Pesticide Information Center at 1-800-858-7378.

General Information

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product controls crabgrass and other annual grasses and broadleaf weeds, and also contains fertilizer for proper turf growth in established lawns. It can be applied 2 to 4 weeks later than other crabgrass preventers, and provides up to 3 to 5 months control of crabgrass and other weed grasses and broadleaf weeds in home lawns. Apply only to lawns that are well established. Use of this product on lawns that are not well established, or weakened by weather, pests or mechanical stress, may increase the chance for turf injury. This product will not harm nearby established ornamentals when used according to label directions. It can be applied according to these directions to the following turfgrasses:

Cool season grasses: creeping bentgrass, Kentucky bluegrass, fine fescue, tall fescue, perennial ryegrass; Warm season grasses: bahiagrass, bermudagrass, buffalograss, carpetgrass, centipedegrass, kikuygrass, St. Augustinegrass, zoysiagrass.

This product will control (at post-emergence rates) or suppress (at pre-emergence rates) these weed grasses and broadleaf weeds when applied before weeds emerge in lawns: crabgrass, annual bluegrass (Poa annua), barnyardgrass, crowfootgrass, foxtail (yellow & green)*, goosegrass, kikuygrass, common lespedeza, buttercup oxalis, creeping oxalis, yellow woodsorrel, pheasantweed, purslane (common), ryegrass (annual & perennial), smartgrass (seedling), corn speedwell, prostrate spurge, spotted spurge, bittercress, chickweed, Carolina geranium, henbit, parsley-plant. *Foxtail is also controlled at the pre-emergence rates.

Application Directions

This product can be applied 2 to 4 weeks later than other crabgrass preventers, because Dimension® herbicide in this product provides preventive (pre-emergence) control when spread on lawns prior to germination of the weed seeds. It also provides post-emergence control of crabgrass when applied 2 to 4 weeks after crabgrass germination. Post-emergence control is limited to crabgrass only. Application of this product later than 4 weeks after crabgrass germination will result in reduced control of crabgrass. For control of the weed grasses and broadleaf weeds listed on this label (other than crabgrass), it must be applied as a preventative treatment before the weed seeds germinate. ONLY crabgrass can be controlled with a delayed application after germination.

This product can be applied in the spring, before weeds become established in the lawn, or in the fall for pre-emergence crabgrass control through the early part of the next spring. Fall applications should be followed by a second (sequential) application the following spring to provide season-long control. Do not apply more than 9.6 pounds per 1,000 square feet per year. Do not make more than one application of this product per year.

The length of time of weed control is directly related to the amount applied to the lawn. Use the application rates in Table 1 to provide 2 to 3 months of weed control. Use the application rates in Table 2 for 3 to 5 months of weed control. The actual length of control will vary depending on weather, weed pressure, and your location in the north or south region.

USE PRECAUTIONS: DO NOT apply more than 9.6 pounds of this product per 1000 square feet per year. DO NOT apply to flowers, vegetables or fruit trees. DO NOT use clippings from treated turf for mulching around flowers, vegetables or fruit trees. DO NOT apply this product to pastures. Keep people and pets off treated areas until dust has settled. Reseeding, overseeding, or application of treated areas within 4 months after applications of this product may inhibit the establishment of desirable turfgrasses. When seeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. For best results, use "sift seeders" designed to give good seed-to-soil contact. When soil testing or professional advice indicates the need for additional plant nutrients, make additional applications of fertilizer products over the growing season.

TIPS FOR IMPROVED CONTROL: For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. For improved weed control, water or irrigate treated areas within a few days after application. Aeration, raking or dethatching should be done before application of this product.


TABLE 1: RATES FOR 2 TO 3 MONTHS OF CRABGRASS CONTROL*

Region	Pre-emergence	Post-emergence
North	8.5 pounds per 1000 square feet	7.2 pounds per 1000 square feet
South	7.2 pounds per 1000 square feet	8.5 pounds per 1000 square feet

TABLE 2: RATES FOR 3 TO 5 MONTHS OF CRABGRASS CONTROL*

Region	Pre-emergence	Post-emergence
North	4.8 pounds per 1000 square feet	8.6 pounds per 1000 square feet
South	8.6 pounds per 1000 square feet	4.8 pounds per 1000 square feet

*Southern states include AL, AR, AZ, CA, FL, GA, HI, LA, MS, NC, NH, NY, OK, SC, TN, TX, VA, and southern areas of IL, IN and OH. All other states included as North Region.



Spreader Settings

SPREADER SETTINGS to apply at the listed pounds per 1000 square feet

	3.6 lbs	4.8 lbs	7.2 lbs	9.6 lbs
Ames (drop and rotary)	15	17	20	24
Earthway (drop and rotary)	15	17	20	24
Precision Red Devil	8	10	12	open
Cyclone SB3000	6.75	7.25	7.75	8.5
Scotts AccuGreen (drop)	7.5	10	13	15
Scotts EasyGreen (rotary)	28	29	30	31
Scotts SpeedyGreen (rotary)	7	8	11	16

Settings shown are approximate for new equipment. To insure proper coverage, ALWAYS CALIBRATE YOUR SPREADER BEFORE USE, due to differences in terrain, your walking speed and the condition of the equipment. Clean and lubricate the spreader immediately after each use.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Storage: Store in original container in a locked storage area out of reach of children and domestic animals. Pesticide disposal: Securely wrap original container in several layers of newspaper and discard in trash. Container disposal: Do not reuse bag. Fold empty bag and discard in trash.

Manufacturer warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label. Manufacturer makes no other warranty, express or implied, and shall not be liable for indirect or consequential damages.

Manufactured and Guaranteed By:



Howard Johnson's Enterprises, Inc.
700 W. Virginia St., Milwaukee, WI 53204



GG24001220C73PE
8503 C75-00880 01/01

NEXT

LABEL

Please read instructions on reverse before completing form. Form Approved OMB No. 2070-0060. Approval expires 2-28-

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration	OPP Identifier Number
		<input checked="" type="checkbox"/> Amendment	
		<input checked="" type="checkbox"/> Other	

Application for Pesticide - Section I

1. Company/Product Number 32802-73	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Turf Food with Crabgrass Preventer	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Howard Johnson's Enterprises, Inc. 700 West Virginia Street #222, Milwaukee, WI 53204		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Check if this is a new pesticide. **NOTIFICATION**

JUN 26 2006

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Additional Brand Name: Hummert's DYNA GREEN Crabgrass Control

Section - III

1. Material This Product Will Be Packaged in:				2. Type of Container:	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	<input type="checkbox"/> Glass	<input checked="" type="checkbox"/> Paper
		If "Yes" Package wgt.	No. per container	<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 20#		5. Location of Label Directions E 1	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Janet P. Baguhn	Title Compliance Manager	Telephone No. (Include Area Code) 414-276-1505
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		3. Date Application Received (Stamped)
2. Signature 	3. Title Compliance Manager	
4. Typed Name Janet P. Baguhn	5. Date June 5, 2006	



NOTIFICATION

JUN 26 2006


Date: June 5, 2006

Product: Turf Food With Crabgrass Preventer
EPA Reg. No. 32802-73

Registrant: Howard Johnson's Enterprises, Inc.
700 W. Virginia Street
Milwaukee, WI 53204-1548

Action: Notification of Additional Brand Name

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152-46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and CFR 152-46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.



Janet P. Baguhn
Compliance Manager

413

HUMMERT'S

NOTIFICATION
JUN 26 2006

DYNA GREEN™

CRABGRASS CONTROL

15.5-10-17.5 inches
with 3.5% Dimension

KEEP OUT OF REACH OF CHILDREN.
CAUTION: See back panel for further cautions,
first aid information and directions for use.

Net Wt. 20 lbs. (9.1 kg) - Free Flowing Conditioner Added

Manufactured by:
Howard Johnson's Enterprises, Inc.
700 W. Virginia St.
Milwaukee, WI 53204



CRABGRASS CONTROL WITH 0.06% DIMENSION

A SELECTIVE HERBICIDE THAT PROVIDES CONTROL OF LISTED ANNUAL GRASSES AND BROADLEAF WEEDS IN ESTABLISHED LAWNS & ORNAMENTAL TURF. CONTAINS A COMPLETE LAWN FOOD.

GUARANTEED ANALYSIS

Total Nitrogen (N)	15%
1.96% Ammoniacal Nitrogen	
13.04% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	5%
Soluble Potash (K ₂ O)	10%
Boron (B)	0.05%
Total Copper (Cu)	0.13%
0.03% Water Soluble Copper (Cu)	
Total Iron (Fe)	1.25%
0.03% Water Soluble Iron (Fe)	
Total Manganese (Mn)	0.13%
0.03% Water Soluble Manganese (Mn)	
Molybdenum (Mo)	0.0013%
Total Zinc (Zn)	0.13%
0.03% Water Soluble Zinc (Zn)	

Derived from: Ammonium Phosphate, Copper Oxide, Copper Sulfate, Iron Oxide, Iron Sulfate, Manganese Oxide, Manganese Sulfate, Sodium Borate, Sodium Molybdate, Zinc Oxide, Zinc Sulfate, Ferrous Sulfate, Muriate of Potash, Urea

Chlorine (Cl) not more than 6.00%
Florida License # F315
FGB95873

KEEP OUT OF REACH OF CHILDREN PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION:

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift or runoff may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

First Aid

If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> Call a poison control center or physician immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or physician. Do not give anything by mouth to an unconscious person.

Hot line number:

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for seed or other commercial use, or for commercial seed production, or for research purposes.

GENERAL INFORMATION: This product controls crabgrass and other annual grasses and broadleaf weeds, and also contains fertilizer for proper turf growth in established lawns. It can be applied 2 to 4 weeks later than other crabgrass preventers, and provides up to 3 to 5 months control of crabgrass and other weed grasses and broadleaf weeds in home lawns. It can be applied according to these directions to the following turfgrasses:

Cool season grasses: creeping bentgrass, Kentucky bluegrass, fine fescue, tall fescue, perennial ryegrass;

Warm season grasses: bahiagrass, bermudagrass, buffalograss, carpetgrass, centipedegrass, kikuyugrass, St. Augustinegrass, zoysiagrass.

Apply only to lawns that are well established. Use of this product on lawns that are not well established, or weakened by weather, pests or mechanical stress, may increase the chance for turf injury. This product will not harm nearby established ornamentals when used according to label directions.

When to apply: This product can be applied 2 to 4 weeks later than other crabgrass preventers, because Dimension® herbicide in this product provides preventive (pre-emergence) control when spread on lawns prior to germination of the weed seeds. It also provides post-emergence control of crabgrass when applied 2 to 4 weeks after crabgrass germination. Post-emergence control is limited to crabgrass only. Application of this product later than 4 weeks after crabgrass germination will result in reduced control of crabgrass. For control of the weed grasses and broadleaf weeds listed on this label (other than crabgrass), it must be applied as a preventative treatment before the weed seeds germinate. ONLY crabgrass can be controlled with delayed application after germination.

APPLICATION DIRECTIONS: This product can be applied in the spring, before weeds become established in the lawn, or in the fall for pre-emergence crabgrass control through the early part of the next spring. Fall applications should be followed by a second (sequential) application the following spring to provide season-long control. Do not apply more than 9.6 pounds per 1000 square feet per year. Do not make more than one application of this product per year.

LENGTH OF WEED CONTROL: The length of time of weed control is directly related to the amount applied to the lawn. Use the application rates in Table 1 to provide 2 to 3 months of weed control. Use the application rates in Table 2 for 3 to 5 months of weed control. The actual length of weed control will vary depending on weather, weed pressure, and your location in the north or south region.

USE PRECAUTIONS: DO NOT apply more than 9.6 pounds of this product per 1000 square feet per year. DO NOT apply to flowers, vegetables or fruit trees. DO NOT use clippings from treated turf for mulching around flowers, vegetables or fruit trees. DO NOT apply this product to pastures. Keep people and pets off treated areas until dust has settled. Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When seeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. For best results, use "slit seeders" designed to give good seed-to-soil contact.

TIPS FOR IMPROVED CONTROL: For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. For improved weed control, water or irrigate treated areas within a few days after application. Aeration, raking or dethatching of lawns should be done before application of this product.

WEEDS CONTROLLED: This product will control (at post-emergence rates) or suppress (at pre-emergence rates) these weed grasses and broadleaf weeds when applied before weeds emerge in lawns:

crabgrass, annual bluegrass (*Poa annua*), barnyardgrass, crowfootgrass, foxtail (yellow & green)*, goosegrass, kikuyugrass (seedling), common lespedeza, buttercup oxalis, creeping oxalis, yellow woodsorrel, pineappleweed, purslane (common), ryegrass (annual & perennial), smartgrass (seedling), corn speedwell, prostrate spurge, spotted spurge, bittercress, chickweed, Carolina geranium, henbit, parsley-plant.

*Foxtail is also controlled at the pre-emergence rates. When soil testing or professional advice indicates the need for additional plant nutrients, make additional applications of fertilizer products over the growing season.

This product may stain driveways, walks, patios, pools, etc. Wash or sweep off thoroughly immediately after each use.

NOTICE: Buyer and user assume all risks and liability of use, storage and/or handling of this product not in accordance with the directions for use on this label.

TABLE 1: RATES FOR 2 TO 3 MONTHS OF CRABGRASS CONTROL*

Region	Preemergence	Post emergence
North	3.6 pounds per 1000 square feet	7.2 pounds per 1000 square feet
South	7.2 pounds per 1000 square feet	9.6 pounds per 1000 square feet

TABLE 2: RATES FOR 3 TO 5 MONTHS OF CRABGRASS CONTROL*

Region	Preemergence	Post emergence
North	4.8 pounds per 1000 square feet	9.6 pounds per 1000 square feet
South	9.6 pounds per 1000 square feet	9.6 pounds per 1000 square feet

*Southern states include AL, AR, AZ, CA, FL, GA, HI, LA, MS, NC, NM, NV, OK, SC, TN, TX, VA, and southern areas of IL, IN and OH. All other states included as North Region.

SPREADER SETTINGS to apply at the listed pounds per 1000 sq. ft.

	3.6 lbs.	4.8 lbs.	7.2 lbs.	9.6 lbs.
Ames Drop or Rotary	15	17	21	24.5
EarthWay Drop or Rotary	15	17	21	24.5
Precision Red Devil Rotary	5	7	11	12
Republic EZ Drop	6.5	7.5	11	12
Republic EZ Rotary	6.5	7.5	11	12
Cyclone SB3000 Rotary	6.75	7.25	7.75	8.5
Scotts EasyGreen	28	29	31	29x2
EG-3 (rotary)				
Scotts SpeedyGreen (rotary)	9	11	16	18.5

Settings shown are approximate for new equipment. Your spreader may need calibration, due to differences in terrain, walking speed and equipment condition. Calibrate your spreader to ensure proper coverage. Clean & lubricate spreader after use.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in original container in a locked storage area out of reach of children and domestic animals. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Fold empty bag and discard in trash.

Manufactured by:

HOWARD JOHNSON'S ENTERPRISES, INC.
700 W. Virginia Street, Milwaukee, WI 53204-1548

NEXT

LABEL

32802-73

6/26/2006

Print Form

1/4

Please read instructions on reverse before using form. Form Approved OMB No. 2070-0060. Approval expires 2-28-



United States
Environmental Protection Agency
Washington, DC 20460

Registration Amendment

Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 32802-73	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Turf Food with Crabgrass Preventer	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Howard Johnson's Enterprises, Inc. 700 West Virginia Street #222, Milwaukee, WI 53204	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Check if this is a new address

NOTIFICATION

Section - II

JUN 26 2006

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Additional Brand Name: Gardens Beautiful Garden Centers Crabgrass Control Plus with 0.06% Dimension

Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input checked="" type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt No. per container

3. Location of Net Contents Information
 Label Container

4. Size(s) Retail Container
20# + 40#

5. Location of Label Directions
E 1

6. Manner in Which Label is Affixed to Product
 Lithograph
 Paper glued
 Stenciled Other _____

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Janet P. Baguhn	Title Compliance Manager	Telephone No. (include Area Code) 414-276-3505
-------------------------	-----------------------------	---

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature 	3. Title Compliance Manager	6. Date Application Received (Stamped)
4. Typed Name Janet P. Baguhn	5. Date June 5, 2006	

NOTIFICATION

JUN 26 2006


Date: June 5, 2006

Product: Turf Food With Crabgrass Preventer
EPS Reg. No. 32802-73


Registrant: Howard Johnson's Enterprises, Inc.
700 W. Virginia Street
Milwaukee, WI 53204-1548

Action: Notification of Additional Brand Name

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152-46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and CFR 152-46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.



Janet P. Baguhn
Compliance Manager





CRABGRASS

Plus with 0.0

20-2-8

A SELECTIVE HERBICIDE THAT PROVIDES CONTROL OF LISTED ANNUAL GRASSES AND BROADLEAF WEEDS IN ESTABLISHED LAWNS & ORNAMENTAL TURF. CONTAINS A COMPLETE LAWN FOOD.

ACTIVE INGREDIENT:

Dithiopyr: 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester 0.06%
 INERT INGREDIENTS: 99.94%
 TOTAL: 100.00%

EPA Reg. No. 32802-73
 EPA Est. No. 32802-WI-1 (A), 32802-MO-1 (B)
 First letter of bag code shows establishment where manufactured.

DIMENSION* is a trademark of Dow AgroSciences LLC

Read the entire "Directions for Use" and "Warranty Statement" before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

GUARANTEED ANALYSIS

Total Nitrogen (N) 20%
 0.78% Ammoniacal Nitrogen
 19.22% Urea Nitrogen *
 Available Phosphate (P₂O₅) 2%
 Soluble Potash (K₂O) 8%
 Total Sulfur (S) 1.73%
 1.73% Free Sulfur (S)

Derived From: Ammonium Phosphate, Muriate of Potash, Polymer Coated-Sulfur Coated Urea, Urea

* 4.00% Slowly Available Nitrogen from Polymer Coated-Sulfur Coated Urea

Item # 95882

Manufactured by:
 Howard Johnson's Enterprises, Inc.
 700 W. Virginia St.
 Milwaukee, WI 53204
 800-298-4656

Manufactured for:
 Garden Supply Distributors, Inc.

Net Wt. 40 lbs (18.20 kg) • Covers 10,000 sq. ft.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

GENERAL INFORMATION: This product controls crabgrass and other annual grasses and broadleaf weeds, and also contains fertilizer for proper turf growth in established lawns. It can be applied 2 to 4 weeks later than other crabgrass preventers, and provides up to 3 to 5 months control of crabgrass and other weed grasses and broadleaf weeds in home lawns. It can be applied according to these directions to the following turfgrasses:

Cool season grasses: creeping bentgrass, Kentucky bluegrass, fine fescue, tall fescue, perennial ryegrass;

Warm season grasses: bahiagrass, bermudagrass, buffalograss, carpet-grass, centipedegrass, kikuyugrass, St. Augustinegrass, zoysiagrass.

Apply only to lawns that are well established. Use of this product on lawns that are not well established, or weakened by weather, pests or mechanical stress, may increase the chance for turf injury. This product will not harm nearby established ornamentals when used according to label directions.

When to apply: This product can be applied 2 to 4 weeks later than other crabgrass preventers, because Dimension* herbicide in this product provides preventive (pre-emergence) control when spread on lawns prior to germination of the weed seeds. It also provides post-emergence control of crabgrass when applied 2 to 4 weeks after crabgrass germination. Post-emergence control is limited to crabgrass only. Application of this product later than 4 weeks after crabgrass germination will result in reduced control of crabgrass. For control of the weed grasses and broadleaf weeds listed on this label (other than crabgrass), it must be applied as a preventive treatment before the weed seeds germinate. ONLY crabgrass can be controlled with delayed application after germination.

APPLICATION DIRECTIONS: This product can be applied in the spring, before weeds become established in the lawn, or in the fall for pre-emergence crabgrass control through the early part of the next spring. Fall applications should be followed by a second (sequential) application the following spring to provide season-long control. Do not apply more than 9.6 pounds per 1000 square feet per year. Do not make more than one application per year.

LENGTH OF WEED CONTROL: The length of time of weed control is directly related to the amount applied to the lawn. Use the application rates in Table 1 to provide 2 to 3 months of weed control. Use the application rates in Table 2 for 3 to 5 months of weed control. The actual length of weed control will vary depending on weather, weed pressure, and your location in the north or south region.

USE PRECAUTIONS: DO NOT apply more than 9.6 pounds of this product per 1000 square feet per year. DO NOT apply to flowers, vegetables or fruit trees. DO NOT use clippings from treated turf for mulching around flowers, vegetables or fruit trees. DO NOT apply this product to pastures. Keep people and pets off treated areas until dust has settled. Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When seeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. For best results, use "slit seeders" designed to give good seed-to-soil contact.

TIPS FOR IMPROVED CONTROL: For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. For improved weed control, water or irrigate treated areas within a few days after application. Aeration, raking or dethatching of lawns should be done before application of this product. **WEEDS CONTROLLED:** This product will control (at post-emergence rates) or suppress (at pre-emergence rates) these weed grasses and broadleaf weeds when applied before weeds emerge in lawns: crabgrass, annual bluegrass (*Poa annua*), barnyardgrass, crowfootgrass, foxtail (yellow & green)*, goosegrass, kikuyugrass (seedling), common lespedeza,

buttercup oxalis, creeping oxalis, yellow oxalis (common), ryegrass (annual & perennial), prostrate spurge, spotted, spurge, henbit, parsley-piert.

TABLE 1: RATES OF CRABGRASS Pre-emergence

Region	Pre-emergence
North	3.6 lbs. per 1000 sq. ft.
South	7.2 lbs. per 1000 sq. ft.

TABLE 2: RATES OF CRABGRASS Pre-emergence

Region	Pre-emergence
North	4.8 lbs. per 1000 sq. ft.
South	9.6 lbs. per 1000 sq. ft.

*Southern states include AL, AR, NV, OK, SC, TN, TX, VA, and southern states included as North Region.

*Foxtail is also controlled at the pre-emergence rate. When soil testing or professional analysis indicates low nutrient levels, make additional applications. This product may stain driveways, walkways, etc. Apply thoroughly immediately after each use. NOTICE: Buyer and user assume responsibility for proper handling of this product not in accordance with label directions.

SPREADER SETTINGS: to apply

- Ames Drop or Rotary
- EarthWay Drop or Rotary
- Precision Red Devil Rotary
- Republic EZ Drop
- Republic EZ Rotary
- Cyclone SB3000 Rotary
- Scotts EasyGreen EG-3 (rotary)
- Scotts SpeedyGreen (rotary)

Settings shown are approximate need calibration, due to different equipment condition. Calibrate before use. Clean & lubricate spreader after use.

STORAGE

Do not contaminate water, food, or feed. **STORAGE:** Store in original container, away from reach of children and domestic animals. **PESTICIDE DISPOSAL:** Securely wrap in several layers of newspaper and discard in a designated area. **CONTAINER DISPOSAL:** Do not reuse. Do not discard in trash.

NOTIFICATION
JUN 26 2006

BGRASS CONTROL

Plus with 0.06% Dimension®

Federal law to use this product in a sale or other commercial use, or for other purposes.

controls crabgrass and other annual broadleaf weeds. Apply 2 to 4 weeks later than other crabgrass control products. It can be applied according to label directions.

controls crabgrass and other annual broadleaf weeds. Apply 2 to 4 weeks later than other crabgrass control products. It can be applied according to label directions.

Use of this product on lawns that are stressed by weather, pests or mechanical stress, may reduce effectiveness. Follow label directions.

Apply 2 to 4 weeks later than other crabgrass control products. This product provides pre-emergence control of crabgrass when applied to lawns prior to germination. Post-emergence control is provided later than 4 weeks after application of crabgrass. For control of weeds listed on this label (other than crabgrass), apply before the weed seeds germinate with delayed application after

can be applied in the spring, before crabgrass germinates, or in the fall for pre-emergence control of the next spring. Fall applications are most effective. Apply according to label directions.

Application rates in Table 1 to provide pre-emergence control of crabgrass. Application rates in Table 2 for 3 to 5 months of weed control will vary depending on region.

Apply 2 to 4 weeks later than other crabgrass control products. This product provides pre-emergence control of crabgrass when applied to lawns prior to germination. Post-emergence control is provided later than 4 weeks after application of crabgrass. For control of weeds listed on this label (other than crabgrass), apply before the weed seeds germinate with delayed application after

Apply 2 to 4 weeks later than other crabgrass control products. This product provides pre-emergence control of crabgrass when applied to lawns prior to germination. Post-emergence control is provided later than 4 weeks after application of crabgrass. For control of weeds listed on this label (other than crabgrass), apply before the weed seeds germinate with delayed application after

buttercup oxalis, creeping oxalis, yellow woodsorrel, pineappleweed, purslane (common), ryegrass (annual & perennial), smutgrass (seedling), corn speedwell, prostrate spurge, spotted, spurge, bittercress, chickweed, Carolina geranium, henbit, parsley-piert.

TABLE 1: RATES FOR 2 TO 3 MONTHS OF CRABGRASS CONTROL*

Region	Pre-emergence	Post-emergence
North	3.6 lbs. per 1000 sq. feet	7.2 lbs. per 1000 sq. feet
South	7.2 lbs. per 1000 sq. feet	9.6 lbs. per 1000 sq. feet

TABLE 2: RATES FOR 3 TO 5 MONTHS OF CRABGRASS CONTROL*

Region	Pre-emergence	Post-emergence
North	4.8 lbs. per 1000 sq. feet	9.6 lbs. per 1000 sq. feet
South	9.6 lbs. per 1000 sq. feet	9.6 lbs. per 1000 sq. feet

*Southern states include AL, AR, AZ, CA, FL, GA, HI, LA, MS, NC, NM, NV, OK, SC, TN, TX, VA, and southern areas of IL, IN and OH. All other states included as North Region.

*Foxtail is also controlled at the pre-emergence rates.

When soil testing or professional advice indicates the need for additional plant nutrients, make additional applications of fertilizer products over the growing season.

This product may stain driveways, walks, patios, pools, etc. Wash or sweep off thoroughly immediately after each use.

NOTICE: Buyer and user assume all risks and liability of use, storage and/or handling of this product not in accordance with the directions for use on this label.

SPREADER SETTINGS: to apply at the listed pounds per 1000 sq. ft.

	3.6 lbs	4.8 lbs	7.2 lbs	9.6 lbs
Ames Drop or Rotary	14	17	20	23.5
EarthWay Drop or Rotary	15	17	21	24.5
Precision Red Devil Rotary	5	7	11	12
Republic EZ Drop	6.5	7.5	11	12
Republic EZ Rotary	6.5	7.5	11	12
Cyclone SB3000 Rotary	6.75	7.25	7.75	8.5
Scotts EasyGreen EG-3 (rotary)	28	29	31	29 x 2
Scotts SpeedyGreen (rotary)	9	11	16	18.5

Settings shown are approximate for new equipment. Your spreader may need calibration, due to differences in terrain, walking speed and equipment condition. Calibrate your spreader to ensure proper coverage. Clean & lubricate spreader after use.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in a locked storage area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Fold empty bag and discard in trash.

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS:

CAUTION: Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift or runoff may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

First Aid

If in eyes • Hold eye open and rinse slowly and gently with water for 15-20 minutes.
• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
• Call a poison control center or doctor for treatment advice.

If inhaled • Move person to fresh air.
• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
• Call a poison control center or doctor for further treatment advice.

If on skin or clothing • Take off contaminated clothing.
• Rinse skin immediately with plenty of water for 15-20 minutes.
• Call a poison control center or physician immediately for treatment advice.

If swallowed • Call a poison control center or physician immediately for treatment advice.
• Have person sip a glass of water if able to swallow.
• Do not induce vomiting unless told to do so by the poison control center or physician.
• Do not give anything by mouth to an unconscious person.

Hot line number:

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.