



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (H7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg. Number:  
32802-74

Date of Issuance:  
MAR 30 1995

NOTICE OF PESTICIDE:  
 Registration  
 Reregistration

Term of Issuance:  
Conditional

Name of Pesticide Product:  
Trimec Weed & Feed  
for Northern  
Turfgrasses

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):  
Howard Johnson's Enterprises, Inc.  
P.O. Box 2990  
Milwaukee, WI 53201

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes:

a. Revise the EPA Registration Number to read, "EPA Reg. No. 32802-74".

b. Add the following groundwater statement, per the 2,4-D exposure reduction labeling instructions: "Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination."

Signature of Approving Official:

Date:  
MAR 30 1995

7505C  
H7505C

2011

- c. Under the section on "Where, When, and How to Use," there is a typographical error in the fourth line of the paragraph. Replace the word "whin" with "within".
3. Submit one copy of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

# TRIMEC® WEED & FEED



Manufactured by

## FOR NORTHERN TURFGRASSES

### A TRIPLE ACTION BROADLEAF HERBICIDE WITH LAWN FOOD

(16, 18 or 20) lbs. Treats  
5,000 Sq.Ft. of Turfgrass.

For Use in Control of Broadleaf Weeds in Home Lawns, Apartment and Condominium Turf, In Parks, Playgrounds and other Ornamental & Recreational Turfgrass.

**BALANCED LAWN FOOD GIVES TURF FAST GREENING WITH EXTENDED FEEDING CONTROLS:** Dandelions, Chickweed, Knotweed, Plantains, Henbit, Spurge, and Many Other Weeds as Listed.

<b>ACTIVE INGREDIENTS:</b>	
2,4-Dichlorophenoxyacetic acid - - - - -	0.5471%
2-(2-methyl-4-chlorophenoxy) propionic acid - - -	0.2448%
Dicamba (3,6 dichloro-o-anisic) acid - - - - -	0.0516%
<b>INERT INGREDIENTS:</b>	99.1565%
Total	100.0000%

\* Equivalent to 10,942 lbs. 2,4-Dichlorophenoxy acetic acid per Ton  
 \*\* Equivalent to 4,896 lbs. 2-(2-methyl-4-chlorophenoxy propionic acid) per Ton  
 \*\*\* Equivalent to 1,032 lbs. (3,6 dichloro-o-anisic) acid per Ton  
 (Isomers Specific by AOAC 14th Edition #6.321)

### KEEP OUT OF REACH OF CHILDREN CAUTION

#### STATEMENTS OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water. For eyes, flush with water for 15 minutes and get medical attention.

See Side/Back Panel For Additional Precautionary Statements

#### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long sleeved shirt, long pants, shoes plus socks. After using product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

#### ENVIRONMENTAL HAZARD

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target areas.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled. Not for use on turf grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in its original labeled container in a cool, dry, locked place out of reach of children and domestic animals. **DISPOSAL:** Do not reuse empty bag. Securely wrap original container in several layers of newspaper and discard in trash.

**IMPORTANT:** Do not apply this product to Dichondra, Carpetgrass, or improved varieties of St. Augustinegrass in Florida.

**NOTICE:** Buyer and user assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label (NEWTRIMEC PM51 12-19-34)

EPA REG. NO. 32802-

EPA EST. NO. 32802-WI-1

Net Weight (16;18; or 20) Pounds

**HOWARD JOHNSON'S ENTERPRISES, INC.** <sup>3</sup>

P.O. Box 2990 - Milwaukee, WI 53201 U.S.A.

### BROADLEAF WEEDS CONTROLLED

#### A PARTIAL LIST OF WEEDS CONTROLLED WITH THIS PRODUCT

Bedstraw	Croton	Kochia	Poison Ivy/Oak	Velvetleaf
Black Medic	Curly Dock	Lambsquarter	Povertyweed	Vetch
Blue Lettuce	Dandelion	Lespedeza	Purslane	Wild Carrot
Buckhorn	English Daisy	Mallow, Venice	Ragweed	Wild Garlic
Bull Thistle	Fleabane	Morning glory	Shepherdspurse	Wild Lettuce
Burdock	Florida Pusley	Mustard	Sheep Sorrel	Wild Onion
Buttercup	Ground Ivy	Oxalis	Smartweed	Yarrow
Chickweed	Hawkweed	Pennywort	Speedwell	and many other Broadleaf Weeds
Chicory	Healall	Peppergress	Spurge	
Clover	Henbit	Pigweed	Thistle	
Cocklebur	Knotweed	Plantains	Vervain	

### WHERE, WHEN, AND HOW MUCH TO USE

Do not use this product where desirable clovers are present or allow the material to "wind drift" onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Do not exceed specified dosage for any area; be particularly careful when dripline of tree and other ornamental species. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning to prevent over-application.

Do not use on newly seeded lawns until after the second cutting. After application delay seeding for at least three weeks and increase the rate of seed put down by 25%.

Apply when weeds are young and actively growing in the spring, summer, or fall. Best results will be obtained when temperatures range from 65 to 90° F., and weeds are in a vigorous state of growth from rain and/or watering. Results will be erratic if applied during drought periods. Lawns that are heavily over-run by hard-to-kill weeds may benefit from an additional treatment in 30 to 45 days. Do not water lawns for 2 days after application.

The maximum rate for turf is 10 pounds of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

Select an appropriate Spreader and distribute 18 lbs. per 5,000 Sq. Ft. (3.6 lbs. / 1,000 Sq. Ft.) of turf for control of the many weeds as listed and to supply the lawn with a complete NPK fertilization at the same time.

#### MAKE SURE YOUR SPREADER IS OPERATING CORRECTLY

For help refer to your Spreader Guide Book. Poor weed control will result if your spreader applies less than recommended. Make sure to apply the total pounds per sq. ft. as shown in the following chart.

#### SPREADER SETTINGS

SPREADER	Amount to Apply per 5,000 Sq. Ft. and Spreader Setting
Scott	16 lbs. #6 1/2      18 lbs. #7      20 lbs. #7 1/2
Cyclone	16 lbs. #3 1/2      18 lbs. #4      20 lbs. #4 1/2

When using a 36 lb. bag, the spreader settings will be the same as for the 18 lb. bag but it will treat twice the Sq. Ft. These settings are based on a "One Time Over" application with a New spreader.

#### GUARANTEED FERTILIZER ANALYSIS

Nitrogen (N)	variable %	variable %
Sulfur Coated Urea	variable %	
Ammoniated Nitrogen	variable %	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	variable %	
Soluble Potash (K <sub>2</sub> O)	variable %	
Derived from Ammonium Sulfate, Sulfur Coated Urea, Ammoniated Phosphate, Muriate of Potash (as variable fertilizer inerts)		

424

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

MAR 30 1995

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

32802-74

f