

WEED & FEED FOR NORTHERN TURFGRASSES

***CONTAINS TRIMEC® HERBICIDE**
TRIPLE ACTION BROADLEAF HERBICIDE WITH LAWN FOOD
CONTROLS BROADLEAF WEEDS IN HOME LAWNS AND OTHER ORNAMENTAL AND RECREATIONAL TURFGRASS LAWN FOOD GIVES TURF FAST GREENING WITH EXTENDED FEEDING

CONTROLS DANDELIONS, CHICKWEED, KNOTWEED, PLANTAINS, HENBIT, SPURGE AND MANY OTHER WEEDS AS LISTED

(16, 18 OR 20) POUNDS TREATS 5000 SQUARE FEET

for commercial seed production, or for research purposes.

BROADLEAF WEEDS CONTROLLED: A partial list of weeds controlled with this product:

Bedstraw	English Daisy	Pennywort	Velvetleaf
Black Medic	Fleabane	Peppergrass	Vetch
Blue Lettuce	Florida Pusley	Pigweed	Wild Carrot
Buckhorn	Ground Ivy	Plantains	Wild Garlic
Bull Thistle	Hawkweed	Poison Ivy/Oak	Wild Lettuce
Burdock	Healall	Povertyweed	Wild Onion
Buttercup	Henbit	Purslane	Yarrow
Chickory	Knotweed	Ragweed	
Chickweed	Kochia	Shepherd's purse	
Clover	Lambsquarter	Sheep Sorrel	and
Cocklebur	Mallow, Venice	Speedwell	many other
Croton	Morningglory	Spurge	broadleaf
Curly Dock	Mustard	Thistle	weeds
Dandelion	Oxalis	Vervain	

HOW, WHEN AND HOW MUCH TO USE: Apply when weeds are young and actively growing in the spring, summer or fall. Best results will be obtained when temperatures range from 65° to 90°F. and weeds are in a vigorous state of growth from rain and/or watering. Results will be erratic if applied during drought periods. Lawns that are heavily over-run by hard-to-kill weeds may benefit from an additional treatment in 30 to 45 days. Do not water lawns for 2 days after application.

The maximum rate for turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

(Optional instructions:

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water before application to hold the material and prevent dust drift onto non-target plants.
4. Do not wash material from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.)

IMPORTANT: Do not apply this product to Dichondra, Carpetgrass, or improved varieties of St. Augustinegrass in Florida. Do not use this product where desirable clovers are present or allow the material to "wind drift" onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Do not exceed specified dosages for any area; be particularly careful within the drip line of trees and other ornamental species. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning to prevent over-application.

Do not use on newly seeded lawns until after the second cutting. After application, delay seeding for at least three weeks and increase the amount of seed put down by 25%.

Select an appropriate spreader and apply 3.2 pounds per 1000 square feet (for 16 pound bag; or 3.6 pounds per 1000 square feet for 18 pound bag, or 4 pounds per 1000 square feet for 20 pound bag) of turf for control of the many weeds as listed and to supply the lawn with a complete NPK fertilization at the same time.

Optional advertising claims:

Prepares your lawn for winter dormancy.

The key to a beautiful lawn next spring.

(Label for EPA Reg. No 32802-74, PAGE 1 OF 3)

ACTIVE INGREDIENTS:

2,4-Dichlorophenoxyacetic acid	0.55%
(+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid	0.12%
Dicamba (3,6-dichloro-o-anisic acid)	0.05%
OTHER INGREDIENTS:	99.28%
TOTAL:	100.00%

This product contains:

- 11. 2,4-Dichlorophenoxyacetic acid equivalent per ton.
 - 2.4 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton.
 - 1.0 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.
- Isomer Specific by AOAC Methods
Contains the single isomer form of Mecoprop-p.

EPA Reg. No. 32802-74

EPA Est. No. 32802-WI-1 (A), 32802-MO-1 (B)

First letter of bag code shows establishment where manufactured.
See tag attached to bag for fertilizer guaranteed analysis.

Manufactured by:

HOWARD JOHNSON'S ENTERPRISES, INC.
700 W. Virginia Street, Milwaukee, WI 53204-1548

TRIMEC® is a registered trademark of PBI/GORDON CORP.

Net Weight (16, 18 OR 20) lbs.

Read the entire "Directions for Use" and "Warranty Statement" before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

STOP - READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

KEEP OUT OF REACH OF CHILDREN PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, shoes plus socks. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

ENVIRONMENTAL HAZARDS: Drift or run-off may adversely affect non-target plants. Do not apply when weather conditions favor drift from target area. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

Not for use on turf grown for sale or other commercial use as sod, or

Promotes deeper green lawns and early rapid growth.

For long-lasting results.

Controls (Kills) dandelions and other tough broadleaf weeds.

Fertilizer may stain driveways, walks, patios, pools, etc. Sweep off thoroughly immediately after each use.

Drop Spreader: First apply two strips at ends, then apply product along the length of the lawn, overlapping very slightly, and shutting off spreader when turning.

Rotary Spreader: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

OPTIONAL Fertilizer labeling language:

For state registrations in Kentucky, add the following paragraph:

A lawn will typically use 1 to 4 pounds of nitrogen per 1,000 sq. ft. per growing season. When used as directed, this product will supplement your regular fertilizer program by providing (XX number of pounds) of nitrogen per square feet. Please consult your local extension office for questions regarding a regular fertilizer program suitable for your geographic region.

All fertilizer products distributed in Washington at a minimum, must have the following labeling statement:

"This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product."

After July 1, 1999, the label must also state: "Information received by the Washington State Department of Agriculture regarding the components in this product is available on the internet at <http://www.wa.gov/agri/>."

OR "Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.xxx.com/>."

For states other than WA, call 1-800-298-4656.

First Aid	
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.</p>	

SPREADER SETTINGS:

The settings shown here are approximate for new equipment. Your spreader may need calibration, due to differences in terrain, walking speed, and equipment condition. Calibrate spreader to ensure correct application rate. Clean and lubricate your spreader after each use.

Spreader setting to apply at listed rate (pounds per 1000 sq. ft.)	X lbs.
Spreader A	
Spreader B	
Spreader C	
Spreader D	
Spreader E	
Spreader F	
Spreader G	
Spreader H	

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original labeled container in a cool, dry, locked place out of reach of children and domestic animals.

Disposal: If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or toll-free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain or toilet.

GUARANTEE: This product is unconditionally guaranteed.

(Alternate wording for warranty statement)

GREEN GUARANTEED: Your satisfaction is guaranteed when this product is used in accordance with instructions. If product performance is less than satisfactory, seller will refund the full purchase price with evidence of purchase.

(Alternate wording for warranty statement)

NOTICE: Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

(Alternate wording for warranty statement)

WARRANTY AND DISCLAIMER: Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

(Alternate wording for warranty statement)

WARRANTY AND DISCLAIMER: Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. No other warranty, expressed or implied, is made concerning the merchantability or fitness for a particular purpose. Manufacturer/seller's liability shall be limited to the purchase price of the product. In no case shall manufacturer/seller be liable for consequential or incidental damages or loss. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

(Alternate wording for warranty statement)

WARRANTY AND DISCLAIMER: Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. No other warranty, expressed or implied, is made concerning the merchantability or fitness for a particular purpose. Manufacturer/seller's liability shall be limited to the purchase price of the product. In no case shall manufacturer/seller be liable for consequential or incidental damages or loss. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label. Products not accepted under these terms should be returned within 20 days in the original, unopened container for return of purchase price.

- WEEDS (KILLED OR CONTROLLED) BY WEED & FEED FOR NORTHERN TURFGRASSES-

Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeye	Pearwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	False dandelion	Plains coreopsis	Tumble mustard
Blessed thistle	Falseflax	Plantain	Tumble pigweed
Bloodflower milkweed	False sunflower	Poison ivy	Velvetleaf
Blue lettuce	Fiddleneck	Poison oak	Venice mallow
Blue vervain	Florida pusley	Pokeweed	Virginia buttonweed
Bracted plantain	Frenchweed	Poorjoe	Virginia creeper
Bristly oxtongue	Galinsoga	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly sida	Western clematis
Broomweed	Gumweed	Prostrate knotweed	Western salsify
Buckhorn	Hairy fleabane	Prostrate pigweed	White mustard
Buckhorn plantain	Hawkweed	Prostrate spurge	Wild aster
Bulbous buttercup	Healall	Prostrate vervain	Wild buckwheat
Bull thistle	Heartleaf drymary	Puncturevine	Wild carrot
Bullnettle	Heath aster	Purshane, common	Wild four-o'clock
Burclover	Hedge bindweed	Ragweed	Wild garlic
Burdock	Hedge mustard	Red sorrel	Wild geranium
Burning nettle	Henbit	Redroot pigweed	Wild lettuce
Burweed	Hoary cress	Redstem filaree	Wild marigold
Buttercup	Hoary plantain	Rough cinquefoil	Wild mustard
Buttonweed	Hoary vervain	Rough fleabane	Wild onion
Canada thistle	Horsenettle	Russian pigweed	Wild parsnip
Carolina geranium	Jimsonweed	Russian thistle	Wild radish
Carpetweed	Knawel	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Knotweed	Scotch thistle	Wild strawberry
Catnip	Kochia	Sheep sorrel	Wild sweet potato
Catsear	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, common	Lespedeza	Slender plantain	Woodsorrel
Chickweed, mouseear	Mallow	Smallflower galinsoga	Woolly croton
Chicory	Matchweed	Smooth dock	Woolly morningglory
Cinquefoil	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, crimson	Milk vetch	Sorrel	Wormseed
Clover, hop	Morningglory	Sowthistle	Yarrow
Clover, red	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, strawberry	Mugwort	Speedwell	Yellowflower pepperweed
Clover, sweet	Musk thistle	Spiny amaranth	and other broadleaf weeds

Optional advertising claims:

Controls (Kills) (desired weed species from above complete list may be inserted here, such as dandelion, chickweed, knotweed, plantain, henbit, spurge, etc.) and many other broadleaf weeds; including those listed on the back (side) panel.

Contains (controlled- or slow-release) nitrogen.
Premium Quality
This product contains a highly effective weed killer and a (winter) lawn conditioner.

Phosphate-free

Controls (Number) weeds!

Place your spreader on the driveway or other paved surface when filling. This guards against spilling fertilizer on the grass. For even coverage, begin walking before you turn the spreader to the open position, to avoid dropping fertilizer in one spot when spreader is opened. Be sure the spreader is closed when filling it, when turning to back-track, or when stopping on the lawn.

ACCEPTED
JAN 30 2003
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 32802-74

