



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (H7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg.  
Number:  
32802-79 ✓

Date of Issuance:  
SEP 23 2003

Term of Issuance:  
Conditional

Name of Pesticide Product:  
RONSTAR® PLUS  
FERTILIZER 1.00%

NOTICE OF PESTICIDE:

Registration  
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Howard Johnson's Enterprises, Inc  
700 West Virginia Street, Suite 222  
Milwaukee, WI 53204

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Add the phrase "EPA Registration No. 32802-79" to the label before you release the product for shipment.

COMMENTS CONTINUED ON PAGE 2 OF THIS NOTICE OF REGISTRATION

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product under the enclosed stamped copy of the label constitutes acceptance of these conditions.

Enclosure

Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505C)

Signature of Approving Official:

*Joanne I. Miller*

Date:

SEP 23 2003

- 2. Submit the following data required for the registration of this pesticide product within 1 year from the date of this Notice of Registration:

<u>EPA Guideline Data Number</u>	<u>Guideline Descriptor</u>
830.6317	Storage Stability Study
830.6320	Corrosion Characteristics Study

- 3. Resubmit the Confidential Statement of Formula (CSF) on EPA Form 8570-4 to reflect the present registrant of Ronstar Intermediate 50% the source of oxadiazon in this pesticide product. Please use your company's correct name on all forms and correspondence.
  - 4. Give name of the present owner of the trade mark name "Ronstar" and move the asterisk to proceed CAS, i.e.: \*CAS No. 19666-30-9; RONSTAR® is a Trade Marked name of Bayer CropScience LP.
  - 5. Under "User Safety Recommendations" section revise the phrase "Users should" to read: "Applicators should".
  - 6. Delete the claim for "Commercial lawns" on page 1. Correct the spelling of "barrier" in the "AUR" section and "chemical" in the PPE statements; and remove dates on this label.
  - 7. Under the "Precautions in Using on Turf Grass" section the sentence that end in the words: "...and normally outgrown within 2 to 3 weeks." must be revised by deleting that phrase. Also, delete the phrase "because of the varying cultural practices used." in the last sentence of this section.
  - 8. Delete (Cotoneaster spp) from the "Varieties" section of "Sensitive Plants"; for consistence give the Latin names of the Plant Species listed in this section.
  - 9. Submit one (1) copy of the final printed labeling before you release this product for shipment.
  - 10. Submit and/or cite all data required for the registration of this product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of this product under FIFRA, section 4. A Stamped copy of the label is enclosed for your records.
-

# RONSTAR® PLUS Fertilizer 1.00%

FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY, TURF MANAGEMENT AND LANDSCAPE PERSONNEL. NOT FOR USE ON HOME LAWNS. FOR USE ON GOLF COURSES, NURSERIES, PARKS, SOD FARMS, ATHLETIC FIELDS, AND OTHER RECREATIONAL AND ORNAMENTAL TURFGRASS.

A Turf Fertilizer plus a Pre-emergent Weed Control Product for use on varieties of Perennial Bluegrass, Bentgrass, Bermuda grass, Buffalograss, Perennial Ryegrass, St. Augustine grass, Seashore Paspalum, Tall Fescue and Zoysiagrass. Also For Use Around Ornamental Shrubs, Vines, Trees, and Ground Covers, as listed.

ACTIVE INGREDIENT: Oxadiazon\* - - - - 1.00%  
 OTHER INGREDIENTS: - - - - - 99.00%  
 \* CAS [19666-30-9] Contains RONSTAR® a Reg. TM of Mobby Corp. 100.00%

**KEEP OUT OF REACH OF CHILDREN  
 WARNING - AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### FIRST AID

**IF SWALLOWED** - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **IF IN EYES** - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. **IF ON SKIN** - Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. **IF INHALED** - Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. **READ ENTIRE LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.**

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING:** Causes substantial but temporary eye injury. Wear protective eyewear (goggles, face shield or safety glasses). Causes skin irritation. Do not get in eyes, on skin or clothing. Harmful if swallowed, absorbed through the skin, or inhaled. Avoid breathing dust, or contact with skin, eyes or clothing. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling. Do not allow domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear coveralls over short-sleeved shirt and short pants; chemical resistant gloves, chemical resistant footwear plus socks; protective eyewear, headgear for overhead exposure; chemical resistant apron when cleaning equipment. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigations system. Apply only as specified on this label.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests and nurseries, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.


Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls; chemical resistant (Barrier Laminate) gloves, chemical resistant footwear plus socks, protective eyewear, and chemical resistant headgear for overhead exposure.

**Howard Johnson's Enterprises, Inc.**

700 West Virginia St. - Suite 222 - Milwaukee, WI 53204

ACCEPTED WITH COMMENTS  
 IN SEP 23 2003  
 SEP 23 2003

Under the National Insecticide and Fungicide Registration Act, this product is registered under EPA Reg. No. 32802-79



Label copy unit 1 of 3  
**NET WEIGHT  
 50 POUNDS**

Page 7 of 17

Controls Crabgrass, Goose grass, Poa Annua, Annual Sedge and other weeds as listed, on Golf Courses, Commercial Lawns, Parks, and Recreational and Ornamental turf.

### EPA Reg. No. -32802-TO EPA EST. NO. 32802-MO-1 WHERE, HOW MUCH AND WHEN TO USE ON TURFGRASS

This product may be used at full rate or as a "split application in established perennial bluegrass, bentgrass (see restrictions), Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum, Tall Fescue, and Zoysiagrass turf, at apartment/condominium complexes, corporate and public parks, golf courses, sod farms, athletic fields, playgrounds, cemeteries and other ornamental and recreational turfgrass area. Not for use on home lawns. The turf should be free of leaves and thatch and in condition to allow granules to reach the soil surface. Know the square footage of the area to be treated and uniformly apply using a granule spreader. Do not apply to turf when it is wet. When using at full rate do not overlap spreader applications, especially on St. Augustinegrass because discoloration "streaking" damage may temporarily occur from the "double rate". See section on this label entitled: **PRECAUTIONS IN USING ON TURFGRASS** for the maximum rate for St. Augustinegrass and Bentgrass of 3 lbs. a.i. per acre which is equiv. to 6.9 lbs. of this product per 1,000 sq. ft.). After application if rain is not expected within 24 hours sprinkler irrigate the treated turf with up to a 1/4 inch of water to release the herbicide for maximum weed control and quicker response from the fertilizer nutrients. This product is a pre-emergence herbicide and must be applied prior to weed seed germination. It will not control growing weeds. \*\*For a SPLIT APPLICATION, see the "FOR SPRING APPLICATION ON GOLF COURSES" section, for instructions on how to apply the 3 lb. or 4 lb. a.i. per acre rate as a split application.

### Pre-emergence Weed Control

Weeds Controlled	lbs. a.i. / Acre	lbs. of Product per 1,000 sq. ft.	Directions - Refer to Map of U.S.
Crabgrass ( <i>Digitaria</i> , spp.)	2 to 4 lbs. a.i./A	4.6 to 9.2 lbs.	Apply in late winter or early spring prior to weed seed germination. See map. In areas 4 & 5 applications made after April or May, may not be fully effective.
Goosegrass ( <i>Elysiine indica</i> )	2 to 4 lbs. a.i./A	4.6 to 9.2 lbs.	
Field Sandbur ( <i>Cynchus incertus</i> )	2 to 4 lbs. a.i./A	4.6 to 9.2 lbs.	
Annual Sedge ( <i>Cyperus compressus</i> )	2 to 4 lbs. a.i./A	4.6 to 9.2 lbs.	
Annual Bluegrass ( <i>Poa annua</i> spp.) (See Note Below)	2 to 4 lbs. a.i./A	4.6 to 9.2 lbs.	Apply in late summer to early fall prior to weed seed germination.
Virginia Buttonweed ( <i>Diodia</i> )	4 lbs. a.i./A	9.2 lbs. per 1,000 sq. ft.	Apply in early spring prior to weed seed germination.
Pigweed ( <i>Amaranthus</i> spp.)	400 lbs. product per Acre.		
Florida Pusley ( <i>Richardia scabra</i> )			
Oxalis ( <i>Oxalis</i> spp.)			
Stinging Nettle ( <i>Urtica dioica</i> )			
Carpetgrass ( <i>Axonopus officinis</i> )			

### Fall Application for Spring Weed Control in Cool Season Turf

Crabgrass, Goosegrass and other annual weeds as listed.	3 to 4 lbs. a.i./A	6.9 to 9.2 lbs. per 1,000 sq. ft.	For use in Bluegrass, Bentgrass, perennial Ryegrass and Tall Fescue. Make application in the fall after turf has become dormant. For late fall applications, seeding into treated areas should be delayed until 4 months after greenup.
---	--------------------	-----------------------------------	---

### Annual Bluegrass Control in Bermudagrass Fairways Overseeded with Perennial Ryegrass Only


Annual Bluegrass ( <i>Poa annua</i> spp.)	2 lbs. a.i./A	4.6 lbs. / 1,000 sq. ft.	Apply in late summer to early fall prior to <i>Poa annua</i> spp., seed germination. Any <i>Poa annua</i> spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where <i>Poa trivialis</i> is present in the overseed mixture. Apply the full 2 lb. rate in one application.
---	---------------	--------------------------	--

**NOTE:** In many areas annual bluegrass exists as at least two biotypes or subspecies of *Poa Annua*. The true annual biotype, "*Poa Annua* spp. annua" which germinates from seed every year is controlled, while the biotype, "*Poa Annua* spp. raptans" which behaves like a perennial and survives through the summer, as an established plant, cannot be controlled. Consult the extension service or University weed specialist in your area for more information concerning your locale.

### Annual Weed Control during turf establishment from "Sprigs" of Bermudagrass, Zoysiagrass, and Seashore Paspalum.

Crabgrass and Goosegrass, Annual Sedge and other annual weeds.	2 to 3 lbs. a.i./A 200 to 300 lbs. product per acre	4.6 to 6.9 lbs. product	Directions - Refer to Map of U.S. on this label. For Bermudagrass make application immediately prior to, or, immediately after sprigging. For Seashore Paspalum and Zoysia make application 10 to 14 days after sprigging. Applications made close to the time of sprigging may cause objectionable injury. Disturbance of the soil surface after application may disrupt the herbicide barrier and reduce effectiveness. For best results sprinkler irrigate the "sprigged" area with up to 1/4 inch water after application. 02-11-03
--	--	-------------------------	---

**Crabgrass Seed Germination Dates**



1. Mid-north & northwestern After May 20-30
2. Upper/mid-states & northeast After May 10
3. Lower mid-states After April 20
4. Upper southern states After Mar 20
5. Warm season states Jan 1 to Mar 20

**NON-AGRICULTURAL USES REQUIREMENTS**

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, and nurseries. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Wear protective clothing including long-sleeved shirt and impervious gloves when handling. Launder exposed clothing before reuse. Do not allow entry to areas treated until dusts have settled or if watering in is required, until treated area and grass is dry.

**FOR SPRING APPLICATION ON GOLF COURSES**

The 3 lb. and 4 lb. active ingredient, per acre rates may be applied as a split application. Weed control and better turf management may be programmed by splitting the per acre rate into "two" separate applications. For example to accomplish this in region #5 (See map of U.S.). Apply 1.5 lbs. a.i. (for a required 3 lbs. a.i./acre rate) or 2 lbs.a.i. (for a required 4 lb. a.i./acre rate) as the first application 1 to 2 weeks before estimated weed seed germination or in late December or early January. Follow with the second "1/2 rate" in 60 days or the first of March. The timing for these applications should be closely programmed to the geographic region and adjusted to an "early" or "late" spring. Do not split the 2 lb. a.i. / acre rate. The 3 lb.a.i. / acre rate should not be split if applied to new turf areas first coming under turf management where weed pressure is expected to be heavy.

**PRECAUTIONS IN USING ON TURFGRASS**

For St. Augustinegrass and Bentgrass turf, DO NOT exceed the rate of 3 lbs. a. i. per acre which is equal to 6.9 lbs. of this product per 1,000 sq.ft. DO NOT apply to Bentgrass turf mowed less than 3/8 inch. This product may cause discoloration of Bentgrass, Bermudagrass and St. Augustinegrass but this discoloration is temporary and normally outgrown within 2 to 3 weeks. Avoid contact with flowers and shrubs except as recommended elsewhere for ornamental plants on this label. DO NOT apply to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment except as noted above. DO NOT apply to Dichondra or Centipedegrass. DO NOT use on golf course putting greens or tees because of the varying cultural practices used.

**GUARANTEED FERTILIZER ANALYSIS**

Total Nitrogen (N).....	variable	%
Ammoniacal Nitrogen	variable	%
Water Soluble Nitrogen	variable	%
and/or Urea Nitrogen	variable	%
Water Insoluble Nitrogen	variable	%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	variable	%
Soluble Potash (K <sub>2</sub> O).....	variable	%

Derived from Ammonium Sulfate, Urea, Ureaform, Muriate of Potash; Manganese as Mn %; Iron as Fe %; Chlorine < %

**WEEDS CONTROLLED**

- Annual Bluegrass
- Annual Sedge
- \*Barnyardgrass
- Bittercress
- Carpetweed
- Carpetgrass
- Common Groundsel
- Common Purslane
- Crabgrass
- \*Evening Primrose
- Fireweed
- Field Sandbur
- Foxtail; Green, Bristly
- Fall Panicum
- Florida Pusley
- \*Galinsoga
- Golden Ragwort
- \*Goosegrass
- \*\*Jungle Ricegrass
- Lambsquarter
- \*\*Liverwort
- Oxalis
- \*\*Panama Paspalum
- Pennsylvania Smartweed
- Redroot Pigweed
- Sensitive plant
- Shepardspurse
- Sow thistle
- Speedwell- Spotted Catsear
- Spurge; Garden, Petty,
- \*\*Prostrate
- Stinging Nettle
- \*Swinecress
- Virginia Buttonweed
- Yellow Woodsorrel.

\*For the weeds with 1 asterisk above, use 3 to 4 lbs. a.i. per acre (equal to 300 lbs. to 400 lbs. of this product per acre or 6.9 lbs. to 9.2 lbs. per 1,000 sq.ft.).

\*\*For the difficult-to-control weeds with 2 asterisks above, use 4 lbs. a.i. / acre (equal to 400 lbs. of this product /acre or 9.2 lbs./ 1,000 sq.ft.)

**NOTE:** For newly seeded lawns do not apply this product until after the second cutting. For best turf management, place new grass seed in the fall. Spring pre-emergence weed control can then be applied.

**SPREADER SETTINGS**

The following spreader settings are for new equipment. Used, or heavily worn equipment may require adjustments. For proper setting calibration refer to the spreader manufacturer's Maintenance Manual. Do not apply this product until equipment is properly calibrated to accurately apply the desired lbs. a. i. per acre, as recommended on this label.

Spreader Settings	for 1.5 lb. a.i./Acre	for 2 lb. a.i./Acre	for 3 lb. a.i./Acre	for 4 lb. a.i./Acre
Cyclone B3-Pro	#XX	#XX	#XX	#XX
Scotts Pro R-8A	#X	#X	#X	#X

Plan on an approximate 2 mph spreader speed to obtain best coverage. Know the exact square footage of the area before application. Incorrect application will result in poor weed control.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Store product in its original bag, in a cool, dry, locked place out of reach of children. In case of spill, sweep up, and use according to directions or dispose of as follows. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. Stay out of smoke.

**READ ENTIRE LABEL BEFORE USING**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties, and Limitation of Liability before using this product. If the terms are not acceptable, return the unopened container at once.

By using this product, user and buyer accept the following: Conditions, Disclaimer of Warranties, and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However it is impossible to eliminate all risks associated with the use of this product. Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Howard Johnsons. All such risks shall be assumed by the user or buyer.

**Disclaimer of Warranties:** There are no warranties express or implied, of merchantability or of fitness for a particular purpose or otherwise, which extend beyond the statements made on this label. No agent of Howard Johnsons is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Howard Johnsons disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

**LIMITATION OF WARRANTIES:** The exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract warranty, tort, negligence, strict liability or otherwise shall not exceed the purchase price, or at Howard Johnsons option, the replacement of the product.

FOR THE FOLLOWING EMERGENCY, PHONE 24 HOURS A DAY  
 TRANSPORTATION: CHEMTREC - 1-800-424-9300

It is important to recognize that this product can be used successfully on the many ground cover plants as here listed, because certain herbicides are not recommended.

For a Wide Variety of listed Woody \*Ornamental Shrubs, Vines, Trees, Landscape Plantings and for Container and Field Grown Nursery Plants.

\* For use only on these plants that will not bear edible fruit within one year after application.

**ESTABLISHED ORNAMENTALS and GROUND COVERS**

COMMON NAME	SCIENTIFIC NAME
Abutilon	Abutilon pictum
Acacia	spp.
Agapanthus	Agapanthus africanus
Allamanda, Common	Allamanda cathartica
Andromeda	Pieris spp.
Aralia	Aralia spinosa
Arborvita	Thuja spp.
Ash, Green	Fraxinus pennsylvanica
Azalea	Rhododendron
Baby's Breath	Gypsophila paniculata
Bald Cypress	Taxodium distichum
Bamboo, Featherly	Bambusa vulgaris
Banyan, Weeping	Ficus benghalensis
Barberry	Berberis spp.
Beefsteak Plant	Acalypha wilkesiana
Beardy	Arctostaphylos uva-ursi
Beech, European	Fagus sylvatica
Birch, Paper	Betula papyrifera
Blanket Flower	Gaillardia aristata
Blazing Star	Liatris spicata
Boxwood, Japanese	Buxus microphylla
Boxwood, Common	Buxus sempervirens
Bougainvillea	Bougainvillea spectabilis
Brassala, Dwarf	Brassala actinophylla
Buddleia	Buddleia davidii
Bugleflower	Ajuga reptans
Callistemon	Callistemon lanceolata
Camellia, Common, Jap	Camellia japonica
Camellia, Sasanqua	Camellia sasanqua
Candytuft	Iberis sempervirens
Carissa	Carissa macrocarpa
Carnation	Dianthus caryophyllus
Ceanothus	Ceanothus americanus
Cedar, Red	Juniperus virginiana
Cedar, Atlas	Cedrus atlantica
Cistus	Cistus purpureus
Citrus, Ornamental*	
Grapefruit	Citrus paradisi
Lemon	Citrus limon
Lime	Citrus aurantiifolia
Mandarin	Citrus reticulata
Tangelo	Citrus X Tangelo
Tangerine	Citrus reticulata var. tangerine
Temple	Citrus X nobilis
Cuban Shaddock	Citrus maxima
Valencia	Citrus sinensis
Washington Navel	Citrus sinensis
Cleyera	Cleyera japonica
Coreopsis	Coreopsis lanceolata
Coremandel	Asystasia gangetica
Cotoneaster	Cotoneaster, spp.
Crabapple	Malus, spp.
Crapemyrtle	Lagerstroemia indica
Chrysanthemum	Chrysanthemum, spp.
Cypress	Cupressus, spp.
Cypress, Leyland	Cupressocyparis leylandii
Dahlia	Dahlia, spp.
Daphne, Fragrant	Daphne odora
Deutzia, Slender	Deutzia gracilis
Dieffenbachia	Dieffenbachia, spp.
Dogwood, Common	Cornus florida
Dogwood, Redosler	Cornus stolonifera
Dogwood, Kousa	Cornus kousa
Douglas fir	Pseudotsuga menziesii
Elm	Ulmus, spp.
Escallonia	Escallonia, spp.
Eucalyptus	Eucalyptus, spp.
Euroyops, Gray Leaved	Euroyops pectinatus
Euroyopus, Spreading	Euroyopus klautshavicus
Euroyopus, Wintercreeper	Euroyopus fortunei
Falscycpress, Hink	Chaemaecyparis obtusa
Falscycpress, Japanese	Chaemaecyparis pisifera
Flowering Quince	Chaenomeles speciosa
Forsythia	Forsythia intermedia
Gardenia	Gardenia jasminoides
Gazania Trailing	Gazania uniflora
Ginko	Ginkgo biloba
Gladiolus	Gladiolus, spp.
Golden Raintree	Koeleruteria paniculata
Heath, Spring	Erica cornea
Heather, Scotch	Calluna vulgaris
Heliotropium	Heliotropium arborescens
Hemlock, Canada	Tsuga canadensis
Hibiscus	Hibiscus, spp.
Honeysuckle	Lonicera, spp.
Holly	Ilex, spp.
Ice Plant	Carpobrotus edule
Ilima	Sida fallax
Ivy, Algerian	Hedera canariensis
Ivy, English	Hedera helix
Jasmine, Orange	Murraya paniculata
Jasmine, Primrose	Jasminum mesnyi
Jasmine, Star	Trachelospermum jasminoides
Jasmine, Showy	Jasminum floridum
Juniper	Juniperis, spp.
Justica	Justica brandegeana
Kwanzan Cherry	Prunus serrulata
Lamb's Ear	Stachys byzantina
Lantana, Trailing	Lantana montivdensis
Lapalapa	Cheirodendron, spp.
Larch, European	Larix decidua
Laurel, English	Prunus laurocerasus
Leopards - Bane	Doronicum cordatum
Leucothoe, Coast	Leucothoe axillaris
Leucothoe, Drooping	Leucothoe foranensis (caesbaei)

**ESTABLISHED ORNAMENTALS and GROUND COVERS Continued**

COMMON NAME	SCIENTIFIC NAME
Lily of the Nile	Agapanthus umbellatus
Lilac	Syringa, spp.
Littleleaf Linden	Tilia cordata
Loquat*	Eriobotrya japonica
Magnolia	Magnolia, spp.
MaHonlia, Grape Holly	MaHonlia aquifolium
Maple	Acer, spp.
Metallic Plant	Hemigraphis colorata
Mexican Orange	Cholisyia temata
Mock Orange	Philadelphus virginalis
Money Tree	Pleomaie marginata
Murraea	Murraea exotica
Murraea	Murraea paniculata
Myoporum	Myoporum laetum
Myrtle	Myrsine communis
Nandina	Nandina domestica
Oak	Quercus, spp.
Ochana	Ochana serrulata
Ohal	Samanea saman
Oleander	erium oleander
Orthosiphon	Orthosiphon stamineus
Osmanthus	Osmanthus heterophyllus
Pachysandra	Pachysandra terminalis
Palm, Bamboo	Rhapi excelsa
Palm, Fishtail	Caryota, spp.
Palm, Licuala	Licuala Thunb. spp.
Palm, Macarthur	Ptychosperma macarthur
Palm, Mabnee	Euterpe, spp.
Palm, Manila	Veitchia merillii
Palm, Parlor	Collinia elegans
Palm, Rattian	Daemonorops, spp.
Palm, Scalling Wax	Cyrtostachyus lakka
Palm, Sugar	Arenga saccharifera
Paperbark Tree	Melaleuca leucadendra
Peony	Peonia, spp.
Periwinkle, Common	Vinca minor
Periwinkle, Large	Vinca major
Photinia, Frasier's	Photinis fraseri
Poinciana	Dolox regae
Pine	Pinus spp.
Pittosporum, Japanese	Pittosporum tobira
Plum, Natal	Carissa macrocarpa
Podocarpus, Yew	Podocarpus macrophyllus
Pomegranite	Punica granatum
Poplar	Populus spp.
Potentilla	Potentilla fruticosa
Potentilla	Potentilla fragiformis
Privet	Ligustrum, spp.
Protea species	Banksia menziesii
	Leucodendron discolor
	Leucospermum cordifolium
	Protea exinia
	Rhododendron, spp.
	Rhododendron, spp.
	Elaeagnus angustifolia
	Picea, spp.
	Osteospermum fruticosum
	Weddella trilobata
	Wisteria sinensis
	Taxus, spp.
Pyracantha	Pyracantha coccinea
Raphiolepis	Raphiolepis umbellata
Redwood, Dawn	Melasequia glyptostroboides
Rhododendron	Rhododendron spp.
Rose, spp.	Rose, spp.
Russian Olive	Elaeagnus angustifolia
Sargent Cherry	Prunus sergentii
Scotch Broom	Cytisus scoparius
Sedum	Sedum album
Serviceberry, Saskatoon	Amelanchier alnifolia
Spruce	Picea, spp.
Starice	Limonium, spp.
St. Johnswort, Golden	Hypericum frondosum
St. Johnswort, Golden	Hypericum patulum
Strawberry Tree	Arbutus unedo
Sumac, Staghorn	Rhus typhina
Sweet William	Dianthus barbatus
Syzygium	Syzygium paniculatum
Trailing African Daisy	Osteospermum fruticosum
Viburnum	Viburnum, spp.
Weddella	Weddella trilobata
Wisteria, chinese	Wisteria sinensis
Wich hazel, common	Hamamelis virginiana
Woodwaxen, common	Genista tinctoria
Wooly Thyme	Thymus pseudolanugosus
Xylosma	Xylosma congestu
Yew	Taxus, spp.

**WHERE, HOW MUCH, AND WHEN TO USE AROUND ORNAMENTALS**

Can be applied anytime of the year prior to weed seed germination to both newly transplanted and established actively growing or dormant ornamental plants where both weed control and fertilization is desired. Rainfall or overhead irrigation will improve weed control activity. Do not mix into soil. Disturbing or cultivating the soil after treatment may reduce weed control. Do not apply granules to wet foliage or under conditions in which granules will collect on leaves, if so, sprinkler-irrigate the plants to move granules off foliage and down to soil level. Do not apply in greenhouses as plant injury may likely result. Use the higher application rates for areas that have a history of heavy weed pressure. For best weed control around ornamentals first remove any existing weed growth and if possible rake, cultivate and smooth the soil surface by breaking-up clumps. Broadcast the granules evenly over the prepared soil by applying 2 lbs. to 4 lbs. a. i. per acre, equal to 200 to 400 lbs. of this product /acre or 4.6 to 9.2 lbs. per 1,000 sq.ft. 08-18-02

**NEWLY TRANSPLANTED ORNAMENTALS AND GROUND COVERS**

COMMON NAME	SCIENTIFIC NAME
Blanket Flower	Gaillardia aristata
Brassala, Dwarf	Brassala actinophylla
Camellia	Camellia japonica
Carnation	Dianthus caryophyllus
Cedar, Red	Juniperus atlantica
Coromandel	Asystasia gangetica
Dogwood, Flwering	Cornus florida
Elm	Ulmus, spp.
Forsythia	Forsythia intermedia
Gazania, Trailing	Gazania uniflora
Hackberry	Celtis occidentalis
Ivy, Algerian	Hedera canariensis
Ivy, English	Hedera helix
Hemlock, Canada	Tsuga canadensis
Holly	Ilex, spp.
Ice Plant	Carpobrotus edule
Juniper	Juniperus, spp.
Leopards - Bane	Doronicum cordatum
Metallic plant	Hemigraphis colorata
Oak	Quercus, spp.
Periwinkle, Common	Vinca minor
Periwinkle, Large	Vinca major
Pine	Pinus, spp.
Poplar	Populus, spp.
Privet	Ligustrum, spp.
Protea Species	Banksia menziesii
	Leucodendron discolor
	Leucospermum cordifolium
	Protea exinia
	Rhododendron, spp.
	Rhododendron, spp.
	Elaeagnus angustifolia
	Picea, spp.
	Osteospermum fruticosum
	Weddella trilobata
	Wisteria sinensis
	Taxus, spp.

**FOREST NURSERY APPLICATION**

COMMON NAME	SCIENTIFIC NAME
Pine, Loblolly	Pinus, reada
Pine, Eastern White	Pinus, Strobus
Pine, Slash	Pinus, Canbaea

**SENSITIVE PLANTS**

Under certain growing conditions, the plant varieties listed below may be sensitive to this product. Growers should evaluate usage on a small scale prior to large scale applications.

PLANT SPECIES	VARIETIES
Andromeda (Pieris japonica)	Snowdrift, Temple Bell
Azalea	Pink Champagne, Hershey, (Rhododendron spp.)
	Reuchtenfuer, Memoire John Hearrens, Hino Crimson, Everest, Girard
Cotoneaster	Eichholz, Himalayan, Shanghai (Cotoneaster spp.)
Crapemyrtle (Lagerstroemia indica)	Petite Red Imperial, and Seminole Bush
Heath, Spring (Erica cameo)	Mediterranean Pink
Hemlock, Canada (Tsuga canadensis)	Pendula
Hibiscus (Hibiscus Syriacus)	Red Heart
Rhododendron	Daphnoides, Nova Zembla, (Rhododendron spp.)
	Catawba Album
Spruce, Norway (Picea abies)	Pumila
Spruce, White (Picea glauca)	Densata
Viburnum (Viburnum spp)	Davidii

EPA Reg. No. 32802--TO = (79) \_\_\_\_\_  
EPA EST. NO. 32802-MO-1 \_\_\_\_\_

**NET WEIGHT (50) POUNDS**

**Howard Johnson's Enterprises, Inc.**  
700 West Virginia Street - Suite 222 - Milwaukee, WI 53204