

32802-81

10/1/2014

1/27



## U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs

Registration Division (7505P)

1200 Pennsylvania Ave., N.W.

Washington, D.C. 20460

EPA Reg. Number:

32802-81

Date of Issuance:

OCT - 1 2014

## NOTICE OF PESTICIDE:

☒ Registration☐ Reregistration

(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Howard Johnson's 0.04%

Penoxsulam Fert MF

## Name and Address of Registrant (include ZIP Code):

Howard Johnson's Enterprises, Inc.

9675 S. 60<sup>th</sup> Street

Franklin, WI 53132

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, "EPA Reg. No. 32802-81."

Signature of Approving Official:

*Kathryn V. Montague*  
Kathryn Montague, Product Manager 23  
Herbicide Branch, Registration Division (7505P)

Date:

OCT - 1 2014


2/27

3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 07/19/2014
- Alternate CSF #1 dated 07/15/2014

If you have any questions, please contact Shanta Adeeb at (703) 347-0502 or [adeeb.shanta@epa.gov](mailto:adeeb.shanta@epa.gov).



Kathryn Montague, Product Manager 23  
Herbicide Branch  
Registration Division (7505P)

[Enclosure]

3/27

[Sub Label A: Residential Uses]

## Howard Johnson's 0.04% Penoxsulam FERT MF

Alternate Brand Names: Pennington® UltraGreen® Southern Weed & Feed MF

For postemergence control of annual and perennial broadleaf weeds in established home lawns and other recreational turfgrass areas.

(See Marketing Claims Section for Additional Claims)

**Active Ingredient:**

penoxsulam: 2-(2,2-difluoroethoxy)-N-(5,8-dimethoxy[1,2,4] triazolo[1,5c]pyrimidin-2-yl)-6-(trifluoromethyl)benzenesulfonamide.....	0.04%
Other Ingredients .....	99.96%
TOTAL .....	100.00%

### KEEP OUT OF REACH OF CHILDREN CAUTION

\_\_\_\_\_ LB covers up to \_\_\_\_\_ sq ft [of turfgrass]


NET WEIGHT [XXX] LB

EPA Reg. No. 32802-TBD

EPA Est. No. 59144-GA-1<sub>A</sub>, 35497-OR-1<sub>B</sub>, 32802-MO-1<sub>C</sub>, 37351-CO-1<sub>D</sub>

Subscript used is the First letter of Run Code on Container

[See tag attached to bag for fertilizer guaranteed analysis.]



9/27

GUARANTEED ANALYSIS XX-X-X	
Total Nitrogen (N)	XX%
x.x% ammoniacal nitrogen	
x.x% urea nitrogen*	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	X%
Soluble Potash (K <sub>2</sub> O)	X%
Sulfur (S)	X%
x.x% free sulfur (S)	
Derived from: polymer-coated, sulfur-coated urea, urea, ammonium phosphate, potassium chloride. *A portion of the urea has been coated to provide x.x% coated slow-release nitrogen. This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx  Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P <sub>2</sub> O <sub>5</sub> and K <sub>2</sub> O may include the following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.	

(Optional additional guarantee statement)

[ALSO CONTAINS NON PLANT FOOD INGREDIENTS.

#### SOIL AMENDING GUARANTEED ANALYSIS

Active Ingredients:

Mycorrhizal Fungi.....1.12 props/gm  
*Glomus intraradices*..... 0.28 props/gm  
*Glomus mosseae*.....0.28 props/gm  
*Glomus etunicatum* .....0.28props/gm  
*Glomus aggregatum* .....0.28props/gm

Inert Ingredients (as a soil amendment):

(The % here needs to be adjusted for this formula which includes herbicides; e.g.)

x.xx% Fertilizer ingredients  
 x.xx% Herbicide ingredients

Lot code:  Expiration Date:
-----------------------------------

Distributed & Guaranteed by: Howard Johnson's Enterprises, Inc.  
 9675 S 60th Street, Franklin, WI 53132

5/27

---

## PRECAUTIONARY STATEMENTS

---

### Hazards to Humans and Domestic Animals

## CAUTION

**Causes Moderate Eye Irritation.**

**Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, and shoes plus socks. Wash thoroughly with soap and water after handling and before eating, drinking chewing gum, or using tobacco.**

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may contact 1-800-265-0761 for emergency medical treatment information.

### Environmental Hazards

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on the label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-265-0761.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

---

## Directions For Use

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions For Use carefully before applying.

### How This Product Works

Penoxsulam FERT 0.04% Southern MF provides postemergence control of annual, biennial and perennial broadleaf weeds in established home lawns and other recreational turfgrass areas.

### Use Precautions

- The active ingredient in this product may leave residues in treated grass clippings which could be harmful to some plants and trees. Do not collect treated grass clippings for use as mulch around plants. The applicator must notify in writing the appropriate residences/individuals of this precaution when this product is applied by a lawn service/commercial applicator.
- To minimize the potential for turfgrass injury, allow four weeks between applications.
- Avoid unnecessary disturbances, including watering and mowing, for approximately 24 hours after application.
- Use Penoxsulam FERT 0.04% Southern MF on solid (monoculture) stands of perennial ryegrass or tall fescue at a maximum rate of 1.2 lb per 1000 sqft. Some yellowing of these turfgrasses may occur within two weeks of application. Any discoloration is temporary and will dissipate under adequate growing conditions.
- Apply Penoxsulam FERT 0.04% Southern MF to newly seeded or sprigged turfgrass only after the third mowing or after the turfgrass seedlings have developed an adequate secondary root system. Reseeding is not recommended for at least three to four weeks after application.
- APPLICATION LIMITS: Certain states have limits on fertilizer application rates and timing. Consult your local State Cooperative Extension for current information on annual application limitations for [turf/lawn] fertilizers in your region.

### Use Restrictions

- Do not apply Penoxsulam FERT 0.04% Southern MF on dichondra, carpetgrass, bahiagrass, or turfgrass containing desirable clovers.
- Do not apply Penoxsulam FERT 0.04% Southern MF directly to, or otherwise permit Penoxsulam FERT 0.04% Southern MF to come into contact with flowers, ornamental shrubs or trees, or other desirable broadleaf plants, as serious plant injury may occur.
- Do not apply to centipedegrass during the period of winter dormancy. Grass should be green and actively growing at application.
- Do not apply more than 3.4/[X] lb of Penoxsulam FERT 0.04% Southern MF per 1000 sq ft per application. (to allow for state restrictions)
- (Note: If registered in the state of Florida, any of the following statements may be added as appropriate)
  - In Florida, do not exceed X applications per year on Bahiagrass in the North, and X per year in Central and South Florida.
  - In Florida, do not exceed X applications per year on Bermudagrass in the North, or X applications per year in Central and X applications per year in South Florida.
  - In Florida, do not exceed X applications per year on Centipedegrass in the North, or X applications per year in Central and South Florida.
  - In Florida, do not exceed X applications per year on St. Augustinegrass in the North, or X applications per year in Central and X applications per year in South Florida.
  - In Florida, do not exceed X applications per year on Zoysiagrass in the North, or X applications per year in Central and South Florida.
  - (Note: may also be expressed in tabular format.)

- In the state of Florida: Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

### **Treatment on Turfgrass Species Not Listed on the Label for Penoxsulam FERT 0.04% Southern MF**

Users who wish to use Penoxsulam FERT 0.04% Southern MF on turfgrass species not listed on this label may determine the suitability for use by treating a small area at a specified rate. Prior to treatment of larger areas, the treated area should be observed for any sign of herbicidal injury during 30 days of normal growing conditions to determine if the target turfgrass species is tolerant. Applications should not be made to turfgrass under stress. The user assumes responsibility for any plant damage or other liability resulting from the use of Penoxsulam turfgrass species not listed below on this label.

### **Weeds Controlled or Suppressed**

<b>Weeds Controlled or Suppressed</b>		<b>Rate</b>
		<b>lb/1000 sq ft</b>
burdock burnweed, American buttercup, hairy chickweed, common chickweed, mouseear clover, hop clover, red clover, white cocklebur, common coffeeweed common vetch creeping beggarweed curly dock dandelion, common dollarweed dwarf beggarweed English lawn daisy false dandelion Florida betony goldenrod hawkweed henbit ivy, ground	lambsquarters lespedenza, common matchweed medic, black parsley-piert plantain, broadleaf plantain, buckhorn purslane, common ragweed, common round leaf mallow sheep sorrel shepherd's purse smartweed speedwell, common spotted catsear spurge, spotted spurweed Virginia buttonweed <sup>1</sup> Virginia pepperweed wild violet woodsorrel, creeping woodsorrel, yellow	<b>2.5</b>
Kyllinga, annual and green (suppression only) <sup>1</sup>	Nutsedge, purple and yellow (suppression only) <sup>1</sup>	

<sup>1</sup> Re-treatment may be necessary

### **How Much to Apply**

**Penoxsulam FERT 0.04% Southern MF is recommended for use on the following established turfgrass species** (numbers in parentheses (-) refer to footnotes below):

**Established Cool Season Turfgrass (1)**

annual bluegrass  
bentgrass  
bluegrass, Kentucky  
fescue, chewing  
fescue, creeping red  
fescue, sheeps  
fescue, tall (2)  
ryegrass, perennial (2)

**Established Warm Season Turfgrass (3)**

bermudagrass (common or hybrid)  
centipedegrass (4)  
fescue, tall (growing in warm season areas) (2)  
kikuyugrass  
seashore paspalum  
St. Augustinegrass  
zoysiagrass

1. Use no more than 2.3 lb of Penoxsulam FERT 0.04% Southern MF per 1000 sq. ft. per application.
2. Use Penoxsulam FERT 0.04% Southern MF on these species at a maximum rate of 1.2 lb per 1000 sq. ft. per application and only if some injury can be tolerated. A slight yellowing of turfgrass may occur within two weeks of application. The discoloration is temporary and is generally gone within four weeks of treatment under adequate growing conditions.
3. Use no more than 3.4 lb of Penoxsulam FERT 0.04% Southern MF per 1000 sq. ft. per application.
4. Do not apply to centipedegrass during the period of winter dormancy. Grass should be green and actively growing at application.

Apply Penoxsulam FERT 0.04% Southern MF using a drop or rotary-type spreader designed to apply granules. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.

To optimize product performance, mow the turfgrass two to three days before applying Penoxsulam FERT 0.04% Southern MF. If the turfgrass is under moisture stress, a thorough sprinkler irrigation of at least ½ inch may be needed at least one to two days before application. For optimal results, apply Penoxsulam FERT 0.04% Southern MF in the



9/27

morning when dew is present. If necessary, sprinkler irrigate with less than ¼ inch of water just prior to application to maximize particle adherence to leaf surfaces. Do not water treated area for 24 hours following application. Do not make applications if rainfall is expected within 24 hours after treatment.

### Application Techniques for Applying Penoxsulam FERT 0.04% Southern MF

- When using a drop-type spreader, a splashboard mounted under the hopper will provide more even granule distribution.
- A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.
- For treating smaller areas, a hand held or push-type rotary applicator such as a whirlybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed.
- A shaker-type applicator made from a small container with holes punched in the bottom is recommended for small, difficult to treat areas. Carefully measure the amount of product needed to avoid over application.

### Spreader Settings as a Guide for Calibration

**Note:** These settings are provided as a suggested starting point in calibrating each individual spreader. Since no two spreaders are alike, these settings should be considered as approximate and are not intended to be used as absolute recommendations by Howard Johnson's Enterprises, Inc.

#### Optional Spreader Chart

Spreader	Setting Delivers 2.5/[X] lb./1000 sq.ft.
Scotts® Turf Builder® Classic Drop	X
Scotts® Turf Builder® Edgeguard® Mini	X
Sta-Green® Broadcast	X
Pennington® Broadcast	X
[Additional spreader brands]	X

### How Much to Apply

To apply Penoxsulam FERT 0.04% Southern MF, use the rate appropriate for the established turfgrass (Cool Season or Warm Season) in your lawn. Possible re-treatment in approximately 30 days may be necessary if turfgrass is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed specified rates because damage to turfgrass may occur. Do not make more than two applications per season.

**Note:** When applying to warm season turfgrass under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor temporary discoloration. Reduce application rates by 50% during hot, dry conditions to avoid possible dehydration or browning of grasses.

10/27

**Broadcast Rates for Penoxsulam FERT 0.04% Southern MF**

Bag Size	Rate	Treated Area
Lb	lb/1000 sq ft	sq ft
12.5	2.5	5,000
18.75	2.5	7,500
37.5	2.5	15,000
XXX	2.5	XXXX

**When to Apply**

Best weed control results are obtained when Penoxsulam FERT 0.04% Southern MF is applied to small, actively growing weeds in the spring or early fall. Extreme growing conditions such as drought or near freezing temperatures prior to, at, or following time of application may reduce weed control. Broadleaf weed species germinate at different times.

This product may also be applied to newly established warm season turfgrass. Delay treatments four weeks after sodding, sprigging or plugging.

**Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container, in a cool, dry place. inaccessible to children and domestic animals.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

**If empty:** Place in trash or offer for recycling if available.

**If partly filled:** Call your local solid waste agency or 1-800-265-0761 for disposal instructions. Never place unused product down any indoor or outdoor drain.

**Warranty**

The directions for use of this product are believed to be adequate and must be followed carefully, it is impossible to eliminate all risks inherently associated with the use of this product. Plant injury, ineffectiveness, or other unintended consequences may result due to such factors as weather conditions, presence or absence of other materials, or the manner of use or application, all of which are beyond the control of Howard Johnson's Enterprises, Inc. the manufacturer, or the seller.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth above.

To the extent permitted by law, Howard Johnson's Enterprises, Inc, the manufacturer, or the seller makes no other warranties, guarantees, or representations of any kind to the buyer or the user, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold or use of the product, including, but not limited to, merchantability, fitness for a particular purpose or use, or eligibility of the product for

11/27

any particular trade usage. To the extent consistent with applicable law, Buyer's or user's exclusive remedy, and Howard Johnson's Enterprises, Inc the manufacturer's or the seller's total liability shall be limited to damages not exceeding the cost of the product. No agent or employee of Howard Johnson's Enterprises, Inc, or the seller is authorized to amend the terms of this warranty disclaimer or the product's label or to make a presentation or recommendation different from or inconsistent with the label of this product.

To the extent consistent with applicable law, Howard Johnson's Enterprises, Inc, the manufacturer, or the seller shall not be liable for consequential, special, or indirect damages resulting from the use, handling, application, storage, or disposal of this product or for damages in the nature of penalties, and the buyer and the user waive any right that they may have to such damages.

**Fertilizer Labeling Requirements by Association of American Plant Food Control Officials (AAPFCO):**

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-xx.com>

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials. Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>.

Certain states have limits on fertilizer application rates and timing. Consult your State Cooperative Extension for current information on annual application limitations for lawn fertilizers in your region.

Information regarding the raw materials composing this product can be obtained by writing to the Guarantor listed on this label and referring to the batch number found on this container.

**APPENDIX FOR LABEL A**

**1. Advertising claims and marketing phrases that may be presented on promotional materials and the container labels of supplemental registrants.**

**Herbicide Benefit Claims**

- Kills Tough Existing Weeds and Controls New Weeds for 3 Months!
- Controls tough weeds [such as] [dollarweed] [clover] [dandelion] [henbit] [and chickweed]
- Use for a beautiful weed-free lawn
- Effective weed control leads to greener, thicker lawns
- Apply anytime when weeds are actively growing

- Kills clover, dollarweed, dandelion, henbit & chickweed
- Controls a wide variety of existing broadleaf weeds [including dollarweed, dandelion and clover] while feeding your grass.

#### **Fertilizer Benefit Claims**

- [Feeds] [Greens] [Feeds and greens] for 3 months [guaranteed] (to be used only on formulations with controlled release technology)
- Feeds with slow release fertilizer for 3 months (to be used only on formulations with controlled release technology)
- Quick Greening [and Extended Feeding] (to be used only on formulations with controlled release technology)
- Helps strengthen lawn against heat & drought
- [Rich in] Nitrogen for a lush, green lawn
- [Enriched with] Potassium for overall plant vigor
- Phosphorus Free [ - Helping reduce Phosphorus in water sources] (to be used only on formulations containing 0% phosphorus)
- Phosphorus Free-Helping Reduce Phosphorus Contamination of Water Sources
- [Contains] Nitrogen for lush, green plant growth
- [Contains] Potassium for [overall plant vigor] [drought & disease resistance]
- Enhanced Efficiency Fertilizer - [feed more lawn with less fertilizer] (to be used only on formulations with enhanced efficiency nitrogen)
- You only need XX pounds [of fertilizer] to cover XXXX square feet
- Promotes thick, full grass
- Delivers nutrients more efficiently than ordinary fertilizers (to be used only on formulations with enhanced efficiency nitrogen)
- Allows grass to use more of the fertilizer you buy & apply (to be used only on formulations with enhanced efficiency nitrogen)
- Contains slow release nitrogen for extended [greening] [feeding]

#### **Fungi Benefit Claims**

- Boosts Nutrient Uptake for a Lush, Green Lawn
- Increases Drought Tolerance & Reduces Watering Needs by Extending Root Absorption Area
- Increased root absorption area for more efficient use of water in [the] soil
- Allows grass to use more of the fertilizer you buy and apply
- [Contains mycorrhizal fungi for] enhanced nutrient uptake
- Increased root absorption area to protect against heat & drought
- Increase absorption [area of roots, of fertilizer]
- Enhance nutrient uptake
- The only brand with MYCO Advantage™ for better results than [fertilizer alone, ordinary fertilizer]

## Marketing Claims

- [For use on residential lawns]
- NOTICE- This product contains chemicals known to the State of California to cause birth defects or other reproductive harm
- Covers up to XXXX sq.ft.

- **The Pennington® UltraGreen® Difference**

Pennington® UltraGreen® products are more efficient than other fertilizers. Half of the nitrogen in this premium line\* is stabilized, which allows grass to use more of the nutrients you put down and reduces nutrient loss to the environment. The other half of the nitrogen is slow-release for extended feeding for 3 months.

Additionally, these unique fertilizer blends contain mycorrhizal fungi, natural organisms that have been present in vigorous soils for millions of years. These microscopic fungi increase the surface absorbing area of roots, greatly improving the ability of the lawn to access vital soil resources such as water and nutrients.

These tiny fertilizer granules are packed with essential plant nutrients and fungi, so you can feed more lawn with less fertilizer.

\*excluding Pennington UltraGreen Starter Fertilizer

- **Benefits of Pennington® UltraGreen® Southern MF Weed & Feed 30-0-4**

- Promotes thick, green grass
- Half of the total nitrogen in this product is slow-release and feeds for 3 months [guaranteed]
- Half of the total nitrogen is stabilized for less leaching into the environment
- Phosphorus-free
- Pennington® UltraGreen® Southern MF Weed & Feed 30-0-4 **WILL NOT BURN** when used as directed. We're so confident you'll love the results of using this product that it comes with the **Pennington Satisfaction Guarantee**. †

† Pennington Satisfaction Guarantee: If, after following the instructions, you are not satisfied with the product results, send your name, address, and proof of purchase (original dated receipt) to:  
 Pennington Refund  
 1000 Parkwood Circle, Suite 700  
 Atlanta, GA 30339

14/27

[Sub Label B: Commercial Uses]

## Howard Johnson's 0.04% Penoxsulam FERT MF

For postemergence control of annual and perennial broadleaf weeds in established turfgrass, including residential lawns, golf courses, parks, sports fields, sod farms and around commercial buildings and other recreational and commercial turfgrass areas.

(See Marketing Claims Section for Additional Claims)

**Active Ingredient:**

penoxsulam: 2-(2,2-difluoroethoxy)-N-(5,8-dimethoxy[1,2,4] triazolo[1,5c]pyrimidin-2-yl)-6-(trifluoromethyl)benzenesulfonamide.....	0.04%
Other Ingredients .....	99.96%
TOTAL .....	100.00%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

\_\_\_\_\_ LB covers up to \_\_\_\_\_ sq ft [of turfgrass]


NET WEIGHT XXX LB

EPA Reg. No. 32802-TBD

EPA Est. No. 59144-GA-1<sub>A</sub>, 35497-OR-1<sub>B</sub>, 32802-MO-1<sub>C</sub>, 37351-CO-1<sub>D</sub>

Subscript used is the First letter of Run Code on Container

[See tag attached to bag for fertilizer guaranteed analysis.]



15/27

GUARANTEED ANALYSIS XX-X-X	
Total Nitrogen (N)	XX%
x.x% ammoniacal nitrogen	
x.x% urea nitrogen*	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	X%
Soluble Potash (K <sub>2</sub> O)	X%
Sulfur (S)	X%
x.x% free sulfur (S)	
Derived from: polymer-coated, sulfur-coated urea, urea, ammonium phosphate, potassium chloride. *A portion of the urea has been coated to provide x.x% coated slow-release nitrogen. This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx  Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P <sub>2</sub> O <sub>5</sub> and K <sub>2</sub> O may include the following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.	

(Optional additional guarantee statement)

[ALSO CONTAINS NON PLANT FOOD INGREDIENTS.

SOIL AMENDING GUARANTEED ANALYSIS

Active Ingredients:

Mycorrhizal Fungi.....1.12 props/gm  
*Glomus intraradices*..... 0.28 props/gm  
*Glomus mosseae*.....0.28 props/gm  
*Glomus etunicatum* .....0.28props/gm  
*Glomus aggregatum* .....0.28props/gm

Inert Ingredients (as a soil amendment):

(The % here needs to be adjusted for this formula which includes herbicides; e.g.)

x.xx% Fertilizer ingredients

x.xx% Herbicide ingredients

Lot code:  Expiration Date:
-----------------------------------

Distributed & Guaranteed by: Howard Johnson's Enterprises, Inc.  
9675 S 60th Street, Franklin, WI 53132

---

## PRECAUTIONARY STATEMENTS

---

### Hazards to Humans and Domestic Animals

**Causes Moderate Eye Irritation.**

**Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking chewing gum, or using tobacco.**

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Directions

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may contact 1-XXX-XXX-XXXX for emergency medical treatment information.

### Environmental Hazards

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

---

### Directions For Use

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions For Use carefully before applying.



Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on the label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-XXX-XXX-XXXX.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

**Non-Agricultural Use Requirements**

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Entry Restrictions for Non-WPS Uses:** Keep unprotected persons out of treated areas until dusts have settled.

**Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in cool dry place in original container.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Nonrefillable rigid containers 5 gallons or less:**

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Nonrefillable nonrigid containers:**

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Refillable rigid containers larger than 5 gal**

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Nonrefillable rigid containers larger than 5 gal:**

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Tip container on side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse** as follows: Empty the remaining contents into application equipment or a

mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

### Information

Penoxsulam FERT 0.04% Southern MF provides postemergence control of annual, biennial and perennial broadleaf weeds in established turfgrass, including residential lawns, golf courses, parks, sports fields, sod farms and around commercial buildings and other recreational and commercial turfgrass areas.

### Use Precautions

- The active ingredient in this product may leave residues in treated grass clippings which could be harmful to some plants and trees. Do not collect treated grass clippings for use as mulch around plants. The applicator must notify in writing the appropriate residences/individuals of this precaution when this product is applied by a lawn service/commercial applicator.
- To minimize the potential for turfgrass injury, allow four weeks between applications.
- Avoid unnecessary disturbances, including watering and mowing, for approximately 24 hours after application.
- Use Penoxsulam FERT 0.04% Southern MF on solid (monoculture) stands of perennial ryegrass or tall fescue at a maximum rate of 50 lb (0.02 lb ai penoxsulam) per acre. Some yellowing of these turfgrasses may occur within two weeks of application. Any discoloration is temporary and will dissipate under adequate growing conditions.
- Apply Penoxsulam FERT 0.04% Southern MF to newly seeded or sprigged turfgrass only after the third mowing or after the turfgrass seedlings have developed an adequate secondary root system. Reseeding is not recommended for at least three to four weeks after application.
- In the state of Florida, this product should be applied at rates not exceeding those recommended in the document titled Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries.
- In the state of Florida: For application to sports turf (non-agricultural land planted exclusively for golf courses, parks, and athletic fields) follow "SL191, Recommendations for N, P, K and Mg for Golf Course & Athletic Field Fertilization Base on Mehlich | Extractant" available at <http://edis.ifas.ufl.edu/SS404>, or "BMP's for Enhancement of Environmental Quality on Florida's Golf Courses" available at <http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/glfbmp07.pdf>
- In the state of Florida: For application to urban turf (non-agricultural land planted in closely mowed, managed grasses except golf courses, parks and athletic fields) follow "Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries" available at [http://www.dep.state.fl.us/central/Home/MeetingsTraining/FLGreen/BMP\\_Book\\_final.pdf](http://www.dep.state.fl.us/central/Home/MeetingsTraining/FLGreen/BMP_Book_final.pdf)

- **APPLICATION LIMITS:** Certain states have limits on fertilizer application rates and timing. Consult your local State Cooperative Extension for current information on annual application limitations for [turf/lawn] fertilizers in your region.

#### **Use Restrictions**

- Do not apply Penoxsulam FERT 0.04% Southern MF on dichondra, carpetgrass, bahiagrass, or turfgrass containing desirable clovers.
- Do not apply Penoxsulam FERT 0.04% Southern MF directly to, or otherwise permit Penoxsulam FERT 0.04% Southern MF to come into contact with flowers, ornamental shrubs or trees, or other desirable broadleaf plants, as serious plant injury may occur.
- Do not apply to centipedegrass during the period of winter dormancy. Grass should be green and actively growing at application.
- Do not apply more than 150 lb/[X lb] of Penoxsulam FERT 0.04% Southern MF (0.06 lb ai penoxsulam) per acre per application. (to allow for state restrictions)
- (Note: If registered in the state of Florida, any of the following statements may be added as appropriate)
  - In Florida, do not exceed X applications per year on Bahiagrass in the North, and X per year in Central and South Florida.
  - In Florida, do not exceed X applications per year on Bermudagrass in the North, or X applications per year in Central and X applications per year in South Florida.
  - In Florida, do not exceed X applications per year on Centipedegrass in the North, or X applications per year in Central and South Florida.
  - In Florida, do not exceed X applications per year on St. Augustinegrass in the North, or X applications per year in Central and X applications per year in South Florida.
  - In Florida, do not exceed X applications per year on Zoysiagrass in the North, or X applications per year in Central and South Florida.
  - (Note: may also be expressed in tabular format.)
- In the state of Florida: Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

### Treatment on Turfgrass Species Not Listed on the Label for Penoxsulam FERT 0.04% Southern MF

Users who wish to use Penoxsulam FERT 0.04% Southern MF on turfgrass species not listed on this label may determine the suitability for use by treating a small area at a specified rate. Prior to treatment of larger areas, the treated area should be observed for any sign of herbicidal injury during 30 days of normal growing conditions to determine if the target turfgrass species is tolerant. Applications should not be made to turfgrass under stress. The user assumes responsibility for any plant damage or other liability resulting from the use of Penoxsulam FERT 0.04% Southern MF on turfgrass species not listed below on this label.

### Weeds Controlled or Suppressed

Weeds Controlled or Suppressed		Rate	
		lb/acre	lb/1000 sq ft
Burdock burnweed, American buttercup, hairy chickweed, common chickweed, mouseear clover, hop clover, red clover, white cocklebur, common coffeeweed common vetch creeping beggarweed curly dock dandelion, common dollarweed dwarf beggarweed English lawn daisy false dandelion Florida betony goldenrod hawkweed henbit ivy, ground	lambsquarters lespedenza, common matchweed medic, black parsley-piert plantain, broadleaf plantain, buckhorn purslane, common ragweed, common round leaf mallow sheep sorrel shepherd's purse smartweed speedwell, common spotted catsear spurge, spotted spurweed Virginia buttonweed <sup>1</sup> Virginia pepperweed wild violet woodsorrel, creeping woodsorrel, yellow	104	2.5
Kyllinga, annual and green (suppression only) <sup>1</sup>	Nutsedge, purple and yellow (suppression only) <sup>1</sup>	108	2.5

<sup>1</sup> Re-treatment may be necessary

### Application Directions

**Penoxsulam FERT 0.04% Southern MF is recommended for use on the following established turfgrass species (numbers in parentheses (-) refer to footnotes below):**

**Established Cool Season Turfgrass (1)**

<b>Common Name</b>	<b>Scientific Name</b>
annual bluegrass	<i>Poa annua</i>
bentgrass	<i>Agrostis</i> species
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, chewing	<i>Festuca rubra</i> var. <i>commutata</i>
fescue, creeping red	<i>Festuca rubra</i>
fescue, sheeps	<i>Festuca ovina</i>
fescue, tall (2)	<i>Festuca arundinaceae</i>
ryegrass, perennial (2)	<i>Lolium perenne</i>

**Established Warm Season Turfgrass (3)**

<b>Common Name</b>	<b>Scientific Name</b>
bermudagrass (common or hybrid)	<i>Cynodon dactylon</i>
centipedegrass (4)	<i>Eremochloa ophiuroides</i>
Fescue, tall (growing in warm season areas) (2)	<i>Festuca arundinaceae</i>
kikuyugrass	<i>Pennisetum clandestinum</i>
seashore paspalum	<i>Paspalum vaginatum</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
zoysiagrass	<i>Zoysia japonica</i>
zoysiagrass	<i>Zoysia tenuifolia</i>

1. Use no more than 100 lb of Penoxsulam FERT 0.04% Southern MF (0.045 lb ai penoxsulam) per acre per application.
2. Use Penoxsulam FERT 0.04% Southern MF on these species at a maximum rate of 50 lb (0.02 lb ai penoxsulam) per acre per application and only if some injury can be tolerated. A slight yellowing of turfgrass may occur within two weeks of application. The discoloration is temporary and is generally gone within four weeks of treatment under adequate growing conditions.
3. Use no more than 150 lb of Penoxsulam FERT 0.04% Southern MF (0.06 lb ai penoxsulam) per acre per application.
4. Do not apply to centipedegrass during the period of winter dormancy. Grass should be green and actively growing at application.

Apply Penoxsulam FERT 0.04% Southern MF using a drop or rotary-type spreader designed to apply granules. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.

To optimize product performance, mow the turfgrass two to three days before applying Penoxsulam FERT 0.04% Southern MF. If the turfgrass is under moisture stress, a thorough sprinkler irrigation of at least ½ inch may be needed at least one to two days before application. For optimal results, apply Penoxsulam FERT 0.04% Southern MF in the

morning when dew is present. If necessary, sprinkler irrigate with less than ¼ inch of water just prior to application to maximize particle adherence to leaf surfaces. Do not water treated area for 24 hours following application. Do not make applications if rainfall is expected within 24 hours after treatment.

#### Application Techniques for Applying Penoxsulam FERT 0.04% Southern MF

- When using a drop-type spreader, a splashboard mounted under the hopper will provide more even granule distribution.
- A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.
- For treating smaller areas, a hand held or push-type rotary applicator such as a whirlybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed.
- A shaker-type applicator made from a small container with holes punched in the bottom is recommended for small, difficult to treat areas. Carefully measure the amount of product needed to avoid over application.

#### Spreader Settings as a Guide for Calibration

**Note:** These settings are provided as a suggested starting point in calibrating each individual spreader. Since no two spreaders are alike, these settings should be considered as approximate and are not intended to be used as absolute recommendations by Howard Johnson's Enterprises, Inc.

#### Optional Spreader Chart

Spreader	Setting for Spreader
	Rate of Penoxsulam FERT 0.04% Southern 104 lb/[X lb]/acre
Spreader Brands	XX

#### Rates and Timing of Application

Apply 104 lb of Penoxsulam FERT 0.04% Southern MF per acre (2.5 lb per 1000 sq ft) per application to established turfgrass. Possible re-treatment in approximately 30 days may be necessary if turfgrass is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed 225 lb of Penoxsulam FERT 0.04% Southern MF per acre (5.4 lb per 1000 sq ft) because damage to turfgrass may occur.

**Note:** When applying to warm season turfgrass under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor temporary discoloration. Reduce application rates by 50% during hot, dry conditions to avoid possible dehydration or browning of grasses.

**Broadcast Rates for Penoxsulam FERT 0.04% Southern MF**

Bag Size	Rate		Treated Area	
	Lb/acre	lb/1000 sq ft	Acre	sq ft
12.5	104	2.5	0.11	5,000
18.75	104	2.5	0.17	7,500
37.5	104	2.5	0.34	15,000
XXX	XXX	XXX	XXX	XXX

Best weed control results are obtained when Penoxsulam FERT 0.04% Southern MF is applied to small, actively growing weeds in the spring or early fall. Extreme growing conditions such as drought or near freezing temperatures prior to, at, or following time of application may reduce weed control. Broadleaf weed species germinate at different times.

This product may also be applied to newly established warm season turfgrass. Delay treatments four weeks after sodding, sprigging or plugging.

**Warranty**

The directions for use of this product are believed to be adequate and must be followed carefully, it is impossible to eliminate all risks inherently associated with the use of this product. Plant injury, ineffectiveness, or other unintended consequences may result due to such factors as weather conditions, presence or absence of other materials, or the manner of use or application, all of which are beyond the control of Howard Johnson's Enterprises, Inc. the manufacturer, or the seller.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth above.

To the extent permitted by law, Howard Johnson's Enterprises, Inc, the manufacturer, or the seller makes no other warranties, guarantees, or representations of any kind to the buyer or the user, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold or use of the product, including, but not limited to, merchantability, fitness for a particular purpose or use, or eligibility of the product for any particular trade usage. To the extent consistent with applicable law, Buyer's or user's exclusive remedy, and Howard Johnson's Enterprises, Inc the manufacturer's or the seller's total liability shall be limited to damages not exceeding the cost of the product. No agent or employee of Howard Johnson's Enterprises, Inc, or the seller is authorized to amend the terms of this warranty disclaimer or the product's label or to make a presentation or recommendation different from or inconsistent with the label of this product.



To the extent consistent with applicable law, Howard Johnson's Enterprises, Inc, the manufacturer, or the seller shall not be liable for consequential, special, or indirect damages resulting from the use, handling, application, storage, or disposal of this product or for damages in the nature of penalties, and the buyer and the user waive any right that they may have to such damages.

**Fertilizer Labeling Requirements by Association of American Plant Food Control Officials (AAPFCO):**

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-xx.com>

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials. Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>.

Certain states have limits on fertilizer application rates and timing. Consult your State Cooperative Extension for current information on annual application limitations for lawn fertilizers in your region.

Information regarding the raw materials composing this product can be obtained by writing to the Guarantor listed on this label and referring to the batch number found on this container.

**APPENDIX FOR LABEL A**

**1. Advertising claims and marketing phrases that may be presented on promotional materials and the container labels of supplemental registrants.**

**Herbicide Benefit Claims**

- Kills Tough Existing Weeds and Controls New Weeds for 3 Months!
- Controls tough weeds [such as] [dollarweed] [clover] [dandelion] [henbit] [and chickweed]
- Use for a beautiful weed-free lawn
- Effective weed control leads to greener, thicker lawns
- Apply anytime when weeds are actively growing
- Kills clover, dollarweed, dandelion, henbit & chickweed
- Controls a wide variety of existing broadleaf weeds [including dollarweed, dandelion and clover] while feeding your grass.

### **Fertilizer Benefit Claims**

- [Feeds] [Greens] [Feeds and greens] for 3 months [guaranteed] (to be used only on formulations with controlled release technology)
- Feeds with slow release fertilizer for 3 months (to be used only on formulations with controlled release technology)
- Quick Greening [and Extended Feeding] (to be used only on formulations with controlled release technology)
- Helps strengthen lawn against heat & drought
- [Rich in] Nitrogen for a lush, green lawn
- [Enriched with] Potassium for overall plant vigor
- Phosphorus Free [ - Helping reduce Phosphorus in water sources] (to be used only on formulations containing 0% phosphorus)
- Phosphorus Free-Helping Reduce Phosphorus Contamination of Water Sources
- [Contains] Nitrogen for lush, green plant growth
- [Contains] Potassium for [overall plant vigor] [drought & disease resistance]
- Enhanced Efficiency Fertilizer - [feed more lawn with less fertilizer] (to be used only on formulations with enhanced efficiency nitrogen)
- You only need XX pounds [of fertilizer] to cover XXXX square feet
- Promotes thick, full grass
- Delivers nutrients more efficiently than ordinary fertilizers (to be used only on formulations with enhanced efficiency nitrogen)
- Allows grass to use more of the fertilizer you buy & apply (to be used only on formulations with enhanced efficiency nitrogen)
- Contains slow release nitrogen for extended [greening] [feeding]

### **Fungi Benefit Claims**

- Boosts Nutrient Uptake for a Lush, Green Lawn
- Increases Drought Tolerance & Reduces Watering Needs by Extending Root Absorption Area
- Increased root absorption area for more efficient use of water in [the] soil
- Allows grass to use more of the fertilizer you buy and apply
- [Contains mycorrhizal fungi for] enhanced nutrient uptake
- Increased root absorption area to protect against heat & drought
- Increase absorption [area of roots, of fertilizer]
- Enhance nutrient uptake
- The only brand with MYCO Advantage™ for better results than [fertilizer alone, ordinary fertilizer]

## Marketing Claims

- NOTICE- This product contains chemicals known to the State of California to cause birth defects or other reproductive harm
- Covers up to XXXX sq.ft.

- **The Pennington® UltraGreen® Difference**

Pennington® UltraGreen® products are more efficient than other fertilizers. Half of the nitrogen in this premium line\* is stabilized, which allows grass to use more of the nutrients you put down and reduces nutrient loss to the environment. The other half of the nitrogen is slow-release for extended feeding for 3 months.

Additionally, these unique fertilizer blends contain mycorrhizal fungi, natural organisms that have been present in vigorous soils for millions of years. These microscopic fungi increase the surface absorbing area of roots, greatly improving the ability of the lawn to access vital soil resources such as water and nutrients.

These tiny fertilizer granules are packed with essential plant nutrients and fungi, so you can feed more lawn with less fertilizer.

\*excluding Pennington UltraGreen Starter Fertilizer

- **Benefits of Pennington® UltraGreen® Southern MF Weed & Feed 30-0-4**

- Promotes thick, green grass
- Half of the total nitrogen in this product is slow-release and feeds for 3 months [guaranteed]
- Half of the total nitrogen is stabilized for less leaching into the environment
- Phosphorus-free
- Pennington® UltraGreen® Southern MF Weed & Feed 30-0-4 **WILL NOT BURN** when used as directed. We're so confident you'll love the results of using this product that it comes with the **Pennington Satisfaction Guarantee**. †

† Pennington Satisfaction Guarantee: If, after following the instructions, you are not satisfied with the product results, send your name, address, and proof of purchase (original dated receipt) to:

Pennington Refund  
1000 Parkwood Circle, Suite 700  
Atlanta, GA 30339