

Airysol Brand Moth Sentry

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water and other aquatic organisms. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE USE.

Affix trigger, pump or other type sprayer head to container for indoor application. Do not transfer to another container. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before spraying or surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry. SHAKE WELL BEFORE USING.

INDOOR APPLICATION: For spot application only. Do not use as a space spray indoors. Do not apply to classrooms while in use. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed. Do not use in commercial food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed.

SURFACE SPRAYING: For 60-Day (Two Months) Control of Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetles, Meal Worms, Grain Mites and Cadelles: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For silverfish: Spray bookcases. For ants: Spray trails, nests and points of entry. **To control Carpet Beetles:** Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. **To control Bedbugs:** Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. **To control Fleas (adults and larvae) and Ticks, including Ticks which may carry Lyme disease (adults and larvae):** Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. **To Allow to dry thoroughly before allowing pets to contact treated areas. To control Clothes Moths:** Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining. **To kill Lice and Louse Eggs:** Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

ON LIVESTOCK: TO PROTECT CATTLE (BEEF AND DAIRY), GOATS, SHEEP, HOGS AND HORSES FROM HORN FLIES, FACE FLIES, HOUSE FLIES, MOSQUITOES AND GNATS: Apply a light mist sufficient to wet the surface of the hair. For face flies: Spray face and head, but do not spray into eyes. **To control Stable Flies, Horse Flies and Deer Flies:** Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily. **To control Blood-sucking Lice:** Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required. **To control Poultry**



Moth Sentry

- For Domestic Use both Indoors and Outdoors
- A Ready-to-Use Insecticide
- For Control of Fleas, Brown Dog Ticks and Lice on Premises and on Dogs
- For Control of Deer Ticks which may cause Lyme disease on Premises
- For Control of Pests on Livestock and Livestock Premises
- Control Fleas and Ticks on Dogs for 14 Days
- Provides 60-day (Two Months) Control of Cockroaches
- Provides 60-day (Two Months) Control of Fleas
- Provides 60-day (Two Months) Control of Stored Product Pests

ACTIVE INGREDIENTS:

*Permethrin: 0.50%

INERT INGREDIENTS: 99.50%

TOTAL: 100.00%

*(3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2-dichloroethenyl) 2,2-dimethylcyclopropanecarboxylate. Cis/trans ratio: min. 35% (+/-) cis and max. 65% (+/-) trans.

ACCEPTED
FEB 9 2004

KEEP OUT OF REACH OF CHILDREN.

CAUTION: Federal Insecticide, Fungicide, and Rodenticide Act.

See back panel for additional precautions.

33176-44

NET CONTENTS 1 GALLON (3.75 LITERS)

Lice: Spray roasts, walls and other surfaces thoroughly. The roasts followed by spraying over the birds with a fine mist. **To control Bedbugs and Mites in poultry houses:** Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

ON DOGS: To control Fleas, Ticks, and Lice: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Use protective gloves or mitts when applying product. **NOTE:** Do not treat puppies less than 12 weeks old. **USE ONLY ON DOGS. DO NOT USE ON CATS OR KITTENS.** Do not use on old, sick or debilitates animals. Consult a veterinarian before using this product on pregnant, medicated or nursing animals. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

OUTDOOR APPLICATION: TO CONTROL TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (PERIMETER TREATMENT): Use this product as a residual treatment to control Ticks including the Deer Tick (*Ixodes dammini*), Lone Star Tick (*Amblyomma americanum*) Brown Dog Tick (*Rhipicephalus sanguineus*), American Dog Tick (*Dermacentor variabilis*) and Western Black-legged Tick (*Ixodes pacificus*): For each area listed below, apply until surface is wet (approximately 1 gal/750-100 ft²). Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways and in crawl spaces where there pests may enter. Treat a band of vegetation and leaf litter 6 to 10 feet wide around and the outside of buildings to help control ticks occurring there. To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

FOR CONTROL OF GARDEN INSECTS (Not for Use on plants grown for sale or other commercial or research purposes. Not for use in commercial greenhouses or nurseries): For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray as well as Whiteflies, Aphids, Exposed Thrips, Spiders, Red Mites and Leafminers. May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas. May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurals, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias. **Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Armyworms:** When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. **STORAGE & DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. **CONTAINER DISPOSAL:** Do not reuse this container IF EMPTY: Place in trash or offer for recycling if available. IF PARTLY FILLED: Never place unused product down any indoor or outdoor drain. Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.

EPA REG. NO. 33176-44

EPA EST. NO. 10807-TX-1

01/04p

FOR EMERGENCY MEDICAL INFORMATION
CHEM • TEL, INC. 1-800-255-3924



MANUFACTURED BY:
MBL INDUSTRIES, INC.
a solely owned subsidiary of Amrep, Inc.
990 Industrial Park Drive
Marietta, GA 30062