

TURF TONIC WEED and FEED

Kills weeds as it feeds your lawn.

Weed and feed your lawn in one easy application.

Fast and easy way to feed your lawn and kill weeds too.

Specially formulated to kill: DANDELIONS, PLANTAIN, and FOXTAIL.

This weed killer also kills:

BUCKHORN PLANTAIN

COMMON CHICKWEED

DOCK

NETTLE

PIGWEEED

RAGWEED

THISTLES

WILD CARROT

WILD GARLIC

WILD MUSTARD

WILD MORNING GLORY

WILD ONION

AND OTHER BROADLEAFED WEEDS

The plant food supplies the quick pick-up needed to fill out bare spots left by dead weeds and helps prevent re-establishment of weeds by encouraging growth and formation of thick, weed resistant turf.

HOW TO USE

Apply only in warm weather when soil moisture is plentiful and weeds are actively growing. Weeds are hard to kill when weather is cold in the early spring or during hot, dry summer weather. For best results, apply while lawn is wet with dew or after sprinkling lightly. Apply several days after mowing to allow for the presence of plenty of leaf surface to catch and retain Turf Tonic Weed and Feed, then do not mow or water lawn for another day.

Apply Turf Tonic Weed and Feed at the rate of 40 lbs. to 4,000 sq. ft. (40 ft. x 100 ft.) using a conventional spreader adjusted to a low setting. For uniform coverage, several passes in opposite directions may be necessary to empty the hopper over the measured area. Avoid application to creeping grasses such as bent, or to newly seeded turf areas until grasses have become well established. Do not apply closer than three feet from ornamental plantings such as trees, shrubs, and flowers.

Suggested Spreader Settings

Spread 2 directions for even coverage at these settings. Setting may vary with age, model and condition of spreader.

Spreader

Approx. Setting

SCOTTS (Model 35-8)	6
ORTHO (Model 3000)	6
LAWN BEAUTY (16", 18", or 24")	7
WARDS GARDEN MARK (Model AX-1093A)	7
SEARS CRAFTSMAN (Model 452-19183 — 18")	5 1/4
(Model 452-19194 — 19")	5 1/4
(Model 452-19203 — 20")	5 1/4

The plant food supplies the quick pick-up needed to fill out bare spots left by dead weeds and helps prevent re-establishment of weeds by encouraging growth and formation of thick, weed resistant turf.

HOW TO USE

Apply only in warm weather when soil moisture is plentiful and weeds are actively growing. Weeds are hard to kill when weather is cold in the early spring or during hot, dry summer weather. For best results, apply while lawn is wet with dew or after sprinkling lightly. Apply several days after mowing to allow for the presence of plenty of leaf surface to catch and retain Turf Tonic Weed and Feed, then do not mow or water lawn for another day.

Apply Turf Tonic Weed and Feed at the rate of 40 lbs. to 4,000 sq. ft. (40 ft. x 100 ft.) using a conventional spreader adjusted to a low setting. For uniform coverage, several passes in opposite directions may be necessary to empty the hopper over the measured area. Avoid application to creeping grasses such as bent, or to newly seeded turf areas until grasses have become well established. Do not apply closer than three feet from ornamental plantings such as trees, shrubs, and flowers.

Suggested Spreader Settings

Spread 2 directions for even coverage at these settings. Setting may vary with age, model and condition of spreader.

Spreader	Approx. Setting
SCOTTS (Model 35-8)	6
ORTHO (Model 3000)	6
LAWN BEAUTY (16", 18", or 24")	7
WARDS GARDEN MARK (Model AX-1093A)	7 1/4
SEARS CRAFTSMAN (Model 452-19183 — 18")	5 1/4
(Model 452-19194 — 19")	5 1/4
(Model 452-19203 — 20")	5 1/4

CAUTION:

Harmful if swallowed. Do not take internally. Avoid contact with eyes. Avoid contact with skin and clothing. Avoid inhalation of dust. Keep out of reach of children. Avoid drift to or contact with vegetables, flowers, shrubs, ornamentals, or other plants that are sensitive to 2,4-D. Do not store near growing plants or fertilizers, seeds, insecticides, and fungicides. Do not contaminate irrigation ditches or water used for domestic purposes. Not recommended for use on bent grass, or on lawns where clover is present and desired. Also not recommended for use on dichondra, centipede grass, and St. Augustine grass. Do not apply to newly seeded lawns. Clean spreader after each use. Keep bag tightly closed. Do not re-use container. Destroy when empty. Do not burn.

Notice: Buyer assumes all risk in use, storage and handling of this product, when not in strict accordance with directions given herewith.

WHERE TO USE

FORAGE UNLIMITED OLV 4D-2T is used to control woody plants in fence rows, vacant lots, roadsides, rangeland and pastures.

WOODY PLANTS CONTROLLED

Aspen
Birch
Blackberry
Box Elder
Buck Brush
Butternut
Crab-Apple
Daw-Berry
Dog Wood
Elder-Berry
 Elm
Ground-Cherry
Honey-suckle
Kudzu-Vine
Oak
Orange-Orange
Poison-Ivy
Raisin-Oak
Raspberr
Sagebrush
Spice-Brush
Sumac
Sweet-Gum
Virginia-Creeper
Wild-Cherry
Wild-Grass
Wild-Rice
Willow
(And many other species)

DIRECTIONS FOR USE

Add the recommended amount of product to diesel oil, No. 1 or No. 2 fuel oil, or kerosene and mix thoroughly. This product is not emulsifiable. DO NOT DILUTE WITH WATER.

RANGELAND AND PASTURES: Use one-half (1/2) to one and one-third (1 1/3) pint in sufficient oil (one to two gallons) to give good coverage to one acre. Use only on established stands of perennial grasses.

BASAL TREATMENT: Dilute one (1) gallon of product with twenty (20) gallons of oil (one quart in five gallons). Wet the lower 18 inches of all stems until material runs down on all sides and collects around the root collar at ground line. May be applied in any season.

STUMP TREATMENT: Dilute one (1) gallon of product with twenty (20) gallons of oil (one quart in five gallons). Spray the entire stump, particularly bark and exposed roots, at any time during the year but always as soon as possible after the trees are cut. A thorough drenching is essential for complete kill.

FRILL TREATMENT: Dilute one (1) gallon of product with twenty (20) gallons of oil (one quart in five gallons). Make frill by using an axe to cut overlapping notches in a continuous ring around the trunk near its base. Cut through the bark but do not remove chips. This method is recommended for trees 5 to 6 inches in diameter and larger. Treat freshly cut frills at any time during the year. Pour in as much of this mixture as the frill will hold without wasting chemical.

ACTIVE INGREDIENTS:	
2-Ethylhexyl Ester+ of 2,4-Dichlorophenoxyacetic Acid	62.8%
2-Ethylhexyl Ester+ of 2,4,5-Trichlorophenoxyacetic Acid	20.0%
INERT INGREDIENTS:	7.2%
TOTAL	100.0%
+ An Isooctyl Ester	
+ An Isooctyl Ester	
(Equivalent to 41.7% Dichlorophenoxyacetic Acid and 20.0% Trichlorophenoxyacetic Acid.)	

EPA Registration No. 33330-1

FORAGE UNLIMITED

RANGELAND BRUSH SPRAY OLV 4D-2T

LOW VOLATILE HERBICIDE

2,4-D and 2,4,5-T

4 lbs. of 2, 4-D and 2 lbs. of 2,4,5-T acid equivalents per gallon

Sold By

FORAGE UNLIMITED

(214) 758-6663
P. O. Box 7346
Longview, Texas USA

CAUTION

This product may cause skin irritation. Avoid inhaling spray mist. Harmful if swallowed. Avoid contact with the eyes, skin or clothing. Flush eyes with clear water and get prompt medical attention.

DO NOT apply this product to susceptible broad-leaved plants that are not desirable such as cotton, soybeans, tomatoes, grapes, clover or ornamentals. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

Minute quantities may cause severe injury by drift or vaporization. Avoid drift of spray and use a coarse spray which is less likely to drift. Although this product contains an active ingredient of low volatility, use it with caution where the spreading of vapors can damage nearby susceptible plants.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DO NOT apply this product on lawns. Legumes, such as white clover, are susceptible and may be severely damaged or killed.

DO NOT graze meat animals on treated areas within two weeks of slaughter.

DO NOT graze dairy animals on treated areas within six weeks after application.

DO NOT store this product near fertilizers, seeds, insecticides, or fungicides.

DO NOT contaminate feed or feedstuffs.

DO NOT use around home, or on recreation sites, or similar areas.

DO NOT use spray equipment contaminated with this product for any other purpose unless the equipment has been cleaned very carefully with a suitable chemical cleaner.

DO NOT mix with liquid fertilizers.

If stored below freezing, it may be necessary to warm to 40 degree F and agitate before using. This does not affect the efficiency of the product.

DO NOT reuse empty container. Destroy by burying in a safe place away from water supplies. Do not burn.

NONWARRANTY

FORAGE UNLIMITED warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purpose. Since time, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

NET CONTENTS: 5 GAL

CAUTION KEEP OUT OF REACH OF CHILDREN