

WHERE TO USE

FORAGE UNLIMITED A-6D is used to control broad-leaved weeds in corn, sorghum, cotton, soybeans, and other crops. It is also used to control weeds in pastures, lawns, and other areas. It is not for use on aquatic weeds.

PLANTS CONTROLLED

FORAGE UNLIMITED A-6D will kill or control the following as well as many other noxious plants susceptible to 2,4-D.

- | | | |
|---------------------------------------|----------------|------------------|
| Arrowhead | Goldenrod | Puncture vine |
| Artichoke | Ground Ivy | Purslane |
| Bindweed (hedge, field, and European) | Hemp | Rush |
| Bitter winter-cress | Henry's cross | Russian thistle |
| Box elder | Honey-suckle | Sagebrush |
| Beckham | Indigo | Shepherd's-purse |
| Buff thistle | Ironweed | Smartweed |
| Bursh | Jimson weed | Sow thistle |
| Burdock | Lambs-quarters | Stinkweed |
| Bur ragweed | Lotusweed | Sumac |
| Buttercup | Mexican weed | Sunflower |
| Canada thistle | Morning-glory | Virginia creeper |
| Catnip | Mustard | Water hyacinth |
| Chickweed | Nutgrass | Water lily |
| Chickory | Parrot feather | Water primrose |
| Cocklebur | Pennywort | Wild garlic |
| Coffee bean | Pigweed | Wild lettuce |
| Crooking Jenny | Plantain | Wild onion |
| Curly indigo | Poison ivy | Wild radish |
| Elderberry | Pokeweed | Willow |
| | Povertyweed | Witchweed |

DIRECTIONS FOR USE

Mix the recommended amount of product with sufficient water to obtain good spray coverage. Do not mix this product with oil. Best results are obtained when applications are made while the weeds are young and actively growing.

On Fall-Planted Wheat, Oats, and Barley—Use 2/3 to 1 pint of product in 5 to 10 gallons of water to cover one acre. Apply in early spring when weeds are small and before the crop has reached the boot stage.

On Spring-Planted Wheat, Oats, and Barley—Use 2/3 pint in 5 to 10 gallons of water to cover one acre. Apply after the fully tillered stage except during the boot to dough stage.

DO NOT forage or graze treated grain fields within two weeks after treatment with 2,4-D. DO NOT feed treated straw to livestock.

On Corn and Sorghum—Use 2/3 pint in 5 to 10 gallons of water to cover one acre when weeds are in active growth. Local climatic conditions determine when treatment should be made. Best results are usually obtained when weeds are small and corn is 4 to 18 inches tall. Do not apply from tasseling to dough stage. Treat sorghum when 4 to 12 inches tall. Use drop nozzles when either crop is over 10 inches tall.

On Rice—Use 2/3 to 1 quart in 5 to 10 gallons of water to cover one acre when weeds are in active growth stage. Rice plants are sensitive to 2,4-D in early stages of growth; therefore, it is advisable to delay spraying until the second or third week after flooding. Water in the field should be shallow enough to permit direct application of the spray material to the weeds. Make all treatments well in advance of heading.

For Aquatic Weeds in lakes, ponds, and drainage ditches.—Use 1/2 to 3 pints of product in 50 to 100 gallons of water per acre. Apply in enough water to wet foliage thoroughly. Application should be made when leaves are fully developed above water line and plants are actively growing.

DO NOT apply to more than 1/3 to 1/2 of a lake or pond in any one month, because excessive decaying vegetation may deplete oxygen content of water, killing fish.

On Pastures, Ornamental Turf, and Noncrop Areas—Use 3/3 to 1-1/3 quarts and enough water to give sufficient coverage to one acre on established stands of perennial grasses, depending on type of weeds and stage of growth. **DO NOT** use on Dichondra and on creeping grasses such as bent except for spot spraying. Newly seeded turf should not be treated until after the second mowing and at the lower dosage rate.

DO NOT graze dairy animals on treated areas within seven days after application.

Perennial and Other Hard-to-Control Weeds may require a repeat application to give adequate control.

FORAGE UNLIMITED

BROADLEAF WEED SPRAY

A-6D

2,4-D AMINE

6 LBS. OF 2,4-D ACID EQUIVALENT PER GALLON

ACTIVE INGREDIENTS:
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid 69.1%

INERT INGREDIENTS: 30.9%

TOTAL 100.0%

(Equivalent to 57.4% of 2,4-Dichlorophenoxyacetic Acid)

EPA REGISTRATION NO. 33330-3

SOLD BY
FORAGE UNLIMITED
(214) 758-6663
P. O. BOX 7346
LONGVIEW, TEXAS, USA

ACCEPTED

JUL 10 1974

UNDER THE FEDERAL INSECTICIDE AND FUNGICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 33330-3 SUBJECT TO ATTACHED COMMENTS.

WARNING

DO NOT apply this product to susceptible broad-leaved plants that are desirable such as cotton, soybeans, and other crops. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth. Minute quantities may cause severe injury by drift. Avoid drift of spray and use a coarse spray which is less likely to drift.

DO NOT contaminate water in irrigation ditches or water used for domestic purposes.

Treatment of aquatic weeds can result in oxygen loss from decomposition of dead weeds. This loss can cause fish kills. Therefore, to minimize this hazard, treat 1/3 to 1/2 of the water area in a single operation and wait at least 10-14 days between treatments. Begin treatment along the shore and proceed outward in bands to allow fish to move into untreated areas.

Users should note that herbicide treatment of public waters requires a permit from appropriate state agencies in most states. Your State Conservation Department or Game and Fish Commission will aid you in securing a permit if required in your state.

DO NOT store this product near fertilizers, seeds, insecticides, or fungicides. **DO NOT** contaminate feed or foodstuffs.

DO NOT use spray equipment contaminated with this product for any other purpose. **DO NOT** apply insecticides, fungicides, or other agricultural products unless the equipment has been cleaned very carefully with a suitable chemical cleaner.

If stored below freezing, it may be necessary to warm to 40 F and agitate before using. This does not affect the effectiveness of the product.

DO NOT mix with liquid fertilizers.

DO NOT reuse empty containers. Destroy by burying in an isolated place away from water supplies. **DO NOT** burn.

CAUTION

This product may cause skin irritation. Avoid inhaling spray mist. Harmful if swallowed. Avoid contact with the eyes, skin, or clothing. Flush eyes with clear water and get prompt medical attention.

NONWARRANTY

FORAGE UNLIMITED warrants that this herbicide conforms to the chemical description on its label. Other warranties are made in accordance with the directions under normal conditions. This herbicide is not warranted for any other purpose. Since the effectiveness of application, weather, plant and soil conditions, and other factors may vary, results may be beyond our control, no warranty is given concerning the use of this product contrary to the label directions or other conditions which are abnormal or not reasonably foreseeable. The user assumes the responsibility of any such use.

NET CONTENTS: 5 GAL.

CAUTION: KEEP OUT OF REACH OF CHILDREN

SEE PANELS FOR ADDITIONAL INFORMATION