BEST AVAILABLE COPY

AQUA KLEAR

HAR 2 - 1987

ACTIVE INGREDIENT:

Poly(oxyethylene (dimethyliminio) ethylene-(dimethyliminio) ethylene dichloride)

__15.0__%

INERT INGREDIENT

85.0 **%**

This product contains $\frac{1.29}{8.58}$ lb. of active ingredient per gallon and weighs $\frac{8.58}{100}$ lb. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water.

Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is tooc to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds, or public water. For gordance, contact the regional office of the Environmental Protection Agency.

DIRECTIONS FOR

and fungi in recirculating commercial and induto its use, systems must be cleaned to remoslime, and other deposits. An initial slug additional fluid ounces of AQUA KLEAR of water to provide a concentration of 32 AQUA KLEAR, but the system, is recommended. Repeat initial

Subsequent slug additions of 1.0

AQUA KLEAR per 1000

_____ parts per million of ______ AQUA_KI,EAR employed every 2 to 5 days, or as needed. The upon the relative amount of bleedoff and the selem. Slug additions should be made in the surplus AQUA_KLEAR ______ is use

air-washing systems that maintain effective m to its use, systems should be cleaned to redeposits. An initial slug dose of 14,8

mended. Repeat initial dosage until control is tions of __10,1__ to __24.7_ fluid ounces of _ per 1000 gallons of water should be employed e frequency of addition depends upon the relativof the bacterial problem. Slug additions may be

collection trays of the airwash system.

STORAGE AND D

Do not contaminate water, food, or feed by storage or disposal STORAGE. Do not stack more than four drums high. Leaking drums for disposal. Spills should be absorbed in sawdust or sand

tainer closed when not in use
PESTICIDE DISPOSAL. Wastes resulting from the use of the
proved waste disposal facility.

CONTAINER DISPOSAL

- © PLASTIC Trule rinse for equivalent). Then offer for recycling sandary fundfill, by incineration or, if allowed by state and to soupe
 - "I METAL Triple rosse for equivalents. Then offer for recycling or samilary landful, or by other procedures approved by state an

	Manufac	ture
	QUALITY CHEMICIA CO.	
	1835 NE 144 ST.	
•	N. MIAMI, FL 33181	
EPA RE	3. NO	E

NET CONTENTS _