

[33404-3]

DIRECTIONS FOR USE INDOOR APPLICATION

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Vacate treated area and ventilate before reoccupying.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, controled flour beetles, rust red flour beetles, swa-toothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, meal worm, grain mites, and cad-elles: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed. *Will indoors*

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings, and floors. Repeat treatment as needed. *Will indoors*

To control fleas and brown dog ticks: Thoroughly spray infested areas. pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, mouldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed. *Will indoors*

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus, and other storages, directing the spray into cracks, joints, and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

site: mothproofing

VIP INSECTICIDE

EPA Reg. No. 33404-3

A Concentrated PYRENONE Micro Emulsion
-Clear

-Stable

-Leaves No Oily or Objectionable Residue

pests
Effective, Fast Acting Insecticide.

For Use in Homes, Institutions, Restaurants.

Food Processing Plants, Dairies, Livestock and Outdoors *Will*
for Mosquito Control.

Apply with Conventional Mechanical or Compressed Air Equipment.

ACTIVE INGREDIENTS:

Pyrethrins	0.1%
*Piperonyl Butoxide, technical	1.0%
Petroleum distillate	0.4%

INERT INGREDIENTS: 98.5%

*Equivalent to 0.8% (butylcarbityl) (6 propylpiperonyl) ether and to 0.2% related compounds.

Pyrenone is a registered trademark of FMC Corporation


CAUTION

Keep Out Of Reach Of Children

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

Net Contents Gallon

MANUFACTURED IN U.S.A. BY



STAT ENTERPRISES INC.

SPACE SPRAYING: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths: direct spray at an upward angle, distributing it uniformly throughout the entire area, at a rate of 3-6 ounces per 1000 cubic feet of space.

FOGGING: To kill exposed or accessible stages of the above named crawling insects, use a rate of one gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

ON ANIMALS: To protect cattle and horses from horse flies, house flies, mosquitoes, and gnats, apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse flies, and deer flies, apply at a rate of 2 ounces per adult cow or horse, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control blood-sucking lice, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 or 3 weeks if required.

OUTDOOR ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest around low marshy places, stagnant pools or ponds and shore lines. Apply at a rate of 1 1/2 gallons per acre (4 1/2 ounces per 1000 square feet).

Using truck mounted conventional misting equipment such as Rotomist models 51C, 91 or 100 HT, traveling at 5 m.p.h. to treat a 300 foot swath, calibrate equipment to deliver 4.5 gallons per minute.

CAUTION: Do not contaminate water.

CAUTION: Harmful if swallowed. Avoid contamination of feed and foodstuffs.

Remove pets, birds and cover fish aquariums before spraying. This product is toxic to fish. Keep out of lakes, streams or ponds.

Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

ACCEPT
25 SEP 19
SEP 19 198
U.S. DEPT. OF AGRICULTURE
FEDERAL AGENCY FOR
FOOD & DRUG ADMINISTRATION
ED. BRADY JR.
TO ATTACH COMB

33404