

Precautionary Statements

Hazards to Humans and domestic animals

**DANGER**

Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

**STORAGE AND DISPOSAL**

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

—OPEN DUMPING IS PROHIBITED.

—DO NOT REUSE EMPTY CONTAINER.

**PESTICIDE DISPOSAL**

PESTICIDE, SPRAY MIXTURE, OR RINSATE THAT CAN NOT BE USED OR CHEMICALLY PROCESSED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

**CONTAINER DISPOSAL**

RESEAL CONTAINER & OFFER RECONDITIONING, OR TRIPLE RINSE (OR EQUIVALENT) AND: OFFER RECYCLING, RECONDITIONING, OR DISPOSAL IN APPROVED LANDFILL, OR BURY IN A SAFE PLACE

**GENERAL**

CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATE PROCEDURES.

Directions for Use  
**GENERAL CLASSIFICATION**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply Bactosep with a cloth mop or mechanical spray device. When applied with a mechanical spray device, surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 30 minutes. Fresh solution should be prepared daily or when the use solution becomes visibly dirty.

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions

For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard, non-porous surfaces.

Add 3 1/2 ounces BACTO-SEP to 4 gallons water. Apply to previously cleaned hard surface with mop or cloth.

At this use-level, BACTO-SEP is effective against *Pseudomonas aeruginosa*.

Disinfection in Institutions, Industry and Schools

For disinfecting floors, walls, bedframes, countertops, tables, chairs, garbage pails, bathroom fixtures and other hard, non-porous surfaces.

Add 2 oz of BACTO-SEP to 4 gallons of water. Apply to previously cleaned hard surface with mop or cloth.

At 2 oz/4 gallon use-level, BACTO-SEP is effective against *Staphylococcus aureus*, *Salmonella choleraesuis* and *Trichophyton interdigitale* (the athlete's foot fungus).

Disinfection of Barber Tools

Precleaned barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a 1% solution of BACTO-SEP.

**BACTO-SEP**

Disinfectant-Sanitizer-Fungicide  
Deodorizer for Hospital, Institutional,  
Industrial, School, Dairy and Other  
Farm Use

Active Ingredient

Didecyl dimethyl ammonium chloride

Inert Ingredients

92.5%

100%

KEEP OUT OF REACH OF CHILDREN

**DANGER**

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Net Contents

EPA Reg. No. 33  
EPA Est. No. 33

38

Manufactured by

**Interchem inc.**

3516 N. 13th Street, P.O. Box 13101

**ONLY FOR SP TO USE AND STORAGE BY SERVICE PERSONS.**