

[33543-1]

Net Weight  
1 Pound, 6 Oz

Kills House Flies,  
Horn Flies, Stable  
Flies, Mosquitoes  
and Gnats.

**CONTENTS**

**ACTIVE INGREDIENTS**

Pyrethrins I & II	25.0%
Technical Piperonyl Butoxide*	50.0%
N-Octyl Cycloheptene dicarboximide	8.0%
Refined Petroleum Oil	12.40%

**INERT INGREDIENTS**

Propellant Gas 80.000%  
\*Equivalent to 40% of n-butyl-  
carbitol, 16-propyl piperonyl ether,  
and 40% of related com-  
pounds.

**CAUTION:** Keep from reach of  
children. Do not store above 120°  
F. or incline it. Do not spray di-  
rectly on shirt, fish, food, open  
flame, or on salt surfaces.

In food processing, foods should be  
removed or covered during treat-  
ment, and processing surfaces cov-  
ered or cleaned before using. Use  
as a space spray only when plant  
is not in operation.

**NEW  
AEROSOL**

**EMBEECIDE**

**FLY & INSECT BOMB**

EMBEECIDE is Recommended for use in:

- Restaurants
- Bakeries
- Packing Houses
- Dairies
- Ice Cream Plants
- Motels
- Locker Plants
- Candy Plants
- Storage Plants
- Institutions
- Hotels
- Hospitals
- Food Processors
- Bottlers

**CAUTION:** KEEP OUT OF REACH OF  
CHILDREN. CONTAINS FLAMMABLE LIQUID.  
SEE PANEL FOR OTHER CAUTIONS.

**DIRECTIONS**

**For Milk Handling & Food  
Processing Areas**

For flies, mosquitoes, gnats, stable flies,  
wasps, flying moths. Obtain cubic feet of  
space to be treated. Example, 10' x 12'  
x 10' equal 1200 cubic feet, or approxi-  
mately 7-second application. Seconds  
may be accurately determined by count-  
ing "one-thousand, two-thousand, three-  
thousand, etc."

Close all doors and windows and dis-  
perse spray upward in all directions  
from five to six seconds per 1000 cubic  
feet. Keep areas closed for 15 minutes  
after treatment.

**FOR HOME USE**

To kill roaches, spiders, silverfish  
and certain other crawling insects,  
thoroughly spray insects and all  
possible hiding places, such as  
cracks, base boards, sinks, cabinets  
and shelves. Repeat treatment when  
necessary.

**HORN FLIES**

For horn flies: Spray the adult cows  
for several seconds on each side,  
being careful to spray back, withers  
and forelegs thoroughly. Hold back  
about 10 feet away from animal  
when spraying. Spray daily, or as  
necessary.

U.S.D.A. Reg. No. 5774-1

Formulated For

