

REVISED
DRAFT LABEL

10FS
P-23
33560-41

PRONONE® POWER PELLET
HERBICIDE

ACTIVE INGREDIENT:	Hexazinone	
	3-cyclohexyl-6-(dimethylamino)-1-methyl-1,3,5-triazine-2,4-(1H,3H)-dione	75%
INERT INGREDIENTS:		25%
TOTAL		100%

EPA Est. No. 33560-TN-1

EPA Reg. No. 33560

(Container Size)

ACCEPTED
MAR 9 1957

KEEP OUT OF REACH OF CHILDREN

WARNING

STATEMENT OF PRACTICAL TREATMENT

REGISTERED UNDER
EPA Reg. No. 33560-41

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air and apply respiration if indicated.

IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly.

IF IN EYES: Rinse eyes with plenty of water. Call a physician.


PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Causes eye irritation. Do not get in eyes, on skin, or on clothing. May cause irritation of nose, throat or skin. Wash clothing and equipment after application. Do not store near food or feed products. Store out of the weather.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water or wetlands. Do not contaminate water used by wildlife and aquatic life or for domestic and irrigation purposes. Do not contaminate water by cleaning of equipment or disposal of waste.



IMPORTANT

Injury to or loss of desirable trees or other plants may result from failure to observe the following. Do not apply or empty equipment near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. DO NOT use on lawns, walks, driveways, tennis courts or similar areas. Prevent contact with desirable plants. Keep from contact with fertilizers, insecticides, fungicides and seeds.

GENERAL INFORMATION

Pro-Serve PRONONE® Power Pellet Herbicide is a concentrated pellet of Hexazinone allowing easy and uniform dosage as an individual stem treatment for control of undesirable woody plants such as elm, hackberry, huisache, mesquite, myrtle, oaks, osageorange, red cedar, tallow, whitebrush and willow in rangeland and for use in forestry site preparation, preharvest control of hardwoods and pine release in pine-hardwood stands for growth enhancement of loblolly, slash, shortleaf and longleaf pine. The treatment provides control of many woody plants such as black cherry, blackgum*, dogwood*, elm, hawthorn, hickory*, oak, red maple*, sweetgum and sumac*. On woody plants, symptoms usually appear within 3 weeks after sufficient rainfall has dissolved the pellet and carried the herbicide into the root zone where it is absorbed during periods of active plant growth. Defoliation and subsequent refoliation may occur, but susceptible plants are killed. The effect of treatment will vary with the amount applied, soil type, rainfall, timing and uniformity of application and other conditions.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Pro-Serve, Inc. will not be responsible for losses or damages resulting

*Considered hard to kill

from use of this product in any manner not specifically recommended to Pro-Serve, Inc.

User assumes all risk associated with such non-recommended use. Make application just before or during the period of active growth of brush species to be controlled. Wear gloves when handling pellets to prevent contact with skin.

Rangeland: Apply the pellet to the soil within 3' of the root collar at the rate of 1 or 2 pellets for each 1" of stem diameter of brush to be controlled. When treating large stems and when more than one pellet is needed per stem, make application on opposite sides of the stem. Do not treat more than 300 one-inch stem equivalents (approximately 70 brush plants) per acre per season. Late winter to mid-spring application is suggested. Do not use on marshy or poorly drained sites or on soils classified as clays. Do not cut hay from treated grasses. Do not use on pastures.

NOTE: For effective brush control and prevention of damage to desirable vegetation do not apply to brush standing in water; do not use water from treated ditches for irrigation; do not use where desirable trees or shrubs are immediately adjacent to treated area.

Forestry: Apply at a rate of 1,892 to 5,676 pellets per acre for preharvest and site preparation, and 946 to 3,784 pellets per acre for pine release. Make the application using a grid pattern as suggested in the table below. Do not apply Power Pellets closer than 36" to seedlings or directly upslope from seedlings.

Selection of the rate per acre and grid pattern will depend on soil texture and woody plant species composition. Use the higher rates on fine-textured soils and when the major component of the hardwoods is a difficult-to-control species such as blackgum, dogwood, hickory and red maple. Use the lower rates on medium to coarse texture soils where elm, cherry, oak and sweetgum are dominant.




TABLE OF SUGGESTED APPLICATION PATTERNS
FOR PRONONE® POWER PELLETS

For Site Preparation and Preharvest Control for Hardwoods

<u>Soil</u>		<u>Grid Spacing (ft)</u>
Coarse	1 Pellet	5'x5' or 4'x5' or 4'x4'
Medium/Fine	1 Pellet	3'x4' or 3'x3' or 2½'x3'

For Pine Release

<u>Soil</u>		<u>Grid Spacing (ft)</u>
Coarse	1 Pellet	6'x7'* or 5'x6'
Medium/Fine	1 Pellet	4'x6'

*Use on deep sands with pines four years of age.

NOTE: As spacing decreases, rate increases.


PRONONE® Power Pellet should be applied during the period from late winter (bud-break) to early summer. DO NOT BURN treated sites after initial foliage browning is evident. Burn only after brush has completely defoliated, at least once, from sufficient root uptake of the herbicide. Inconsistent results may occur if the site is burned 3 to 6 months prior to treatment.

NOTE: Use on 1-year old or 4-year old or older seedlings is acceptable; however, injury may result from use on 2- and 3-year seedlings where root growth is extensive but hardiness is lacking.

Results may be unsatisfactory where stump sprouts of less than one year's growth predominates.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.



50F5

STORAGE: Store product in original container only, away from other pesticides, fertilizer, food or feed.

CONTAINER DISPOSAL: Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

NOTICE OF WARRANTY

Pro-Serve warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risk inherently associated with use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner or use or application, all of which are beyond the control of Pro-Serve, Inc. In no case, shall Pro-Serve be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by buyer. Pro-Serve **MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS** for a particular purpose nor any other express or implied warranty except as stated above.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

