

PLEASE NOTE

**This image contains more than one label
approved for this product on this date.**

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

APR 13 2010

Ms. Peggy Galloway
Landis International
P.O. Box 5126
Valdosta, GA 31603-5126

Authorized agent for Mitsui Chemicals

Dear Ms. Galloway:

Subject: Label amendment; add greenhouse grown tomato use
Dinotefuran 20% Turf, Ornamental and Vegetable Transplant
EPA Registration Number 33657-16
Your submission dated July 24, 2009

The amendment referred to above is conditionally acceptable in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) section 3(c)(7)(B), subject to the comments listed below:

1. Registration does not eliminate the need for continual reassessment of pesticides. If the Agency determines that, at any time, additional data are required to maintain in effect an existing registration, the Agency will require submission of such data.
2. We have approved this label on the condition that you will submit a data matrix and a Certification with Respect to Citation of Data form within 4 weeks from the date of this letter.
3. Please make the following label changes:
 - a. Provide the new registration number when the transfer of this product from MITSUI CHEMICALS, INC. (Company No. 33657) to MITSUI CHEMICALS AGRO, INC. (Company No. 86203) is complete, as indicated by you in a previous e-mail dated 4/6/2010 regarding the transfer of all Dinotefuran registrations.

- b. Provide an American address on the label.
- c. On page 9, revise "Important Notes" to read "Restrictions".
- d. On page 9, revise "For optimal control, apply irrigation if rainfall does not...." to read, "Irrigate turf if rainfall does not.....".
- e. Delete forestry, Christmas trees, tree plantations, reforestation nurseries, forests and all use directions related to these sites from pages 1, 10, 11, 12, 15, 19, 20, and elsewhere on the label.
- f. Make the following changes in the Greenhouse Grown Tomatoes use directions:
 - Delete "enclosed structures" from this site. This use was assessed for greenhouse grown tomatoes only, not for any other enclosed structures.
 - In the remarks column, add the heading "Use Restrictions" after the gallonage text, and before all the "Do not" statements in both foliar and soil sections.
 - Add "Do not use adjuvants or surfactants." under the restrictions for foliar and soil sections. Submitted field trial data indicates that an adjuvant or surfactant was not used in these residue studies.
 - In the soil section add the gallonage text indicated below:

 "At 0.75 lb/ 100 gallons, 360 gallons of drench solution will deliver 2.7 lbs of Dinotefuran 20% Turf and Ornamental per acre. At 1.5 lb/100 gallons, 180 gallons of drench solution will deliver 2.7 lbs. of Dinotefuran 20% Turf & Ornamental per acre."
 - Provide application rate in lbs a.i. per 100 gallons for soil drench.
 - There are two grammatical errors on page 13. Add "that" to the fourth restriction under soil drench: "...plants *that* have received..."; and add "foliar" in the third sentence of general directions below the use rates: "...more than two *foliar* sprays.....".
- g. Delete the use directions for Tree Injection from page 20. This is a new use for dinotefuran, and must be submitted as a separate PRIA action.
- h. Add the following revised container disposal statement from your notification

dated Feb 10, 2010:

- Container Handling and Disposal


Nonrefillable container: Do not reuse or refill this container.

Triple rinse as follows. Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or reconditioning if appropriate or puncture and dispose off ina sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Submit two (2) copies of your final printed labeling before you release the product for shipment. Also submit a data matrix and a Certification with Respect to Citation of Data form within 4 weeks from the date of this letter. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(a).

A stamped copy of the label is enclosed for your records. A copy of the review of submitted residue data is also enclosed for your perusal. If you have any questions about this letter, please contact Rita Kumar at (703) 308-8291 or kumar.rita@epa.gov.

Sincerely,



Kable Bo Davis
Acting Product Manager 7
Insecticide Rodenticide Branch
Registration Division (7505P)

Enclosures (2)

4
25

GROUP	4A	INSECTICIDE
-------	----	-------------

DINOTEFURAN 20 %

TURF, ORNAMENTAL, FORESTRY AND VEGETABLE TRANSPLANTS IN ENCLOSED STRUCTURES

For foliar and systemic insect control in ornamental plants, vegetable transplants in enclosed structures, and turfgrass.

For Greenhouse, Nursery, Interior Plantscape, Outdoor Landscape, Forestry and Greenhouse Tomato Use Only
Active Ingredient:

Dinotefuran, [N-methyl-N'-nitro-N''-(tetrahydro-3-furanyl)methyl]guanidine)	20%
Other Ingredients	80%
Total:	100%

KEEP OUT OF REACH OF CHILDREN
CAUTION
See side panels for additional precautionary statements

FIRST AID	
If On Skin Or Clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.
If Swallowed	<ul style="list-style-type: none"> Call poison control center or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by the poison control center or doctor. Have person sip a glass of water if able to swallow. Do not give anything by mouth to an unconscious person.
If In Eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.
If Inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact CHEMTREC (800) 424-9300 (24 hours) for emergency medical treatment information.</p> <p><i>See side/back panels for additional precautionary statements</i></p>	

EPA Reg No. 33657-16

Net Contents:
Manufactured By:
MITSUI CHEMICALS, INC.
Shiodome City Center
1-5-2 Higashi-Shimbashi
Minato-ku, Tokyo 105-7117
JAPAN

EPA Establishment No. 67545-AZ-01

ACCEPTED
With COMMENTS
In EPA Letter Dated:
APR 13 2010
**Under the Federal Insecticide, Fungicide
and Rodenticide Act, As amended, for the
pesticide Registered under EPA Reg. No:**
33657-16

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Do not allow contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to shrimp. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not dispose equipment washwaters or rinsate into a natural drain or water body.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Dinotefuran and its degradate, MNG have the properties and characteristics associated with chemicals detected in ground water. The high water solubility of dinotefuran, and its degradate, MNG, coupled with its very high mobility, and resistance to biodegradation indicates that this compound has a strong potential to leach to the subsurface under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination. Periodic monitoring of shallow groundwater in the use area is recommended.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

SPRAY DRIFT ADVISORY

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crop thereof rendered for sale, use or consumption.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL, USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural insecticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

EXCEPTION: If product is drenched or soil-injected, workers may enter the area at any time if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. **Do not allow others to enter treated areas until sprays have dried.**

APPLICATION INFORMATION

- Applications of Dinotefuran 20% Turf and Ornamental in residential areas may be made by commercially licensed applicators.

Application to Turfgrass:

- Dinotefuran 20% Turf and Ornamental can be used for the control of soil inhabiting pests of turfgrass such as Masked Chafers, European Chafer, Green June Beetle, May or June Beetle, Japanese Beetle, Oriental Beetle, Billbugs, Annual Bluegrass Weevil, Black Turfgrass Ataenius and Mole Crickets. Dinotefuran 20% Turf and Ornamental can also be used for the suppression of cutworms and chinchbugs in turfgrass areas.
- Dinotefuran 20% Turf and Ornamental can be used as directed on outdoor residential, recreational and commercial turfgrass in sites such as home lawns, commercial lawns, multi-family residential and apartment complexes, grounds or lawns around business and office complexes, shopping centers, airports, military and other institutions, cemeteries, golf courses, playgrounds, parks, athletic fields and sod farms.

- Target timing of Dinotefuran 20% Turf and Ornamental applications at or just prior to or during egg laying of the target pests. The need for an application can be based on historical and/or physical monitoring of the site, current season adult trapping, previous experience or other methods. Optimum control will be achieved when applications are made prior to or at egg hatch of the target pests followed by sufficient irrigation or rainfall to move the active ingredient through the turf thatch layer. Consult your State Extension Service for information regarding specific application timing.
- Do not apply when the target site is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist.

Application to Ornamental plants:

- Dinotefuran 20% Turf and Ornamental can be applied as a foliar spray, a broadcast spray, a soil drench, soil injection and via chemigation for insect control in ornamental plants in greenhouses, nurseries, outdoor landscapes and interior landscapes.
- Dinotefuran 20% Turf and Ornamental is a systemic product and will be taken up by the root system and translocated upward throughout the plant. When applied as a foliar spray, the product offers translaminar and locally systemic control of foliar pests.
- When applied to the soil, Dinotefuran 20% Turf and Ornamental will be translocated more quickly in herbaceous plants than in woody shrubs and trees. Speed of insect control will range from as little as one day for small herbaceous plants in containers, to several weeks in large trees in growing in the landscape.
- For outdoor and landscape ornamentals, broadcast applications cannot exceed a total of 2.7 lbs of product (0.54 lb. active ingredient) per acre per year.

MIXING INSTRUCTIONS:

Dinotefuran 20% Turf and Ornamental Alone: Add half of the required amount of water to the mix tank. With the agitator running, add the desired amount of Dinotefuran 20% Turf and Ornamental to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after Dinotefuran 20% Turf and Ornamental has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Dinotefuran 20% Turf and Ornamental + Tank Mixtures: Add half of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. In general, add tank mix partners in this order: products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables), liquid flowables, liquids, emulsifiable concentrates, and surfactants/adjuvants. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all the mixture has been applied.

NOTE: When using Dinotefuran 20% Turf and Ornamental in tank mixtures, add all products in water-soluble packaging to the tank before any other tank mix partner, including Dinotefuran 20% Turf and Ornamental. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using Dinotefuran 20% Turf and Ornamental in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label. Do not exceed label dosage rate, and follow the most restrictive label precautions and limitations. Do not mix this product with any product that prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

Compatibility

IMPORTANT - The safety of all potential tank mixes has not been tested on all crops. Before applying any tank mixture not specifically listed on this label, confirm the safety to the target crop.

Dinotefuran 20% Turf and Ornamental is compatible with most commonly used pesticides, crop oils, adjuvants, and nutritional sprays. However, since it is not possible to test all possible mixtures, pre-test to assure the physical compatibility and lack of phytotoxic effect of any proposed mixtures with Dinotefuran 20% Turf and Ornamental. To determine the physical compatibility of Dinotefuran 20% Turf and Ornamental with other products, use a jar test, as described below:

Using a quart jar, add the proportionate amounts of the products to 1 quart of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for additional required ingredients to the spray tank.

RESISTANCE MANAGEMENT

Dinotefuran 20% Turf and Ornamental contains a Group 4A insecticide. Insect biotypes with acquired resistance to Group 4A may eventually dominate the insect population if Group 4A insecticides are used repeatedly in the same crop or in successive years as the primary method of control for a targeted species. This may result in partial or total loss of control of those species by Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides.

To delay the development of insecticide resistance in greenhouse, nursery and interiorscape use sites, strongly consider the following guidelines:

- Do not apply Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides to consecutive generations of the same insect pest species.
- Do not drench soil media with Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides more than one time per crop cycle or three months, whichever is shorter.
- Do not make more than two foliar or broadcast sprays of Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides to a single crop during a two-month period.
- Do not make more than one soil drench and one foliar or broadcast spray with Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides during a two-month period.
- Base insecticide use on a comprehensive IPM program.
- Monitor treated insect populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturers for insecticide resistance management and/or IPM guidelines for the specific site and resistant pest problems.
- For further information or to report suspected resistance, you may contact Landis International, Inc., a representative of Mitsui Chemicals, Inc., at toll free number: 1-800-526-3471 or at their web site: www.landisintl.com

APPLICATION PROCEDURES AND SPRAY EQUIPMENT

Ground Application: Select spray nozzles that will provide accurate and uniform spray deposition. Use spray nozzles that provide medium-sized droplets and reduce drift. To help insure accuracy, calibrate sprayer before each use. For information on spray equipment and calibration, consult nozzle manufacturers and/or State Extension Service specialists.

Apply Dinotefuran 20% Turf and Ornamental using sufficient water volume to provide thorough and uniform coverage. In situations where a dense canopy exists and/or pest pressure is high, use greater water volumes. The use of a spray adjuvant may improve spray coverage. Do not apply under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Applications to turfgrass: Apply Dinotefuran 20% Turf and Ornamental through conventional spray equipment in a minimum of 1 gallon of finished spray per 1000 sq. ft. Ensure adequate distribution in the treated area using accurately calibrated equipment normally used for application of turfgrass insecticides. Use equipment that will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off target drift. Check calibration periodically to ensure that equipment is working properly. Prevent skips by using marker dyes or foam aids.

Applications to ornamental plants: Dinotefuran 20% Turf and Ornamental can be applied using many different types of application equipment. Apply in sufficient water to ensure good coverage of ornamental plants. When making applications to plants with hard to wet foliage such as holly or pine, the product may be tank mixed with a surfactant. If concentrate or mist type spray equipment is used, apply the same amount of product on the sprayed area as would be used in a dilute solution. To assure optimum effectiveness, the product must be placed where the growing portion of the target plant can absorb the active ingredient. Applications can be made to foliage or as a soil drench.

RESTRICTIONS

- Do not graze treated areas or use clippings from treated areas for feed or forage.
- Prevent runoff or puddling of irrigation water following application.
- Do not apply to areas that are water logged or saturated, or frozen, which will not allow penetration into the root zone of the plant.
-

APPLICATION THROUGH IRRIGATION SYSTEMS (CHEMIGATION):

Dinotefuran 20% Turf and Ornamental may be applied by injection into an irrigation system, either alone or in combination with other pesticides or chemicals that are registered for application through irrigation systems. Dilution ratios are normally 1:100 to 1:200, depending on the system. Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems (Turfgrass) or microirrigation (individual spaghetti tube), drip irrigation, overhead irrigation, or motorized calibrated irrigation equipment (Ornamentals). Do not apply through any other type of irrigation system. Lack of effectiveness can result from non-uniform distribution of treated water. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make adjustments when necessary.

Using Water from Public Water Systems:

DO NOT APPLY DINOTEFURAN 20% TURF & ORNAMENTAL THROUGH ANY IRRIGATION SYSTEM PHYSICALLY CONNECTED TO A PUBLIC WATER SYSTEM.

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year. Dinotefuran 20% Turf and Ornamental may be applied through irrigation systems which may be supplied by a public water system only if the water from the public water system is discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. Before beginning chemigation, always make sure that the air gap exists and that there is no blockage of the overflow of the reservoir tank.

Any irrigation system using water supplied from a public water system must also meet the following requirements:

Operating Instructions for Irrigation Systems:

1. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts.
2. The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
6. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
7. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
8. Do not apply when wind speed favors drift beyond the area intended.

Calibration and Application Instructions:

Apply Dinotefuran 20% Turf and Ornamental under the schedule specified in the specific use instructions, not according to the irrigation schedule unless the events coincide. In general, set the equipment to apply the minimum amount of water per acre. Run the system at 86 - 90% of the manufacturer's maximum rated travel speed.

The following calibration and application techniques are provided for user reference, but do not constitute a warranty of fitness for application through sprinkler irrigation equipment. Check with state and local regulatory agencies for potential use restrictions before applying any agricultural chemical through sprinkler irrigation equipment.

Center Pivot Irrigation Equipment:

NOTES: 1) Use only drive systems that provide uniform water distribution. 2) Do not use end guns when chemigating Dinotefuran 20% Turf and Ornamental through center pivot systems because of non-uniform application. 3) Plug the first nozzle closest to the wellhead to protect the water source.

1. Determine the size of the area to be treated.
2. Determine the time required to apply 0.1 - 0.25 inches of water over the area to be treated when the system and injection equipment are operated at normal pressures as recommended by the equipment manufacturer. Run the system at 80-95% of the manufacturer's rated maximum travel speed.
3. Using water, determine the injection pump output when operated at normal line pressure.
4. Determine the amount of Dinotefuran 20% Turf and Ornamental, and any tank mix partners, required to treat the area covered by the irrigation system.
5. Add the required amount of Dinotefuran 20% Turf and Ornamental, and any tank mix partners, and sufficient water to meet the injection time requirements to the solution tanks. (See "Mixing Instructions" section of this label.)
6. Make sure the system is fully charged with water before starting injection of the Dinotefuran 20% Turf and Ornamental solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
7. Inject the specified amount of Dinotefuran 20% Turf and Ornamental per acre continuously for one complete revolution of the system.
8. Stop the injection equipment after treatment is complete. Continue to operate the system until the Dinotefuran 20% Turf and Ornamental solution has cleared all of the sprinkler heads.
9. Allow time for all lines to flush the pesticide through all nozzles before turning off irrigation water.

Solid Set, Hand Move, and Moving Wheel Irrigation Equipment:

1. Determine the acreage covered by the sprinklers.
2. Fill injector solution tank with plain water and calibrate the flow rate of the system to deliver the contents of the tank over a 20-40 minute time interval.
3. Determine the amount of Dinotefuran 20% Turf and Ornamental required to treat the area covered by the

irrigation system.

- 4. Add the required amount of Dinotefuran 20% Turf and Ornamental, and any other tank mix partners, into the same quantity of water used to calibrate the injection period. (See "Mixing Instructions" section of this label.)
- 5. Operate the system at the same pressure and time interval established during the calibration.
- 6. Inject specified amount of Dinotefuran 20% Turf and Ornamental per acre for either a 20-40 minute period at the end of a regular irrigation set, or as a 20-40 minute injection as a separate application not associated with a regular irrigation to maximize retention of the insecticide by the foliage.
- 7. Stop injection equipment after treatment is completed. Continue to operate the system until the Dinotefuran 20% Turf and Ornamental solution has cleared the last sprinkler head. To ensure lines are flushed and free from remaining pesticides, a dye indicator may be injected into the lines to mark the end of the application period.

MINIMIZING SPRAY DRIFT

As with all crop protection products, it is important to minimize off-target movement. Do not allow spray to drift onto adjacent land, crops, or aquatic areas. To minimize spray drift:

- 1. Make applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 10 mph. Do not apply when wind gusts approach 10 mph.
- 2. Risk of exposure to sensitive aquatic areas can be reduced by not applying when wind direction is toward the aquatic area.
- 3. Do not cultivate or plant crops within 25 feet of the aquatic area as to allow growth of a vegetative filter strip.
- 4. Do not make applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with increased height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.
- 5. Use the largest droplet size consistent with good pest control. Small droplets are more prone to spray drift and can be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by not using excessive spray boom pressure.
- 6. Apply as close to target plants as practical to obtain a good spray pattern for adequate coverage. Don not apply more than 10 ft. above the crop canopy.
- 7. For aerial applications, moun spray boom on the aircraft so as to minimize drift caused by wing tip vortices. Use minimum practical boom length and do not use boom that exceeds 75% of wing span or rotor diameter.

Air Assisted (Air Blast) Tree and Vine Sprayers (Ornamentals Only):

Air assisted tree and vine sprayers carry droplets into the canopy of trees and vines via a radially or laterally directed air stream. In addition to the general drift management principles already described, the following specific practices will further reduce the potential for drift:

- 1. Adjust deflectors and aiming devices so that spray is only directed into the canopy.
- 2. Block off upward pointed nozzles when there is no overhanging canopy.
- 3. Use only enough air volume to penetrate the canopy and provide good coverage. Use a minimum of 50 gallons finished spray per acre.
- 4. Do not allow spray to go beyond the edge of the cultivated area. Spray the outside row only from outside the planting.

TURFGRASS

Crop	Pest	Product Rate	Remarks
Turfgrasses Residential Recreational Commercial	Mole cricket Southern mole cricket Tawny mole cricket	2.7 lbs. per acre (1 oz. per 1000 sq. ft.) 0.54 lbs. a.i. per acre	Make application prior to or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, tank mix with a curative insecticide
	White grub larvae such as: Annual bluegrass weevil Asiatic garden beetle Billbug Black Turfgrass ataeinus European chafer Green June beetle Japanese beetle May/June beetle Northern masked chafer Oriental beetle Southern masked chafer	2.7 lbs. per acre (1 oz. per 1000 sq. ft.) 0.54 lbs. a.i. per acre	For optimum control of grubs, billbugs, and annual bluegrass weevil, make application prior to or during egg hatch of the target pest.
	Cutworms Chinchbug Sod webworm	2.7 lbs. per acre (1 oz. per 1000 sq. ft.) 0.54 lbs. a.i. per acre	For suppression of chinchbugs, make application prior to hatching of the first instar nymphs.
	European Cranefly	2.7 lbs. per acre (1 oz. per 1000 sq. ft.) 0.54 lbs. a.i. per acre	Apply in the spring, when larvae are mature but prior to pupation or in the fall prior to egg hatch

Important Notes:

- Apply in sufficient water to ensure thorough coverage of target area. Use a minimum of 50 gallons finished spray per acre.
- Consult your local State Extension Service or State Extension Turfgrass Specialists for more specific information on timing of insecticide applications.
- For optimal control, apply irrigation if rainfall does not occur within 24 hours after application to ensure movement of the active ingredient through the thatch.
- Do not apply more than a total of 2.7 lbs. of Dinotefuran 20% Turf and Ornamental (0.54 lbs. a.i.) per acre of turf per year.

ORNAMENTAL PLANTS, VEGETABLE TRANSPLANTS AND FORESTS

Foliar or broadcast spray application

For foliar insect control on ornamental plants in nurseries, greenhouses, interior plantscapes, lath and shadehouses, outdoor landscapes (commercial, industrial, recreational and residential), tree plantations, reforestation nurseries and forests.

Crop	Pest	Product Rate	Remarks
------	------	--------------	---------

<p>Ornamental plants including:</p> <p>Shrubs</p> <p>Bedding Plants</p> <p>Flowering Plants</p> <p>Foliage Plants</p> <p>Ground Covers</p> <p>Evergreens</p> <p>Ornamental Trees</p> <p>Non-Bearing Fruit Trees</p> <p>Non-Bearing Nut Trees</p> <p>Non-Bearing Vines</p> <p>Christmas Trees</p> <p>Trees in Plantations including:</p> <p>Conifers</p> <p>Deciduous trees</p> <p>Reforestation Nurseries</p> <p>Forests and Wooded Areas</p> <p>National, Private and State</p> <p>Not for use on house plants grown inside private residences</p>	<p>Adelgids including:</p> <p>Hemlock Woolly</p> <p>Balsam Woolly</p> <p>Aphids (suppression) including</p> <p>Balsam</p> <p>Crepe Myrtle</p> <p>Green Peach</p> <p>Melon</p> <p>Japanese beetles (adults)</p> <p>Lacebugs including:</p> <p>Azalea</p> <p>Cotoneaster</p> <p>Hawthorne</p> <p>Rhododendron</p> <p>Leaf beetles</p> <p>Viburnum</p> <p>Leafhoppers including</p> <p>Glassy-Winged Sharpshooter</p> <p>Potato</p> <p>Leafminers including:</p> <p>Serpentine</p> <p>Mealybugs including:</p> <p>Citrus</p> <p>Long-Tailed</p> <p>Madeira</p> <p>Obscure</p> <p>Phornium</p> <p>Pink Hibiscus</p> <p>Psyllids including:</p> <p>Asian Citrus</p> <p>Root Weevils (adults) including:</p> <p>Black Vine</p> <p>Diaprepes</p> <p>Sawflies (larvae)</p> <p>Scales (Armored and Soft) including:</p> <p>Cryptomeria</p> <p>Cycad Aulacaspis</p> <p>Elongate Hemlock</p> <p>Euonymus</p> <p>Florida Red</p> <p>Florida Wax</p> <p>Tea</p> <p>Thrips including</p> <p>Chilli</p> <p>Gynaikothrips uzeli</p> <p>Western Flower (Suppression)</p> <p>Whiteflies including:</p> <p>Fig (Ficus)</p> <p>Giant</p> <p>Greenhouse</p> <p>Silverleaf /Sweetpotato (B and Q Biotypes)</p>	<p>Foliar Spray</p> <p>¼ to ½ lb. per 100 gallons</p> <p>(4 to 8 oz. per 100 gallons)</p> <p>(0.05 to 0.1 lbs. a.i. per 100 gallons)</p> <p>8-16 oz per Acre</p> <p>(0.1 to 0.2 lbs. a.i./A)</p> <p>0.2-0.4 oz per 1,000 sq. ft.</p> <p>For treatment of small areas:</p> <p>½-1.0 tsp per gallon</p>	<p>For optimal control, make first application just before pest populations reach an economic threshold. If necessary, make a second application after 14-21 days.</p> <p>Tank mixing with a surfactant may improve control of pests such as whitefly, mealybug and scale. Confirm plant safety of tank mix in small area before using on a commercial scale.</p> <p>100 gals. of spray mix will treat 20,000 sq. ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.</p>
---	--	--	---

15/19

1 level teaspoon contains 2.4 grams and 1 cup (8 fl oz) contains 4.0 oz by weight of Dinotefuran 20% Turf and Ornamental.

Do not apply more than 2.7 lbs (0.54 lbs. a.i.) per acre of nursery, landscape or forest per year. To delay the development of resistance: Do not apply Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides to consecutive generations of the same insect species without switching to a different mode of action. Do not make more than two sprays of Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides to a single crop during a two-month period. Refer to "Resistance Management" section of label for further guidelines.

VEGETABLE TRANSPLANTS

Foliar or broadcast spray application			
For foliar insect control on vegetable transplants grown in enclosed structures.			
Crop	Pest	Product Rate (By Weight)	Remarks
Cucurbits (Transplants only) Cantaloupe, Cucumber, Melons, Squash	Aphids Leafminers Mealybugs Thrips (suppression)	3.5 - 7.0 oz per 100 gal	Do not make more than one application per crop.
		7-14 oz per Acre	Apply only to cucurbits and brassica being grown as transplants and before transplants are sold.
Fruiting Vegetables Eggplant, Peppers, Tomato Head and Stem Brassica Broccoli, Brussel Sprouts, Cabbage, Cauliflower, Kohlrabi	Whiteflies including Silverleaf/Sweetpotato (B and Q Biotypes)	0.16 - 0.32 oz per 1,000 sq ft	100 gals. of spray mix will treat 20,000 sq. ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.
		(0.09 to 0.18 lbs. a.i. per Acre)	
Leafy Vegetables (Transplants only) (Excluding Brassica spp.)	Aphids Leafminers Mealybugs Thrips (suppression) Whiteflies including Silverleaf/Sweetpotato (B and Q Biotypes)	3.5 - 5.5 oz. Per 100 gal	Do not make more than one application per crop.
		7-11 oz per Acre	Apply only to leafy vegetables being grown as transplants and before transplants are sold
		0.16 - 0.25 oz. per 1,000 sq ft	100 gals. of spray mix will treat 20,000 sq. ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area
		(0.09 to 0.134 lbs. a.i. per Acre)	
1 level teaspoon contains 2.4 grams and 1 cup (8 fl oz) contains 4.0 oz by weight of Dinotefuran 20% Turf and Ornamental.			
Do not apply more than 1.34 lbs (0.268 lbs. a.i.) per acre of nursery per year.			
To delay the development of resistance: Do not apply Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides to consecutive generations of the same insect species without switching to a different mode of action. Do not make more than two sprays of Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides to a single crop. Refer to "Resistance Management" section of label for further guidelines.			

GREENHOUSE GROWN TOMATOES

Foliar spray, broadcast spray or soil drench application

For foliar insect control on tomatoes grown and harvested in enclosed structures.

Crop	Pest	Product Rate (By Weight)	Remarks
Tomatoes	Aphids Leafminers Mealybugs Thrips (suppression) Whiteflies including Silverleaf/Sweetpotato (B and Q Biotypes)	Foliar Spray 4 to 8 oz. per 100 gal 8-16 oz per Acre (0.1 to 0.2 lbs. a.i. per Acre) 0.2-0.4 oz per 1,000 sq. ft.	100 gals. of spray mix will treat 20,000 sq. ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area Do not apply more than two foliar sprays per crop Do not apply more than 2.0 lbs (0.4 lbai) per acre per year as foliar sprays Do not apply soil drench to plants that have received foliar sprays Do not apply within one (1) day of harvest.
		Soil Drench ¾ to 1½ pounds per 100 gallons 12 to 24 ounces per 100 gallons 1.5-3.0 teaspoons per gallon Apply 4 fl oz of drench solution per gallon of potting media	Only apply soil drench to moist soil media. Do not apply to dry or saturated media. Do not leach treated soil media for at least 7 days after application or performance may be reduced. Do not apply more than 2.7 lb (0.54 lbai) per acre per year as a soil drench. Do not apply foliar sprays to plants have received a soil drench Do not apply within one (1) day of harvest.

One level teaspoon contains 2.4 grams and 1 cup (8 fl oz) contains 4.0 oz by weight of Dinotefuran 20% Turf and Ornamental.

To delay the development of resistance: Do not apply Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides to consecutive generations of the same insect species without switching to a different mode of action. Do not make more than two sprays or one soil drench of Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides to a single crop. Refer to "Resistance Management" section of label for further guidelines.

PESTS CONTROLLED BY DINOTEFURAN 20% TURF AND ORNAMENTAL WHEN APPLIED TO THE SOIL

<p>Adelgids including: Hemlock Woolly Balsam Woolly</p> <p>Aphids including Balsam Crepe Myrtle Green Peach Melon Bagworms Eastern Tent Caterpillar Erythrina Gall Wasp Flatheaded Borers including Alder Bronze Birch Emerald Ash Flatheaded Appletree Two-Lined Chestnut Froghoppers</p> <p>Fungus Gnats (larvae) Gypsy Moth (larvae) Horned Oak Gall Japanese Beetle (Adults)</p> <p>Lacebugs including: Azalea Cotoneaster</p> <p>Hawthorne Rhododendron</p> <p>Leaf Beetles Elm Viburnum</p> <p>Leafhoppers including Glassy-Winged Sharpshooter Potato</p> <p>Leafminers including: Birch</p> <p>Boxwood Holly Serpentine</p>	<p>Mealybugs Citrus Longtailed Madeira Obscure Phormium Pink Hibiscus</p> <p>Root Mimosa Webworm Pine Tip Moth (larvae)</p> <p>Plantbugs</p> <p>Psyllids including: Asian Citrus Boxwood</p> <p>Root Weevils (larvae and adults) including Black Vine</p> <p>Diaprepes</p> <p>Roundheaded Borers (excluding Asian Longhorned)</p> <p>Eucalyptus Longhorned</p> <p>Linden Locust Royal Palm Bug Sawfly larvae Scales (Armored and Soft)</p> <p>Including: Azalea Bark Brown Soft Calico California Red</p> <p>Cottony Cushion Cottony Maple Cryptomeria</p>	<p>Scales (Armored and Soft) continued Cycad Aulacaspis Duplachionaspis Elongate Hemlock Euonymus False Oleander Fig (Ficus) Wax Fletcher Florida Red Florida Wax Indian Wax</p> <p>Lecanium Lobate Lac Obscure Oystershell Poplar (Aspen) Pine Needle</p> <p>Tea Tuplptree Spittlebugs</p> <p>Tent Caterpillar</p> <p>Thrips including: Chilli (Suppression) Citrus Cuban Laurel Gladiolus Gynaikothrips uzeli (Suppression) Western Flower (Suppression)</p> <p>Treehoppers Walnut Twig Beetle</p> <p>Whiteflies including: Fig (Ficus) Giant Greenhouse Silverleaf/Sweetpotato (B and Q biotypes)</p> <p>White Grubs including: Oriental Beetle White Pine Weevil</p>
--	--	--

Application to soil

For systemic insect control on containerized and field grown (in-ground) ornamental plants in nurseries, greenhouses, interior plantscapes, lath and shadehouses, outdoor landscapes (commercial, industrial, recreational and residential), tree plantations, reforestation nurseries and forests when applied via soil drench, soil injection, micro-irrigation (spaghetti tube or emitter), drip irrigation, overhead irrigation, ebb and flood irrigation equipment or motorized irrigation equipment.

Crop	Pest	Product Rate (By Weight)		Remarks
<p>Ornamental plants including:</p> <ul style="list-style-type: none"> Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines <p>Christmas Trees</p> <p>Trees in Plantations including:</p> <ul style="list-style-type: none"> Conifer Deciduous <p>Reforestation Nurseries</p> <p>Forests and Wooded Areas National, Private and State</p>		<p>Containerized Plants</p> <p>Soil Media Drench</p> <p>¼ to 1½ pounds per 100 gallons</p> <p>12 to 24 ounces per 100 gallons</p> <p>1.5-3.0 teaspoons per gallon</p>		<p>Only apply to moist soil media.</p> <p>Do not apply to dry or saturated media.</p> <p>For optimal performance, do not apply media drench until roots from transplanted plugs or liners have extended at least half way to the edge of pots.</p> <p>Do not leach treated soil media for at least 7 days after application or performance may be reduced.</p> <p>Heavy rainfall or excessive irrigation following application may decrease performance.</p>
		<p>Media Drench Volume for Individual Pots</p>		
		Pot diameter (inches)	Fl oz of dilute solution per pot	<p>In general, higher rates will be needed to control insects on woody plants than on herbaceous plants.</p> <p>Poinsettia: For optimal control of whiteflies, treat plants 1-3 weeks after pinch. Late season drenches will take longer to provide effective control.</p>
		4	2	
		5	3	
		6	4	
		7	5	
		8	6	
<p>For larger pot volumes, apply 3 – 4 fl oz of dilute solution (0.11 to 0.22 g product per 4 fl oz water,) per gallon of potting media. Use a drench volume that is sufficient to wet soil media without resulting in overflow or runoff through drain holes in pot.</p>				

		<p>Containerized Plants</p> <p>Media Drench Volume for Plants in Raised Beds, Benches, Bedding Flats, Plug and Liner Trays:</p> <p>Apply sufficient dilute solution to wet soil media without loss of liquid from bottom of bed or liner.</p>													
		<p>Containerized Plants</p> <p>Ebb and Flood</p> <table border="1"> <thead> <tr> <th>Pot diameter (inches)</th> <th>Ounces per 1,000 pots</th> </tr> </thead> <tbody> <tr> <td>4</td> <td>1.9 - 3.7</td> </tr> <tr> <td>5</td> <td>2.8 - 5.6</td> </tr> <tr> <td>6</td> <td>3.7 - 7.5</td> </tr> <tr> <td>7</td> <td>4.7 - 9.3</td> </tr> <tr> <td>8</td> <td>5.6 - 11.2</td> </tr> </tbody> </table>	Pot diameter (inches)	Ounces per 1,000 pots	4	1.9 - 3.7	5	2.8 - 5.6	6	3.7 - 7.5	7	4.7 - 9.3	8	5.6 - 11.2	<p>Bring several pots to field capacity, let soil dry and then measure amount of water required to bring pots back to field capacity. Multiply the average volume of water required to rehydrate one pot by the number of pots to be treated. Add this volume of water to the minimum amount of water needed to flood the area to be treated. Re-use any returned volume in subsequent irrigation of same plants.</p> <p>For pot diameter greater than 8", use 3.7-7.5 ounces of Dinotefuran 20% Turf and Ornamental per 1,000 gallons of potting soil media.</p>
Pot diameter (inches)	Ounces per 1,000 pots														
4	1.9 - 3.7														
5	2.8 - 5.6														
6	3.7 - 7.5														
7	4.7 - 9.3														
8	5.6 - 11.2														
		<p>Chemigation of individual containers using a micro-irrigation system (spaghetti tube)</p> <table border="1"> <thead> <tr> <th>Injection ratio</th> <th>Ounces per gallon of injector tank water</th> </tr> </thead> <tbody> <tr> <td>1:100</td> <td>12 - 24</td> </tr> </tbody> </table>	Injection ratio	Ounces per gallon of injector tank water	1:100	12 - 24	<p>Use typical injection ratio for injectors (e.g. 1:100, which equals 1 part injector tank solution: 100 parts irrigation water). Do not mix more than 24 oz of Dinotefuran 20% Turf and Ornamental per gallon of injector tank water, or some product may settle out of solution. Calibrate irrigation system to deliver 3-4 fl oz of dilute solution per gallon of potting media.</p>								
Injection ratio	Ounces per gallon of injector tank water														
1:100	12 - 24														

		<p>Field Grown (In-Ground) Plants</p> <p>Shrubs</p> <p>3 - 6 grams (1.25 - 2.5 level teaspoons) per foot of height</p> <p>1.0 - 2.1 ounces per 10 feet of height</p>	<p>When applied to the soil, Dinotefuran 20% Turf and Ornamental is taken up by actively growing trees and shrubs. Speed of control will be dependent on tree size, tree health, environmental conditions and how actively pests are feeding. In actively growing and transpiring plants, control may be evident within one week after application in small shrubs, and within three weeks after application in large trees. For optimal control, time soil applications so that Dinotefuran 20% Turf and Ornamental has had time to move to insect feeding sites by time target life stage is present. Control may be less effective when applied to dry, saturated, or frozen soil, or at times when plants are not actively taking up water from soil.</p> <p>If possible, irrigate dry soils 1-3 days before application, or apply irrigation within 3 days after application.</p> <p>Heavy rainfall or inadequate irrigation immediately following application may decrease performance.</p> <p>Use higher labeled rates for broadleaf evergreens with dense foliage (ex. hollies), and with very large trees.</p> <p>Soil Drench: Mix required dose in water and uniformly apply to soil around base of shrub or tree. Pull back mulch before drenching. For optimal performance, apply 1-4 pints of drench solution per foot of height (shrubs) or inch of trunk diameter (trees). Adjust drench volume based on soil type, soil moisture and thickness of mulch so that product is moved into root zone. To enhance soil penetration in heavy soils and sloping terrain, dig shallow holes around tree or shrub, and apply drench solution in holes. Lower drench volumes may be less effective in dry soils or when applied over heavy mulch unless there is adequate rainfall or irrigation after application to move product into root zone.</p>
--	--	--	--

		<p>Small Trees (Less than 24" diameter at breast height)</p> <p>3 - 12 grams (1.25 - 5.0 level teaspoons) per inch of trunk diameter at breast height (DBH)</p> <p>1.05 - 4.2 ounces per 10 inches of trunk diameter at breast height (DBH)</p> <p>For multi-stem trees, base rate on cumulative inches of diameter of all stems at breast height.</p>	<p>Soil Injection: Mix required dose in water and make at least four injections per shrub or tree with a low-pressure applicator. Use same amount of solution per hole. Injections can be made using the following methods:</p> <p><i>Grid System</i>-Space injections on a 2.5 ft center extending to drip line.</p> <p><i>Circle System</i>-Make injections in concentric circles extending inward from drip line.</p> <p><i>Basal System</i>- Space injections evenly around trunk no more than 24" out from the base.</p>														
		<p>Large Trees (24" diameter or greater at breast height)</p> <p>6 - 12 grams (2.5 - 5.0 level teaspoons) per inch of trunk diameter at breast height (DBH)</p> <p>2.1 - 4.2 ounces per 10 inches of trunk diameter at breast height (DBH)</p> <p>For multi-stem trees, base rate on cumulative inches of diameter of all stems at breast height.</p>	<p>Dinotefuran 20% Turf and Ornamental may be soil injected with low volume (e.g. Kioritz injector) or high volume injection equipment. For optimal performance, inject 1 - 32 fl oz of dilute solution per foot of height or inch of trunk diameter depending on application equipment. Make shallow injections where feeder roots are most concentrated.</p>														
		<p>Hedges (Soil drench)</p> <p>0.5 - 2.0 oz per foot of height per 100 linear feet of row</p>	<p>Apply 25-100 gallons of drench solution per 100 linear feet of hedge row. Use higher volume for drier soils. Apply in a one foot wide band over base of trunk and soil down center of hedgerow. To improve performance, rake back mulch before application.</p>														
		<p>Banded spray application to soil surface</p> <p>(2.7 lbs per acre)</p> <table border="1" data-bbox="792 1270 1104 1659"> <thead> <tr> <th>Row spacing in feet</th> <th>Ounces per 1,000 linear feet of row</th> </tr> </thead> <tbody> <tr> <td>3</td> <td>3</td> </tr> <tr> <td>4</td> <td>4</td> </tr> <tr> <td>5</td> <td>5</td> </tr> <tr> <td>6</td> <td>6</td> </tr> <tr> <td>7</td> <td>7</td> </tr> <tr> <td>8</td> <td>8</td> </tr> </tbody> </table>	Row spacing in feet	Ounces per 1,000 linear feet of row	3	3	4	4	5	5	6	6	7	7	8	8	<p>Apply as a uniform band in row over root zone. Apply from peak adult flight to peak egg hatch.</p> <p>Apply in at least two gallons of water per 1,000 linear feet. Irrigate after application to move product into soil profile.</p> <p>Control any weeds in treated area prior to application, or performance may be reduced.</p> <p>Adjust rates accordingly for other row spacing.</p> <p>For optimal performance, irrigate after application to move Dinotefuran 20% Turf and Ornamental into the root zone.</p>
Row spacing in feet	Ounces per 1,000 linear feet of row																
3	3																
4	4																
5	5																
6	6																
7	7																
8	8																

		<p>Broadcast spray to soil of plant beds</p> <p>2.7 lbs per acre</p>	<p>Apply over the top of ornamental plant beds in a water volume sufficient to move product to soil surface. If necessary, irrigate after application to move product off of foliage and into upper root zone of soil. May be less effective on large woody shrubs than on herbaceous annuals and perennials.</p>
<p>1 level teaspoon contains 2.4 grams, and 1 cup (8 fl oz) contains 4.0 oz by weight of Dinotefuran 20% Turf and Ornamental.</p> <p>Do not apply more than 2.7 lbs (0.54 lbs. a.i.) per acre of nursery, landscape or forest per year.</p> <p>To delay the development of resistance in greenhouses, nurseries and interiorscapes, do not make more than one soil application per crop cycle or three-months, whichever is shorter. Refer to "Resistance Management" section of the label for additional guidelines.</p>			

<p>Trunk sprays in trees and large shrubs</p> <p>For systemic insect control in containerized and field grown (in-ground) ornamental trees and shrubs in nurseries, interior plantscapes, lath and shadehouses and outdoor landscapes (commercial, industrial, recreational and residential), tree plantations, and forests when applied as a trunk spray.</p>			
Crop	Pest	Product Rate (By Weight)	Remarks

<p>Shrubs</p> <p>Ornamental trees</p> <p>Non-Bearing Fruit Trees</p> <p>Non-Bearing Nut Trees</p> <p>Christmas Trees</p> <p>Trees in Plantations including: Conifer Deciduous</p> <p>Reforestation Nurseries</p> <p>Forests and Wooded Areas National, Private and State</p>	<p>Adelgids including Hemlock Woolly Aphids</p> <p>Flatheaded Borers including Alder Bronze Birch Emerald Ash Flatheaded Appletree Two-lined Chestnut</p> <p>Lacebugs</p> <p>Leaf Beetles</p> <p>Leafhoppers</p> <p>Leafminers</p> <p>Mealybugs</p> <p>Mountain Pine Beetle</p> <p>Pine Tip Moth (Larvae)</p> <p>Psyllids</p> <p>Roundheaded Borers (Excluding Asian Longhorned)</p> <p>Scales including Calico Elongate Hemlock Fig (Ficus) Wax</p> <p>Thrips (Suppression)</p> <p>Walnut Twig Beetle</p> <p>Whiteflies Fig (Ficus)</p>	<p>12 – 24 oz per gallon</p> <p>Depending on bark type and thickness, one gallon of spray solution will cover 40-80" of trunk diameter when measured at 4.5 feet above ground.</p>	<p>When sprayed on the trunk, Dinotefuran 20% Turf and Ornamental will be absorbed through the bark and into the vascular system, and then transported throughout the tree. Speed of control will be dependent on tree size, tree health, environmental conditions and how actively pests are feeding. In actively transpiring trees, control may be evident within 1-4 weeks after application.</p> <p>Spray bark on root flare (buttress roots) and on trunk between soil surface and 4-5 feet above the soil surface.</p> <p>Adjust nozzle to uniformly distribute spray over the entire circumference of the tree trunk and buttress roots. Wet bark just to the point of saturation and run off onto soil.</p> <p>Apply ONLY with a low volume sprayer operated at 10-20 PSI to prevent tree damage, bounce back and drift.</p> <p>For optimal control, apply to actively growing trees, and time application so that Dinotefuran 20% Turf and Ornamental has had time to move to insect feeding sites at when target life stage is present.</p> <p>Do not apply to wet bark, during rainfall or if rain is expected within 12 hours.</p> <p>Control may be less effective in trees with thick bark, and at times when trees are not actively growing or transpiring.</p> <p>For Mountain Pine Beetle; apply between two weeks before and two weeks after peak of expected flight activity.</p>
--	--	--	--

1 level teaspoon contains 2.4 grams, and 1 cup (8 fl oz) contains 4.0 oz by weight of Dinotefuran 20% Turf and Ornamental

Do not apply more than 2.7 lbs (0.54 lbs. a.i.) per acre of nursery or landscape per year.

Trunk injection in trees and large shrubs

For systemic insect control in containerized and field grown (in-ground) ornamental trees and large shrubs in greenhouses, nurseries, interior plantscapes, lath and shadehouses, outdoor landscapes (commercial, industrial, recreational and residential), tree plantations, reforestation nurseries and forests when applied via trunk injection.

Crop	Pest	Product Rate	Remarks
------	------	--------------	---------

<p>Ornamental trees Non-Bearing Fruit Trees Non-Bearing Nut Trees Christmas Trees Trees in Plantations including: Conifer Deciduous Reforestation Nurseries Forests and Wooded Areas - National, Private and State</p>	<p>Adelgids Aphids Erythrina Gall Wasp Flatheaded Borers Lacebugs Leaf Beetles Leafhoppers Leafminers Mealybugs Pine Tip Moth (Larvae) Psyllids Roundheaded Borers Scales Thrips (Suppression) Whiteflies</p>	<p>1.0-2.0 grams per inch of trunk diameter at breast height</p>	<p>Trunk Injection: Measure diameter of tree at breast height. Using a 1/4-3/8" drill bit, drill two holes into trunk for each inch of trunk diameter. Drill holes 3/8-1/2" deep into healthy xylem. Rill holes at a 45-degree downward slant, and evenly space at 4-8" above the ground. Place feeder tube in each hole and position so that the tip is seated in the conductive zylem tissue.</p> <p>Mix required amount of product in water, and inject solution into each feeder tube. Adjust water volume according to type of injection equipment. Uptake can take several minutes. Remove tubes immediately after uptake is completed.</p> <p>For optimal control, make treatments before egg hatch.</p> <p>Do not inject trees less than two inches in diameter.</p>
<p>1 level teaspoon contains 2.4 grams, and 1 cup (8 fl oz) contains 4.0 oz by weight of Dinotefuran 20% Turf and Ornamental Do not inject trees more than one time per year.</p>			

25
/26

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Pesticide Storage: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in a cool dry place. Do not store diluted spray. For help with any spill, leak, fire or exposure involving this material, call day or night 1-800-CLEANUP.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Triple rinse (or equivalent). Do not reuse container. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE

MITSUI CHEMICALS, INC., warrants that this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions to the crops specified. To the extent consistent with applicable law, there are no other warranties, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, this warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or conditions not reasonably foreseeable to seller, and buyer assumes all risk of any such use.

070909

NEXT

LABEL

33657-16

4/13/2010

1/22



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

APRIL
~~FEB~~ 13 2010

Name Ms. Peggy Galloway
Authorized Agent for Mitsui Chemicals

Company Landis International

Address P.O. Box 5126
Valdosta, GA 31603-5126

Subject: Notification: Minor label revision
Amended container handling language per PR Notice 2007-4
Dinotefuran 20% Turf, Ornamental and Vegetable Transplants
EPA Registration Number 33657-16

Your submission pin punched Feb. 12, 2010

Dear Ms. Galloway:


The Agency is in receipt of your Application dated Feb. 12, 2010 for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 for the product Dinotefuran 20% Turf, Ornamental and Vegetable Transplants. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-308-8291.

Sincerely,

Rita Kumar

Rita Kumar
Senior Regulatory Specialist
Insecticide-Rodenticide Branch
Registration Division (7505P)
Office of Pesticide Programs
E-mail: kumar.rita@epa.gov

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
---	---	--	-----------------------

Application for Pesticide - Section I

1. Company/Product Number 33657-16	2. EPA Product Manager Rita Kumar	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Dinotefuran 20% Turf, Ornamental, & Vegetable Transplants	PM# IRB-Team 1	
5. Name and Address of Applicant (Include ZIP Code) MITSUI CHEMICALS, INC. c/o Landis International, Inc. P. O. Box 5126, Valdosta, GA 31603-5126 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: <input checked="" type="checkbox"/> EPA Reg. No. _____ Product Name _____	

NOTIFICATION
APR 13 2010

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of PR Notice 2007-4 Container Handling statement amendment per PR Notice 98-10. See attached statement.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No			
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt No. per container	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information		4. Size(s) Retail Container		5. Location of Label Directions	
<input type="checkbox"/> Label <input type="checkbox"/> Container				<input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product				<input type="checkbox"/> Other _____	
<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					


Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name PEGGY M. GALLOWAY	Title Regulatory Agent	Telephone No. (Include Area Code) 229-247-6422
----------------------------------	----------------------------------	--

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.


2. Signature 	3. Title Regulatory Agent	6. Date Application Received (Stamped) _____
4. Typed Name PEGGY M. GALLOWAY	5. Date Feb 10, 2010	

**U.S. ENVIRONMENTAL PROTECTION AGENCY
Application of Pesticide Amendment (EPA Form 8570-1)**

**DINOTEFURAN 20% TURF, ORNAMENTAL AND
VEGETABLE TRANSPLANTS IN ENCLOSED STRUCTURES
EPA Registration No. 33657-16**


Section - II. Explanation (cont.)

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the Confidential Statement of Formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this Notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.



Peggy M. Galloway, Regulatory Agent
MITSUI CHEMICALS, INC.

02/10/2010
Date





4
22

February 10, 2010

VIA FEDERAL EXPRESS

NOTIFICATION
APR 13 2010

Ms. Rita Kumar, Senior Regulatory Specialist
Insecticide-Rodenticide Branch, Team 1
U. S. EPA Office of Pesticide Programs - H7505P
Document Processing Desk (NOTIF)
One Potomac Yard, Room S4900
2777 S. Crystal Drive
Arlington, VA 22202

**SUBJECT: DINOTEFURAN 20% TURF, ORNAMENTAL AND VEGETABLE
TRANSPLANTS - EPA Registration No. 33657-16**

86203-11

Dear Ms. Kumar:

Please find the following documents enclosed on behalf of MITSUI CHEMICALS, INC.:

- Application for Pesticide Amendment - NOTIFICATION.
- One copy of amended labeling with changes "highlighted".
- Two copies of "clean" amended labeling.

In accordance with PR Notice 98-10, we are submitting the enclosed amended labeling by NOTIFICATION which amends the "Container Handling" statement in compliance with PR Notice 2007-4.


Please let me know if you have any questions. We appreciate your continued assistance with this registration.

Best regards,

A handwritten signature in cursive script that reads "Peggy M. Galloway".

Peggy M. Galloway
Regulatory Agent for
MITSUI CHEMICALS, INC.

Enclosures: Enumerated Above



H:\MITSUI 44620% T&O\RK021010L.NOTIFICATION

5
22
NOTIFICATION

APR 13 2010

GROUP	4A	INSECTICIDE
-------	----	-------------

DINOTEFURAN 20 %

TURF, ORNAMENTAL AND VEGETABLE TRANSPLANTS IN ENCLOSED STRUCTURES

For foliar and systemic insect control in ornamental plants, vegetable transplants in enclosed structures,
and turfgrass.

For Greenhouse, Nursery, Interior Plantscape and Outdoor Landscape Use Only

Active Ingredient:

Dinotefuran, [N-methyl-N'-nitro-N''-((tetrahydro-3-furanyl)methyl)guanidine]	20%
Other Ingredients	80%
Total:	100%

KEEP OUT OF REACH OF CHILDREN
CAUTION
See side panels for additional precautionary statements

FIRST AID	
If On Skin Or Clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for further treatment advice.
If Swallowed	<ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Have person sip a glass of water if able to swallow.• Do not give anything by mouth to an unconscious person.
If In Eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for further treatment advice.
If Inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact CHEMTREC (800) 424-9300 (24 hours) for emergency medical treatment information.	
<i>See side/back panels for additional precautionary statements</i>	

EPA Reg No. 33657-16

EPA Establishment No. 67545-AZ-01

Net Contents:
Manufactured By:
MITSUI CHEMICALS, INC.
Shiodome City Center
1-5-2 Higashi-Shimbashi
Minato-ku, Tokyo 105-7117
JAPAN

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Do not allow contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to shrimp. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not dispose equipment washwaters or rinsate into a natural drain or water body.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Dinotefuran and its degradate, MNG have the properties and characteristics associated with chemicals detected in ground water. The high water solubility of dinotefuran, and its degradate, MNG, coupled with its very high mobility, and resistance to biodegradation indicates that this compound has a strong potential to leach to the subsurface under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination. Periodic monitoring of shallow groundwater in the use area is recommended.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

SPRAY DRIFT ADVISORY

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crop thereof rendered for sale, use or consumption.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL, USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural insecticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

EXCEPTION: If product is drenched or soil-injected, workers may enter the area at any time if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. **Do not allow others to enter treated areas until sprays have dried.**

APPLICATION INFORMATION

- **Applications of Dinotefuran 20% Turf and Ornamental in residential areas may be made by commercially licensed applicators.**

Application to Turfgrass:

- Dinotefuran 20% Turf and Ornamental can be used for the control of soil inhabiting pests of turfgrass such as Masked Chafers, European Chafer, Green June Beetle, May or June Beetle, Japanese Beetle, Oriental Beetle, Billbugs, Annual Bluegrass Weevil, Black Turfgrass Ataenius and Mole Crickets. Dinotefuran 20% Turf and Ornamental can also be used for the suppression of cutworms and chinchbugs in turfgrass areas.
- Dinotefuran 20% Turf and Ornamental can be used as directed on outdoor residential, recreational and commercial turfgrass in sites such as home lawns, commercial lawns, multi-family residential and apartment complexes, grounds or lawns around business and office complexes, shopping centers, airports, military and other institutions, cemeteries, golf courses, playgrounds, parks, athletic fields and sod farms.

- Target timing of Dinotefuran 20% Turf and Ornamental applications at or just prior to or during egg laying of the target pests. The need for an application can be based on historical and/or physical monitoring of the site, current season adult trapping, previous experience or other methods. Optimum control will be achieved when applications are made prior to or at egg hatch of the target pests followed by sufficient irrigation or rainfall to move the active ingredient through the turf thatch layer. Consult your State Extension Service for information regarding specific application timing.
- Do not apply when the target site is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist.

Application to Ornamental plants:

- Dinotefuran 20% Turf and Ornamental can be applied as a foliar spray, a broadcast spray, a soil drench, soil injection and via chemigation for insect control in ornamental plants in greenhouses, nurseries, outdoor landscapes and interior plantscapes.
- Dinotefuran 20% Turf and Ornamental is a systemic product and will be taken up by the root system and translocated upward throughout the plant. When applied as a foliar spray, the product offers translaminar and locally systemic control of foliar pests.
- When applied to the soil, Dinotefuran 20% Turf and Ornamental will be translocated more quickly in herbaceous plants than in woody shrubs and trees. Speed of insect control will range from as little as one day for small herbaceous plants in containers, to several weeks in large trees in growing in the landscape.
- For outdoor and landscape ornamentals, broadcast applications cannot exceed a total of 2.7 lbs of product (0.54 lb. active ingredient) per acre per year.

MIXING INSTRUCTIONS:

Dinotefuran 20% Turf and Ornamental Alone: Add half of the required amount of water to the mix tank. With the agitator running, add the desired amount of Dinotefuran 20% Turf and Ornamental to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after Dinotefuran 20% Turf and Ornamental has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Dinotefuran 20% Turf and Ornamental + Tank Mixtures: Add half of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. In general, add tank mix partners in this order: products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables), liquid flowables, liquids, emulsifiable concentrates, and surfactants/adjuvants. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all the mixture has been applied.

NOTE: When using Dinotefuran 20% Turf and Ornamental in tank mixtures, add all products in water-soluble packaging to the tank before any other tank mix partner, including Dinotefuran 20% Turf and Ornamental. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using Dinotefuran 20% Turf and Ornamental in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label. Do not exceed label dosage rate, and follow the most restrictive label precautions and limitations. Do not mix this product with any product that prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

Compatibility

IMPORTANT - The safety of all potential tank mixes has not been tested on all crops. Before applying any tank mixture not specifically listed on this label, confirm the safety to the target crop.

Dinotefuran 20% Turf and Ornamental is compatible with most commonly used pesticides, crop oils, adjuvants, and nutritional sprays. However, since it is not possible to test all possible mixtures, pre-test to assure the physical compatibility and lack of phytotoxic effect of any proposed mixtures with Dinotefuran 20% Turf and Ornamental. To determine the physical compatibility of Dinotefuran 20% Turf and Ornamental with other products, use a jar test, as described below:

Using a quart jar, add the proportionate amounts of the products to 1 quart of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for additional required ingredients to the spray tank.

RESISTANCE MANAGEMENT

Dinotefuran 20% Turf and Ornamental contains a Group 4A insecticide. Insect biotypes with acquired resistance to Group 4A may eventually dominate the insect population if Group 4A insecticides are used repeatedly in the same crop or in successive years as the primary method of control for a targeted species. This may result in partial or total loss of control of those species by Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides.

To delay the development of insecticide resistance in greenhouse, nursery and interiorscape use sites, strongly consider the following guidelines:

- Do not apply Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides to consecutive generations of the same insect pest species.
- Do not drench soil media with Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides more than one time per crop cycle or three months, whichever is shorter.
- Do not make more than two foliar or broadcast sprays of Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides to a single crop during a two-month period.
- Do not make more than one soil drench and one foliar or broadcast spray with Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides during a two-month period.
- Base insecticide use on a comprehensive IPM program.
- Monitor treated insect populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturers for insecticide resistance management and/or IPM guidelines for the specific site and resistant pest problems.
- For further information or to report suspected resistance, you may contact Landis International, Inc., a representative of Mitsui Chemicals, Inc., at toll free number: 1-800-526-3471 or at their web site: www.landisintl.com

APPLICATION PROCEDURES AND SPRAY EQUIPMENT

Ground Application: Select spray nozzles that will provide accurate and uniform spray deposition. Use spray nozzles that provide medium-sized droplets and reduce drift. To help insure accuracy, calibrate sprayer before each use. For information on spray equipment and calibration, consult nozzle manufacturers and/or State Extension Service specialists.

Apply Dinotefuran 20% Turf and Ornamental using sufficient water volume to provide thorough and uniform coverage. In situations where a dense canopy exists and/or pest pressure is high, use greater water volumes. The use of a spray adjuvant may improve spray coverage. Do not apply under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Applications to turfgrass: Apply Dinotefuran 20% Turf and Ornamental through conventional spray equipment in a minimum of 1 gallon of finished spray per 1000 sq. ft. Ensure adequate distribution in the treated area using accurately calibrated equipment normally used for application of turfgrass insecticides. Use equipment that will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off target drift. Check calibration periodically to ensure that equipment is working properly. Prevent skips by using marker dyes or foam aids.

Applications to ornamental plants: Dinotefuran 20% Turf and Ornamental can be applied using many different types of application equipment. Apply in sufficient water to ensure good coverage of ornamental plants. When making applications to plants with hard to wet foliage such as holly or pine, the product may be tank mixed with a surfactant. If concentrate or mist type spray equipment is used, apply the same amount of product on the sprayed area as would be used in a dilute solution. To assure optimum effectiveness, the product must be placed where the growing portion of the target plant can absorb the active ingredient. Applications can be made to foliage or as a soil drench.

RESTRICTIONS

- Do not graze treated areas or use clippings from treated areas for feed or forage.
- Prevent runoff or puddling of irrigation water following application.
- Keep children and pets off treated areas until spray has dried.
- Do not apply to areas that are water logged or saturated, or frozen, which will not allow penetration into the root zone of the plant.
- APPLICATION THROUGH IRRIGATION SYSTEMS (CHEMIGATION):

Dinotefuran 20% Turf and Ornamental may be applied by injection into an irrigation system, either alone or in combination with other pesticides or chemicals that are registered for application through irrigation systems. Dilution ratios are normally 1:100 to 1:200, depending on the system. Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems (Turfgrass) or microirrigation (individual spaghetti tube), drip irrigation, overhead irrigation, or motorized calibrated irrigation equipment (Ornamentals). Do not apply through any other type of irrigation system. Lack of effectiveness can result from non-uniform distribution of treated water. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make adjustments when necessary.

Using Water from Public Water Systems:

DO NOT APPLY DINOTEFURAN 20% TURF & ORNAMENTAL THROUGH ANY IRRIGATION SYSTEM PHYSICALLY CONNECTED TO A PUBLIC WATER SYSTEM.

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year. Dinotefuran 20% Turf and Ornamental may be applied through irrigation systems which may be supplied by a public water system only if the water from the public water system is discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. Before beginning chemigation, always make sure that the air gap exists and that there is no blockage of the overflow of the reservoir tank.

Any irrigation system using water supplied from a public water system must also meet the following requirements:

Operating Instructions for Irrigation Systems:

1. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts.
2. The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
6. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
7. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
8. Do not apply when wind speed favors drift beyond the area intended.

Calibration and Application Instructions:

Apply Dinotefuran 20% Turf and Ornamental under the schedule specified in the specific use instructions, not according to the irrigation schedule unless the events coincide. In general, set the equipment to apply the minimum amount of water per acre. Run the system at 86 - 90% of the manufacturer's maximum rated travel speed.

The following calibration and application techniques are provided for user reference, but do not constitute a warranty of fitness for application through sprinkler irrigation equipment. Check with state and local regulatory agencies for potential use restrictions before applying any agricultural chemical through sprinkler irrigation equipment.

Center Pivot Irrigation Equipment:

NOTES: 1) Use only drive systems that provide uniform water distribution. 2) Do not use end guns when chemigating Dinotefuran 20% Turf and Ornamental through center pivot systems because of non-uniform application. 3) Plug the first nozzle closest to the wellhead to protect the water source.

1. Determine the size of the area to be treated.
2. Determine the time required to apply 0.1 - 0.25 inches of water over the area to be treated when the system and injection equipment are operated at normal pressures as recommended by the equipment manufacturer. Run the system at 80-95% of the manufacturer's rated maximum travel speed.
3. Using water, determine the injection pump output when operated at normal line pressure.
4. Determine the amount of Dinotefuran 20% Turf and Ornamental, and any tank mix partners, required to treat the area covered by the irrigation system.
5. Add the required amount of Dinotefuran 20% Turf and Ornamental, and any tank mix partners, and sufficient water to meet the injection time requirements to the solution tanks. (See "Mixing Instructions" section of this label.)
6. Make sure the system is fully charged with water before starting injection of the Dinotefuran 20% Turf and Ornamental solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
7. Inject the specified amount of Dinotefuran 20% Turf and Ornamental per acre continuously for one complete revolution of the system.
8. Stop the injection equipment after treatment is complete. Continue to operate the system until the Dinotefuran 20% Turf and Ornamental solution has cleared all of the sprinkler heads.
9. Allow time for all lines to flush the pesticide through all nozzles before turning off irrigation water.

Solid Set, Hand Move, and Moving Wheel Irrigation Equipment:

1. Determine the acreage covered by the sprinklers.
2. Fill injector solution tank with plain water and calibrate the flow rate of the system to deliver the contents of the tank over a 20-40 minute time interval.
3. Determine the amount of Dinotefuran 20% Turf and Ornamental required to treat the area covered by the

irrigation system.

4. Add the required amount of Dinotefuran 20% Turf and Ornamental, and any other tank mix partners, into the same quantity of water used to calibrate the injection period. (See "Mixing Instructions" section of this label.)
5. Operate the system at the same pressure and time interval established during the calibration.
6. Inject specified amount of Dinotefuran 20% Turf and Ornamental per acre for either a 20-40 minute period at the end of a regular irrigation set, or as a 20-40 minute injection as a separate application not associated with a regular irrigation to maximize retention of the insecticide by the foliage.
7. Stop injection equipment after treatment is completed. Continue to operate the system until the Dinotefuran 20% Turf and Ornamental solution has cleared the last sprinkler head. To ensure lines are flushed and free from remaining pesticides, a dye indicator may be injected into the lines to mark the end of the application period.

MINIMIZING SPRAY DRIFT

As with all crop protection products, it is important to minimize off-target movement. Do not allow spray to drift onto adjacent land, crops, or aquatic areas. To minimize spray drift:

1. Make applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 10 mph. Do not apply when wind gusts approach 10 mph.
2. Risk of exposure to sensitive aquatic areas can be reduced by not applying when wind direction is toward the aquatic area.
3. Do not cultivate or plant crops within 25 feet of the aquatic area as to allow growth of a vegetative filter strip.
4. Do not make applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with increased height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.
5. Use the largest droplet size consistent with good pest control. Small droplets are more prone to spray drift and can be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by not using excessive spray boom pressure.
6. Apply as close to target plants as practical to obtain a good spray pattern for adequate coverage. Don not apply more than 10 ft. above the crop canopy.
7. For aerial applications, moun spray boom on the aircraft so as to minimize drift caused by wing tip vortices. Use minimum practical boom length and do not use boom that exceeds 75% of wing span or rotor diameter.

Air Assisted (Air Blast) Tree and Vine Sprayers (Ornamentals Only):

Air assisted tree and vine sprayers carry droplets into the canopy of trees and vines via a radially or laterally directed air stream. In addition to the general drift management principles already described, the following specific practices will further reduce the potential for drift:

1. Adjust deflectors and aiming devices so that spray is only directed into the canopy.
2. Block off upward pointed nozzles when there is no overhanging canopy.
3. Use only enough air volume to penetrate the canopy and provide good coverage. Use a minimum of 50 gallons finished spray per acre.
4. Do not allow spray to go beyond the edge of the cultivated area. Spray the outside row only from outside the planting.

TURFGRASS

Crop	Pest	Product Rate	Remarks
Turfgrasses Residential Recreational Commercial	Mole cricket Southern mole cricket Tawny mole cricket	2.7 lbs. per acre (1 oz. per 1000 sq. ft.) 0.54 lbs. a.i. per acre	Make application prior to or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, tank mix with a curative insecticide
	White grub larvae such as: Annual bluegrass weevil Asiatic garden beetle Billbug Black Turfgrass ataenius European chafer Green June beetle Japanese beetle May/June beetle Northern masked chafer Oriental beetle Southern masked chafer	2.7 lbs. per acre (1 oz. per 1000 sq. ft.) 0.54 lbs. a.i. per acre	For optimum control of grubs, billbugs, and annual bluegrass weevil, make application prior to or during egg hatch of the target pest.
	Cutworms Chinchbug Sod webworm	2.7 lbs. per acre (1 oz. per 1000 sq. ft.) 0.54 lbs. a.i. per acre	For suppression of chinchbugs, make application prior to hatching of the first instar nymphs.
	European Cranefly	2.7 lbs. per acre (1 oz. per 1000 sq. ft.) 0.54 lbs. a.i. per acre	

Important Notes:

- Apply in sufficient water to ensure thorough coverage of target area. Use a minimum of 50 gallons finished spray per acre.
- Consult your local State Extension Service or State Extension Turfgrass Specialists for more specific information on timing of insecticide applications.
- For optimal control, apply irrigation if rainfall does not occur within 24 hours after application to ensure movement of the active ingredient through the thatch.
- Do not apply more than a total of 2.7 lbs. of Dinotefuran 20% Turf and Ornamental (0.54 lbs. a.i.) per acre of turf per year.

ORNAMENTAL PLANTS

Foliar or broadcast spray application			
For foliar insect control on ornamental plants in nurseries, greenhouses, interior plantscapes, lath and shadehouses, and outdoor landscapes (commercial, industrial, recreational and residential).			
Crop	Pest	Product Rate	Remarks
Ornamental plants including: Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Not for use on house plants grown inside private residences	Adelgids including: Hemlock Woolly Balsam Woolly Aphids (suppression) Balsam Crepe Myrtle Green Peach Melon Japanese beetles (adults) Lacebugs including: Azalea Cotoneaster Hawthorne Rhododendron Leaf beetles Viburnum Leafhoppers Glassy-Winged Sharpshooter Potato Leafminers including: Serpentine Mealybugs including: Citrus Long-Tailed Madeira Obscure Phormium Pink Hibiscus Psyllids including: Asian Citrus Root Weevils (adults) including: Black Vine Diaprepes Sawflies (larvae) Scales (Armored and Soft) including: Cryptomeria Cycad Aulacaspis Elongate Hemlock Euonymus Florida Red Florida Wax Tea Thrips including Chilli Gynaikothrips uzeli Western Flower (Suppression) Whiteflies including: Giant Greenhouse Silverleaf /Sweetpotato (B and Q Biotypes)	<p style="text-align: center;">Foliar Spray</p> ¼ to ½ lb. per 100 gallons (4 to 8 oz. per 100 gallons) (0.05 to 0.1 lbs. a.i. per 100 gallons) 8-16 oz per Acre (0.1 to 0.2 lbs. a.i./A) 0.2-0.4 oz per 1,000 sq. ft. For treatment of small areas: ½-1.0 tsp per gallon	For optimal control, make first application just before pest populations reach an economic threshold. If necessary, make a second application after 14-21 days. Tank mixing with a surfactant may improve control of pests such as whitefly, mealybug and scale. Confirm plant safety of tank mix in small area before using on a commercial scale. 100 gals. of spray mix will treat 20,000 sq. ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.

1 level teaspoon contains 2.4 grams and 1 cup (8 fl oz) contains 4.0 oz of Dinotefuran 20% Turf and Ornamental.

Do not apply more than 2.7 lbs (0.54 lbs. a.i.) per acre of nursery or landscape per year To delay the development of resistance: Do not apply Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides to consecutive generations of the same insect species without switching to a different mode of action. Do not make more than two sprays of Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides to a single crop during a two-month period. Refer to "Resistance Management "section of label for further guidelines.

VEGETABLE TRANSPLANTS

Foliar or broadcast spray application			
For foliar insect control on vegetable transplants grown in enclosed structures.			
Crop	Pest	Product Rate	Remarks
Cucurbits (Transplants only) Cantaloupe, Cucumber, Melons, Squash Fruiting Vegetables Eggplant, Peppers, Tomato Head and Stem Brassica Broccoli, Brussel Sprouts, Cabbage, Cauliflower, Kohlrabi	Aphids Leafminers Mealybugs Thrips (suppression) Whiteflies including Silverleaf/Sweetpotato (B and Q Biotypes)	3.5 - 7.0 oz per100 gal 7-14 oz per Acre 0.16 - 0.32 oz Per 1,000 sq ft (0.09 to 0.18 lbs. a.i. per Acre)	Do not make more than one application per crop. Apply only to cucurbits and brassica being grown as transplants and before transplants are sold. . 100 gals. of spray mix will treat 20,000 sq. ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.
Leafy Vegetables (Transplants only) (Excluding Brassica spp.)	Aphids Leafminers Mealybugs Thrips (suppression) Whiteflies including Silverleaf/Sweetpotato (B and Q Biotypes)	3.5 - 5.5 oz. per100 gal 7-11 oz per Acre 0.16 - 0.25 oz. Per 1,000 sq ft (0.09 to 0.134 lbs. a.i. per Acre)	Do not make more than one application per crop. Apply only to leafy vegetables being grown as transplants and before transplants are sold 100 gals. of spray mix will treat 20,000 sq. ft of area when using a typical high volume sprayer. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area
1 level teaspoon contains 2.4 grams and 1 cup (8 fl oz) contains 4.0 oz of Dinotefuran 20% Turf and Ornamental. Do not apply more than 1.34 lbs (0.268 lbs. a.i.) per acre of nursery per year. To delay the development of resistance: Do not apply Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides to consecutive generations of the same insect species without switching to a different mode of action. Do not make more than two sprays of Dinotefuran 20% Turf and Ornamental or other Group 4A insecticides to a single crop during a two-month period. Refer to "Resistance Management "section of label for further guidelines.			

PESTS CONTROLLED BY DINOTEFURAN 20% TURF AND ORNAMENTAL WHEN APPLIED TO THE SOIL

<p>Adelgids including: Hemlock Woolly Balsam Woolly Aphids including Balsam Crepe Myrtle Green Peach</p> <p>Melon Bagworms Flatheaded Borers including Alder Bronze Birch Emerald Ash Flatheaded Appletree Two-Lined Chestnut Froghoppers</p> <p>Fungus Gnats (larvae) Horned Oak Gall Lacebugs including:</p> <p>Azalea Cotoneaster Hawthorne Rhododendron</p> <p>Leaf Beetles Elm Viburnum</p> <p>Leafhoppers including Glassy-Winged Sharpshooter Potato</p> <p>Leafminers including: Birch</p> <p>Boxwood Holly Serpentine</p>	<p>Mealybugs Citrus Longtailed Madeira Obscure Phormium Pink Hibiscus</p> <p>Root Pine tip moth (larvae)</p> <p>Plantbugs Psyllids including: Asian Citrus Boxwood Root Weevils (larvae and adults) including Black Vine Diaprepes Roundheaded Borers (excluding Asian Longhorned)</p> <p>Eucalyptus Longhorned Linden Locust Royal Palm Bug Sawfly larvae Scales (Armored and Soft) Including: Azalea Bark Brown Soft Calico Cottony Cushion</p> <p>Cottony Maple Cryptomeria</p>	<p>Scales (Armored and Soft) continued Cycad Aulacaspis Duplacionaspis Elongate Hemlock Euonymus False Oleander Fig Wax Fletcher Florida Red Florida Wax</p> <p>Lecanium Obscure</p> <p>Oystershell Poplar (Aspen) Pine Needle</p> <p>Tea Tuliptree Spittlebugs</p> <p>Thrips including: Chilli (Suppression) Cuban Laurel Gladiolus Gynaikothrips uzeli (Suppression) Western Flower (Suppression)</p> <p>Treehoppers Whiteflies including: Ficus Giant Greenhouse Silverleaf/Sweetpotato (B and Q biotypes)</p> <p>White Grubs including: Oriental Beetle White Pine Weevil</p>
--	--	---

Application to soil				
For systemic insect control on containerized and field grown (in-ground) ornamental plants in nurseries, greenhouses, interior plantscapes, lath and shadehouses, and outdoor landscapes (commercial, industrial, recreational and residential) when applied via soil drench, soil injection, micro-irrigation (spaghetti tube or emitter), drip irrigation, overhead irrigation, ebb and flood irrigation equipment or motorized irrigation equipment.				
Crop	Pest	Product Rate	Remarks	
Ornamental plants including: Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines		Containerized Plants Soil Media Drench ¼ to 1½ pounds per 100 gallons 12 to 24 ounces per 100 gallons 1.5-3.0 teaspoons per gallon	Only apply to moist soil media. Do not apply to dry or saturated media. For optimal performance, do not apply media drench until roots from transplanted plugs or liners have extended at least half way to the edge of pots. Do not leach treated soil media for at least 7 days after application or performance may be reduced. Heavy rainfall or excessive irrigation following application may decrease performance. In general, higher rates will be needed to control insects on woody plants than on herbaceous plants. Poinsettia: For optimal control of whiteflies, treat plants 1-3 weeks after pinch. Late season drenches will take longer to give control.	
		Media Drench Volume for Individual Pots		
		Pot diameter (inches)		Fl oz of dilute solution per pot
		4		2
		5		3
		6		4
		7		5
		8		6
		For larger pot volumes, apply 3 – 4 fl oz of dilute solution (0.11 to 0.22 g product per 4 fl oz,) per gallon of potting media. Use a drench volume that is sufficient to wet soil media without resulting in overflow or runoff through drain holes in pot.		

		<p>Containerized Plants</p> <p>Media Drench Volume for Plants in Raised Beds, Benches, Bedding Flats, Plug and Liner Trays:</p> <p>Apply sufficient dilute solution to wet soil media without loss of liquid from bottom of bed or liner.</p>													
		<p>Containerized Plants</p> <p>Ebb and Flood</p> <table border="1"> <thead> <tr> <th>Pot diameter (inches)</th> <th>Ounces per 1,000 pots</th> </tr> </thead> <tbody> <tr> <td>4</td> <td>1.9 - 3.7</td> </tr> <tr> <td>5</td> <td>2.8 - 5.6</td> </tr> <tr> <td>6</td> <td>3.7 - 7.5</td> </tr> <tr> <td>7</td> <td>4.7 - 9.3</td> </tr> <tr> <td>8</td> <td>5.6 - 11.2</td> </tr> </tbody> </table>	Pot diameter (inches)	Ounces per 1,000 pots	4	1.9 - 3.7	5	2.8 - 5.6	6	3.7 - 7.5	7	4.7 - 9.3	8	5.6 - 11.2	<p>Bring several pots to field capacity, let soil dry and then measure amount of water required to bring pots back to field capacity. Multiply the average volume of water required to rehydrate one pot by the number of pots to be treated. Add this volume of water to the minimum amount of water needed to flood the area to be treated. Re-use any returned volume in subsequent irrigation of same plants.</p> <p>For pot diameter greater than 8", use 3.7-7.5 ounces of Dinotefuran 20% Turf and Ornamental per 1,000 gallons of potting soil media.</p>
Pot diameter (inches)	Ounces per 1,000 pots														
4	1.9 - 3.7														
5	2.8 - 5.6														
6	3.7 - 7.5														
7	4.7 - 9.3														
8	5.6 - 11.2														
		<p>Chemigation of individual containers using a micro-irrigation system (spaghetti tube)</p> <table border="1"> <thead> <tr> <th>Injection ratio</th> <th>Ounces per gallon of injector tank water</th> </tr> </thead> <tbody> <tr> <td>1:100</td> <td>12 - 24</td> </tr> </tbody> </table>	Injection ratio	Ounces per gallon of injector tank water	1:100	12 - 24	<p>Use typical injection ratio for injectors (e.g. 1:100, which equals 1 part injector tank solution: 100 parts irrigation water). Do not mix more than 24 oz of Dinotefuran 20% Turf and Ornamental per gallon of injector tank water, or some product may settle out of solution. Calibrate irrigation system to deliver 3-4 fl oz of dilute solution per gallon of potting media.</p>								
Injection ratio	Ounces per gallon of injector tank water														
1:100	12 - 24														

		<p>Field Grown (In-Ground) Plants</p> <p>Shrubs</p> <p>3 - 6 grams (1.25 - 2.5 level teaspoons) per foot of height</p> <p>1.0 - 2.1 ounces per 10 feet of height</p>	<p>When applied to the soil, Dinotefuran 20% Turf and Ornamental is taken up by actively growing trees and shrubs. Speed of control will be dependent on tree size, tree health, environmental conditions and how actively pests are feeding. In actively growing and transpiring plants, control may be evident within one week after application in small shrubs, and within three weeks after application in large trees. For optimal control, time soil applications so that Dinotefuran 20% Turf and Ornamental has had time to move to insect feeding sites by time target life stage is present.</p> <p>Apply to moist soil. Control may be less effective when applied to dry, saturated, or frozen soil, or at times when plants are not actively taking up water from soil.</p> <p>If possible, irrigate dry soils 1-3 days before application, or apply irrigation within 3 days after application.</p> <p>Heavy rainfall or inadequate irrigation immediately following application may decrease performance.</p> <p>Use higher labeled rates for broadleaf evergreens with dense foliage (ex. hollies), and with very large trees.</p> <p>Soil Drench: Mix required dose in water and uniformly apply to soil around base of shrub or tree. Pull back mulch before drenching. For optimal performance, apply 1-4 pints of drench solution per foot of height (shrubs) or inch of trunk diameter (trees). Adjust drench volume based on soil type, soil moisture and thickness of mulch so that product is moved into root zone. To enhance soil penetration in heavy soils and sloping terrain, dig shallow holes around tree or shrub, and apply drench solution in holes. Lower drench volumes may be less effective in dry soils or when applied over heavy mulch unless there is adequate rainfall or irrigation after application to move product into root zone.</p>
--	--	--	--

		<p>Small Trees (Less than 24" diameter at breast height)</p> <p>3 - 12 grams (1.25 - 5.0 level teaspoons) per inch of trunk diameter at breast height (DBH)</p> <p>1.05 - 4.2 ounces per 10 inches of trunk diameter at breast height (DBH)</p> <p>For multi-stem trees, base rate on cumulative inches of diameter of all stems at breast height.</p>	<p>Soil Injection: Mix required dose in water and make at least four injections per shrub or tree with a low-pressure applicator. Use same amount of solution per hole. Injections can be made using the following methods:</p> <p><i>Grid System</i>-Space injections on a 2.5 ft center extending to drip line.</p> <p><i>Circle System</i>-Make injections in concentric circles extending inward from drip line.</p> <p><i>Basal System</i>- Space injections evenly around trunk no more than 24" out from the base.</p>														
		<p>Large Trees (24" diameter or greater at breast height)</p> <p>6 - 12 grams (2.5 - 5.0 level teaspoons) per inch of trunk diameter at breast height (DBH)</p> <p>2.1 - 4.2 ounces per 10 inches of trunk diameter at breast height (DBH)</p> <p>For multi-stem trees, base rate on cumulative inches of diameter of all stems at breast height.</p>	<p>Dinotefuran 20% Turf and Ornamental may be soil injected with low volume (e.g. Kioritz injector) or high volume injection equipment. For optimal performance, inject 1 - 32 fl oz of dilute solution per foot of height or inch of trunk diameter depending on application equipment. Make shallow injections where feeder roots are most concentrated.</p>														
		<p>Banded spray application to soil surface</p> <p>(2.7 lbs per acre)</p> <table border="1" data-bbox="795 1071 1104 1449"> <thead> <tr> <th>Row spacing in feet</th> <th>Ounces per 1,000 linear feet of row</th> </tr> </thead> <tbody> <tr><td>3</td><td>3</td></tr> <tr><td>4</td><td>4</td></tr> <tr><td>5</td><td>5</td></tr> <tr><td>6</td><td>6</td></tr> <tr><td>7</td><td>7</td></tr> <tr><td>8</td><td>8</td></tr> </tbody> </table>	Row spacing in feet	Ounces per 1,000 linear feet of row	3	3	4	4	5	5	6	6	7	7	8	8	<p>Apply as a uniform band in row over root zone. Apply from peak adult flight to peak egg hatch.</p> <p>Apply in at least two gallons of water per 1,000 linear feet. Irrigate after application to move product into soil profile.</p> <p>Control any weeds in treated area prior to application, or performance may be reduced.</p> <p>Adjust rates accordingly for other row spacing.</p> <p>For optimal performance, irrigate after application to move Dinotefuran 20% Turf and Ornamental into the root zone.</p>
Row spacing in feet	Ounces per 1,000 linear feet of row																
3	3																
4	4																
5	5																
6	6																
7	7																
8	8																
		<p>Broadcast spray to soil of plant beds</p> <p>2.7 lbs per acre</p>	<p>Apply over the top of ornamental plant beds in a water volume sufficient to move product to soil surface. If necessary, irrigate after application to move product off of foliage and into upper root zone of soil. May be less effective on large woody shrubs than on herbaceous annuals and perennials.</p>														

1 level teaspoon contains 2.4 grams, and 1 cup (8 fl oz) contains 4.0 oz of Dinotefuran 20% Turf and Ornamental.

Do not apply more than 2.7 lbs (0.54 lbs. a.i.) per acre of nursery or landscape per year.

To delay the development of resistance in greenhouses, nurseries and interiorscapes, do not make more than one soil application per crop cycle or three-months, whichever is shorter. Refer to "Resistance Management" section of the label for additional guidelines.

Trunk sprays in trees and large shrubs

For systemic insect control in containerized and field grown (in-ground) ornamental trees and shrubs in nurseries, interior plantscapes, lath and shadehouses and outdoor landscapes (commercial, industrial, recreational and residential) when applied as a trunk spray.

Crop	Pest	Product Rate	Remarks
Shrubs Ornamental trees Non-Bearing Fruit Trees Non-Bearing Nut Trees	Adelgids including Hemlock Woolly Aphids Flatheaded Borers including Alder Bronze Birch Emerald Ash Flatheaded Appletree Two-lined Chestnut Lacebugs Leaf Beetles Leafhoppers Leafminers Mealybugs Pine Tip Moth (Larvae) Psyllids Roundheaded Borers (Excluding Asian Longhorned) Scales including Calico Thrips (Suppression) Whiteflies Ficus	12 – 24 oz per gallon Depending on bark type and thickness, one gallon of spray solution will cover 40-50" of trunk diameter when measured at 4.5 feet above ground.	<p>When sprayed on the trunk, Dinotefuran 20% Turf and Ornamental will be absorbed through the bark and into the vascular system, and then transported throughout the tree. Speed of control will be dependent on tree size, tree health, environmental conditions and how actively pests are feeding. In actively transpiring trees, control may be evident within 1-4 weeks after application.</p> <p>Spray bark on root flare (buttress roots) and on trunk between soil surface and 4-5 feet above the soil surface.</p> <p>Adjust nozzle to uniformly distribute spray over the entire circumference of the tree trunk and buttress roots. Wet bark just to the point of saturation and run off onto soil.</p> <p>Apply with a low volume sprayer operated at 10-20 PSI to prevent tree damage, bounce back and drift.</p> <p>For optimal control, apply to actively growing trees, and time application so that Dinotefuran 20% Turf and Ornamental has had time to move to insect feeding sites at when target life stage is present.</p> <p>Do not apply to wet bark, during rainfall or if rain is expected within 12 hours.</p> <p>Control may be less effective in trees with thick bark, and at times when trees are not actively growing or transpiring.</p>

1 level teaspoon contains 2.4 grams, and 1 cup (8 fl oz) contains 4.0 oz of Dinotefuran 20% Turf and Ornamental

Do not apply more than 2.7 lbs (0.54 lbs. a.i.) per acre of nursery or landscape per year.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Pesticide Storage: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in a cool dry place. Do not store diluted spray. For help with any spill, leak, fire or exposure involving this material, call day or night 1-800-CLEANUP.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank.

Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE

MITSUI CHEMICALS, INC., warrants that this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions to the crops specified. To the extent consistent with applicable law, there are no other warranties, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, this warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or conditions not reasonably foreseeable to seller, and buyer assumes all risk of any such use.

021010