

#### U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) **Ariel Rios Building** 1200 Pennsylvania Ave., NW

Washington, D.C. 20460

EPA Reg. Number:

Date of Issuance:

33658-10

DEC 3 1 2007

NOTICE OF PESTICIDE:

Registration Reregistration

(under FIFRA, as amended)

Term of Issuance: Unconditional

Name of Pesticide Product:

Pilot 15G Chlorpyrifos Agricultural Insecticide

Name and Address of Registrant (include ZIP Code):

Gharda Chemicals, Ltd. 660 Newtown-Yardley Road, Suite 105 Newtown, PA 18940

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number. On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the Reregistration Eligibility Decision, EPA has reregistered the product listed above. This action is taken under the authority of section 4(g)(2)(C) of the Federal Insecticide. Fungicide, and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.

Generic data requirements that have been identified in the RED as confirmatory data for the active ingredient contained in the subject product will continue to be tracked and assessed in the Agency's continual reassessment of pesticides.

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable with the following comments:

#### Please see page 2 for label comments

You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this Notice or the next printing of the label whichever occurs first, must bear the new revised label. A stamped copy of the label is enclosed for your records. Should you have any questions, please contact Registration Review Manager, Julie Chao, by phone at: 703-308-8735 or by email at: chao.julie@epa.gov.

Signature of Approving Official:

Meredith F. Laws, Chief

Insecticide-Rodenticide Branch

Registration Division (7505P)

Date:

DEC 3 1 2007

EPA Form 8570-6

#### **Label Comments:**

- NOTE: If the product is packaged in a container weighing less than 50 pounds, then it must be labeled as a "Restricted Use Pesticide."
- On page 1, delete the phrase "Low Odor Formula."
- On page 1, delete the phrase "various" and replace it with "listed" so the sentence reads, "For control of listed insects infesting certain field and vegetable crops."
- On page 1, under the First Aid section, reorder the statements as follows:
  - IF SWALLOWED:...
  - IF IN EYES:...
  - IF ON SKIN OR CLOTHING:...
  - IF INHALED:...
- On page 2, under Hazards to Humans and Domestic Animals, modify the eye irritation statement so it reads, "Causes **moderate** eye irritation."
- On page 2, under the PPE section, modify the first paragraph so it reads, "Some materials that are chemical-resistant to this product are **barrier laminate or viton**. If you want more instructions, follow the instructions for category **H** on an EPA chemical resistance category selection chart."
- On page 2, under the PPE requirements, "chemical-resistant apron" and "chemical-resistant headgear" are not required and may be removed.
- On page 2, the final sentence under the PPE section should be revised as follows: "A NIOSH-approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any N, R, P, or HE filter."
- On page 2, under User Safety Recommendations, modify the second statement so it reads, "Remove clothing and/or PPE immediately if pesticide gets inside..."
- On page 2, modify the fifth sentence of the Environmental Hazards section so it reads, "Do not contaminate water when cleaning equipment or disposing of equipment wash water or rinsate."
- On page 2, in the Agricultural Use Requirements box, delete the second paragraph and replace it with the following text:
  - "Do not enter or allow entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the directions for use associated with each crop."
- On page 2, under Storage and Disposal, in the Pesticide Disposal paragraph, delete the phrase "Pesticide wastes are toxic."

- For consistency and clarity, list the restrictions for each crop as bullets (as with the corn directions).
- For consistency and clarity, list all application rates and pests controlled in a table (as with the directions for corn, sorghum, peanuts, soybeans, and sugar beets).
- For consistency and clarity, list crops alphabetically.
- On page 3, for Corn, under the Pest Specific Use Directions, delete fire ants. This pest/use pattern requires additional efficacy data.
- On page 4, for Onions (Dry Bulb), modify the last sentence of the At Plant In-Furrow Treatment Paragraph so it reads, "Do not exceed 6.6 lb. Pilot 15G per acre (1 lb. ai chlorpyrifos)." The rate 6.7 lb. Pilot 15G exceeds the use restriction of 1 lb ai per acre.
- On page 4, for Onions (Dry Bulb), add the following statements under the Restrictions section:
  - Restricted Entry Interval (REI) is 24 hours.
  - Do not apply last treatment within 7 days before harvest.
  - Do not apply more than 1 lb. ai chlorpyrifos per crop per season.
  - Do not make more than 1 application of any product containing chlorpyrifos per year.
- On page 4, for Citrus Grove Floors, delete fire ants. The claim "other ant species" is overly broad, and also unacceptable. You may claim ants, but only if you make a note excluding broadcast spray treatment for fire, harvester, carpenter, and pharaoh ants. These pests/use patterns require additional efficacy data.
- On page 4, for Citrus Grove Floors, the application rate must be limited to 6.6 lb Pilot 15G per acre." The rate 6.7 lb. Pilot 15G exceeds the use restriction of 1 lb ai per acre.
- On page 4, for Citrus Grove Floors, under the Postplant Broadcast Treatment paragraph, delete the sentence "Retreat as needed." This sentence contradicts the required use restrictions.
- On page 4, for Citrus Grove Floors, modify the first sentence of the Restrictions section so it reads, "Do not apply more than 20 lbs. of Pilot 15G per year (3 lb. ai per acre per season)."
- On page 4, for Citrus Grove Floors, add the following statements under the Restrictions section:
  - Restricted Entry Interval (REI) is 5 days.
  - Do not make a second application within 10 days of any application of chlorpyrifos to the orchard.
  - Do not apply more than 1 lb. ai chlorpyrifos per application.

- On page 4, for Sorghum/Milo, delete fire ants from the Application Rates and Pests Controlled table. This pest/use pattern requires additional efficacy data.
- On page 4, for Sorghum/Milo, the application rate (oz per 1000 ft of row) listed in the table for lesser cornstalk borer must be changed to "4 to 8."
- On page 4, for Sorghum/Milo, add the following statements under the Restrictions section:
  - Restricted Entry Interval (REI) is 24 hours.
  - Do not make a foliar application of any other product containing chlorpyrifos within 10 days of an at-plant application of Chlorpyrifos 15G.
  - Do not apply last treatment within 7 days before harvest.
- On page 5, for Peanuts, remove the Aerial Broadcast column in the Application Rates and Pests Controlled table. Aerial application is prohibited. Also delete all subsequent references or directions related to aerial application rates (i.e., delete the Broadcast Rescue Treatment paragraph and the phrases "or 13.3 pounds of Pilot 15G per acre per application" and "or 26.6 pounds per acre per crop season" under the Restrictions section).
- On page 5, for Peanuts, add the following statements under the Restrictions section:
  - Restricted Entry Interval (REI) is 24 hours.
  - Do not make more than 2 applications of Pilot 15G per year.
  - Do not make a second application of Pilot 15G or any other product containing chlorpyrifos within 10 days of the first application.
- On page 5, for Soybeans, the reference to the "accompanying table" should be clarified (i.e., refer to Table 1 by name).
- On page 5, for Soybeans, add the following statements under the Restrictions section:
  - Restricted Entry Interval (REI) is 24 hours.
  - Do not apply within 28 days of harvest.
  - The maximum single application rate is 2 lb ai chlorpyrifos per acre for preplant/at-plant incorporation and 1 lb ai chlorpyrifos per acre for foliar and postharvest application.
  - Do not make more than 2 applications of other products containing chlorpyrifos
  - Do not make a foliar application of any other product containing chlorpyrifos within 10 days of an at-plant application of Chlorpyrifos 15G.
- On page 5, for Sugar Beets, in the At Plant Band Treatment section, change all occurrences of "6.75 lb." to "6.7 lb." for consistency with the value provided in Table 1.
- On page 5, for Sugar Beets, in the Postemergence Band Treatment paragraph, change "10 to 13.5 lb." to "9.7 to 13.4 lb." for consistency with the values provided in Table 1.

- On page 5, for Sugar Beets, add the following statements under the Restrictions section:
  - Restricted Entry Interval (REI) is 24 hours.
  - Do not apply within 30 days of harvest.
  - Do not apply more than a total of 3 lb. ai chlorpyrifos per acre per year, or make more than 3 applications of products containing chlorpyrifos per season.
  - Do not make a foliar application of any other product containing chlorpyrifos within 10 days of an at-plant application of Chlorpyrifos 15G.
- On page 6, for Sunflowers, add the following statements under the Restrictions section:
  - Restricted Entry Interval (REI) is 24 hours.
  - Do not apply within 30 days of harvest.
  - The maximum single application rate is 2 lb ai chlorpyrifos per acre for preplant/at-plant incorporation and 1 lb ai chlorpyrifos per acre for foliar and postharvest application.
  - Do not make more than 2 applications of other products containing chlorpyrifos
  - Do not make a foliar application of any other product containing chlorpyrifos within 10 days of an at-plant application of Chlorpyrifos 15G.
- On page 6, for Sweet Potatoes, add the following statements under the Restrictions section:
  - Restricted Entry Interval (REI) is 24 hours.
  - The maximum single application rate is 2 lb ai chlorpyrifos per acre.
- On page 6, for Tobacco, delete "to 20" from the application rate instructions. The maximum rate is 13.5 lb per acre.
- On page 6, for Tobacco, add the following statements under the Restrictions section:
  - Restricted Entry Interval (REI) is 24 hours.
  - Do not apply within 7 days before harvest.
  - The maximum single application rate is 2.025 lb. ai chlorpyrifos per acre.
- On page 6, for Brassica (Cole) Crops, add the following statements under the Restrictions section:
  - Restricted Entry Interval (REI) is 24 hours, except for cauliflower. The REI for cauliflower is 3 days.
  - Do not apply within 30 days of harvest.
  - The maximum single application rate is 1.4 oz ai chlorpyrifos per 1000 ft of row, except for cauliflower. For cauliflower, the maximum application rate is 1.2 oz ai/1,000 ft. of row.
  - Do not make a foliar application of any other product containing chlorpyrifos within 10 days of an at-plant application of Chlorpyrifos 15G.

- On page 6, for Brassica (Cole) Crops, Rutabagas must be listed separately. Therefore, delete all references to rutabaga in the Brassica (Cole) Crops section (i.e., remove "Do not use rutabaga tops of food or feed purposes").
- In the Rutabaga section, add the following statements:
  - Restricted Entry Interval (REI) is 24 hours.
  - Do not apply within 7 days before harvest.
  - Application rate is 10.56 oz Chlorpyrifos 15G per 1000 ft of row.
  - The maximum single application rate is 1.6 oz ai chlorpyrifos per 1000 ft of row (8.8 lb ai chlorpyrifos per acre).
  - Do not make more than one application per crop season.
  - Do not use rutabaga tops of food or feed purposes.
- On page 6, for Radishes, add the following statements under the Restrictions section:
  - Restricted Entry Interval (REI) is 24 hours.
  - Do not apply within 7 days before harvest.
  - The maximum single application rate is 0.5 oz ai chlorpyrifos per 1000 ft of row (2.75 lb ai chlorpyrifos per acre).
- For consistency and clarity, on page 7, for Table 1, include all values for 18" row spacing and extend the table to 16.0 oz (as on the Lorsban 15G label).
- For consistency and clarity, on page 8, for Table 2, include values for the application rate in oz per 1000 ft of row (as on the Lorsban 15G label).
- On page 8, under the Notice of Warranty and Disclaimer, fix the typo in the second to last sentence so that "ll" reads "all."

# PILOT™ 15G

### Chlorpyrifos Agricultural Insecticide

#### LOW ODOR FORMULA

For control of various insects infesting certain field and vegetable crops.

Active Ingredient:

Chlorovrifos: O.O-diethyl O-(3,5,6-trichloro-2-pyridinyl) phosphorothioate ......15.0% Inert Ingredients: ......85.0%

#### KEEP OUT OF REACH OF CHILDREN **PRECAUCION** CAUTION

Si usted no entiende la etiqueta, busque a aiguien para que se la explique a usted en detaile. (If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. No.: 33658-10 EPA Est. No.: 39578-TX-1

Net Contents: 50 pounds

Pilot is a registered trademark of Gharda Chemicals Limited Newtown, PA 18940

## Gharda Chemicals Limited

FIRST AID (Organophosphate Insecticide)						
If inhaled:	<ul> <li>Remove person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>Call a poison control center or doctor for further treatment advise.</li> </ul>					
If on skin or clothing:	<ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advise.</li> </ul>					
If in eyes:	<ul> <li>Hold eye open and rinse slowly and gently with water for15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advise.</li> </ul>					
If swallowed:	<ul> <li>Call poison control center or doctor immediately for treatment advise.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by</li> </ul>					
	the poison control center or doctor.  Do not give anything by mouth to an unconscious person.					

#### HOT LINE NUMBER (Organophosphate Insecticide)

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical treatment information call: 1-(866)-359-5660

#### NOTE TO PHYSICIAN

Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate significance of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidote immediately after establishing an open airway and respiration.

## with COMMENTS In EPA Letter Dated:

DEC 3 1 2007 Under the Federal Record Fungicide, and Rodentiettle A as amended, for the pesti registered under EPA Reg. No.

O entact w 2 14

#### PRECAUTIONARY STATEMENTS

#### **Hazards To Humans And Domestic Animals**

**CAUTION.** Harmful if swallowed. Causes eye imitation. Avoid contact with eyes, skin or clothing. Avoid breathing dust.

Personal Protective Equipment

Some materials that are chemical-resistant to this product are natural rubber and polyethylene. If you want more options, follow the instructions for category A on an EPA chemical resistance category selections chart.

#### All mixers, loaders, other applicators and other handlers must wear:

- · coveralls over long-sleeved shirt and long pants;
- · chemical-resistant gloves;
- chemical-resistant apron when mixing or loading or exposed to the concentrate;
- · chemical resistant footwear plus socks:
- · chemical-resistant headgear for overhead exposures;
- a NIOSH-approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator any R,P or HE filter.

**User Safety Requirements** 

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry

#### **User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside.
   Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### **Engineering Controls**

Pilots must use an enclosed cockpit in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

When applicators use closed cab equipment in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### **Environmental Hazards**

This pesticide is toxic to fish, aquatic invertebrates, small mammals and birds. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water when disposing of equipment wash water. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

This product is not registered in California and Arizona. California and Arizona law prohibits sale, distribution, and use within the State of any products not registered by the State.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use before applying.

Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. Do not apply by aircraft at a rate greater than 6.5 pounds of formulated product (1 pound of active ingredient) per acre. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the required restricted entry interval (REI). The required REI's are:

For all crops: 24 hours, unless specifically otherwise noted below

- Fruit trees: 4 days
- Citrus trees: 5 days
- Cauliflower: 3 days

Also see specific Use Directions under **Approved Crops** Section of this label

**Exception:** If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- · coveralls over short-sleeved shirt and short pants;
- · chemical-resistant gloves made out of water proof material;
- · chemical-resistant footwear plus socks;
- · chemical-resistant headgear for overhead exposure.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a secured dry storage area. Prevent cross contamination with other pesticides and fertilizers. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as indicated below.

In Case of Spill: Isolate the spill. Hold this package, other cargo and vehicles involved. For Emergency spill assistance Call CHEMTREC (24-hour service): 1-800-535-5053.

Pesticide Disposal: Open dumping is prohibited. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Offer for recycling if available, or, dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

#### **APPROVED USES**

#### Field Corn, Sweet Corn, and Corn Grown for Seed \*

**Application Rates and Pests Controlled** 

	Bar (Ou	Aerial Broadcast Application (lb/Acre)		
Pests Controlled				
	. At Plant	Applications	Postplant Treatment	
	T-Band	In-Furrow		
armyworms	-		6-8	-
billbugs	8	-	-	-
Chinch bug (1)	8	8	-	-
Cutworms (1)	8	8	-	-
European and southwestern corn borer(3)	·		,	
1st Generation 2nd Generation	-	· <del>-</del>	3.5 to 8 6 to 8	5.0 to 6.5 6.5
grubs	8	8		-
lesser cornstalk borer	8	· •		· -
Northern, Western and Southern corn rootworm larvae	8	8	8	-
seed corn beetle	8	8	-	-
seed corn maggots	8	8	-	
Southern corn Rootworm larvae	. 8	8	8	-
symphylans	8	-	-	
Wireworms (2)	. 8	8 '	-	-

Numbers in parentheses (-) refer to Pest-Specific Use Directions.

**NOTE:** Pilot 15G insecticide is compatible with all ALS inhibitor herbicides, including Accent and Beacon herbicides, applied in accordance with label recommendations. Refer to product label for additional Precautionary Statements, Mixing and Application instructions.

#### **Pest Specific Use Directions:**

- 1. Cutworms, fire ants and chinch bugs: The 8 oz rate provides suppression only for in-furrow treatments.
- 2. **Wireworms:** For best control, apply as an in-furrow treatment. Consider using a hopper box insecticidal seed treatment with T-band applications.
- 3. European corn borer: When using post plant banded applications, use rates of 3.5 to 4 oz of Pilot 15G per 1000 feet of row for low to moderate first generation infestations before larvae have entered corn stalks. Use application rates of 6 to 8 oz of Pilot 15G per 1000 feet of row for severe first generation infestations and all second generation infestations before larvae have entered corn stalks.

At Plant T-Band Application: Apply 8 oz of Pilot 15G per 1000 ft of row as a T-band over an open seed furrow over the row behind the planter shoe, in front of the press wheel. In conventional and minimum-till corn, incorporate into the top ½ to 1 inch of soil using suitable equipment. A soil applied T-band treatment may be followed by post-applied herbicides. Pilot 15G has demonstrated suppression of certain soil-borne pathogens that may result in physiological and agronomic advantages to corn under environmental stress conditions when compared to corn not treated with Pilot 15G.

At Plant In-Furrow Application: Apply 8 oz of Pilot 15G per 1000 ft of row at planting as an in-furrow treatment in conventional, minimum and no-till corn. Direct the granules into the planter shoe with the seed, or place the applicator tube directly behind the planter shoe so that the granules drop into the seed furrow, or place the granular band applicator behind the planter shoe so that the granules fall on the soil surface and into the open seed furrow and are covered with soil.

**Postplant Application:** To control corn rootworm larvae, apply 8 oz of Pilot 15G per 1000 ft of row at cultivation by placing the granules at the base of the plant on both sides of the row just ahead of the cultivation shovels and covering the granules with soil. To control European and southwestern corn borer larvae, apply Pilot 15G in a band over the row so that the granules are directed into the whorl or use a postplant broadcast treatment. Consult your state agricultural experiment station or extension service specialist for proper time to treat and local threshold information. Scouting for insect damage is strongly encouraged.

Postplant Broadcast Treatment: To control European and southwestern corn borers, apply Pilot 15G by uniformly broadcasting the granules over the corn plants by aerial application or by applying the granules into the corn whorls by ground application. For aerial applications, do not apply within 150 feet of rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries and commercial fishponds. Apply at a rate of 5 lb per acre for low to moderate first generation infestations or at 6.5 lb per acre for severe first generation infestations and all second generation infestations. Apply before larvae have entered corn stalks. Consult your state agricultural experiment station or extension service specialist for local threshold information. Scouting for insect damage is strongly encouraged.

#### **Specific Use Restrictions:**

- Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.
- Preharvest Interval: Do not apply within 21 days before harvest of grain or ears.
- Do not apply by aircraft at a rate greater than 1 lb ai per acre.
- Do not make more than 1 at-plant application and 1 foliar application of Pilot 15G per season at the 1 lb ai chlorpyrifos rate.
- Do not make more than 3 applications of any product containing chlorpyrifos per season, including the maximum allowed of 2 granular applications, at the 1 lb ai chlorpyrifos rate. Re-treatment with a second soil application of Pilot 15G is allowed under replant situations due to loss of crop during establishment only when initially applied at the rate of 1 lb.

- Do not apply more than a total of 3 lb ai chlorpyrifos per acre per season.
- Do not make a second application of Pilot 15G or other product containing chlorpyrifos within 10 days of the first application.
- Maximum single application rate for at-plant applications is 8 oz of Pilot 15G per 1000 ft of row (1.3 lb ai chlorpyrifos per acre).
- Maximum single application rate for postplant applications is 6.5 lb of Pilot 15G (1 lb ai chlorpyrifos) per acre.
- If more than 1 lb ai granular chlorpyrifos per acre is applied at-plant (for a maximum of 1.3 lb ai per acre per season), only 1 additional application of a liquid product containing chlorpyrifos at 1 lb ai per acre is allowed per season, for a total of 2.3 lb ai chlorpyrifos per acre per season

#### Onions (Dry Bulb)

Pest Controlled: onion maggot

Application Rate: 3.7 ounces per 1,000 feet of row at an 18-inch row spacing

At Plant In-Furrow Treatment: Apply as an at planting in-furrow treatment. In Colorado, Idaho, Washington, and Oregon, to control onion maggots in onions planted in double rows with rows spaced 2 to 4 inches apart, apply Pilot 15G at the rate of 3.7 oz per 1,000 feet of double row. Place the granules in a 5 to 7 inch wide band over both rows behind the planter shoe and in front of the press wheel to achieve shallow incorporation. Do not exceed 6.7 lb per acre.

Restrictions: Do not make more than one application per year.

#### **Citrus Grove Floors**

Pests Controlled: fire ants and other ant species

Application Rate: 6.7 lb/acre

Postplant Broadcast Treatment: To control foraging ants and suppress mounds, apply Pilot 15G with ground application equipment. Use a suitable granular applicator, such as a cyclone fertilizer spreader, that will uniformly broadcast the granules over the grove floor. Retreat as needed. Pilot 15G may be custom blended with granular fertilizers provided that application of the blended Pilot 15G plus fertilizer mixture can be applied uniformly to the grove floor. Do not apply where weed growth or other obstructions would impede uniform coverage of the grove floor.

**Restrictions:** Do not apply more than 3 applications per season. Do not apply more than 20 pounds of Pilot 15G per acre per crop season. Do not apply last treatment within 28 days before harvest. Do not allow livestock to graze in treated areas.

#### Sorghum/Milo

Application Rates and Pests Controlled

	Banded Applications (Ounces per 1,000 feet of row)			
	At Plant Treatments			
Pests Controlled	T-Band Band			
lesser cornstalk borer	4 to 12	4 to 12		
corn rootworm and		·		
cutworms	8	8		
fire ants*	8	-		
chinch bug*	8	-		

<sup>\* 8</sup> oz. rate - suppression only

At Plant T-Band or Band Treatments: Apply in a 6 to 8 inch band over the row and incorporate into the top 1-inch of soil using suitable equipment. Equivalent rates of Pilot 15G per acre for various row spacing is given in Table 1. Use the lowest rate for lesser cornstalk borer control when protection is desired for 2 to 3 weeks and higher rates for longer residual activity. It is absolutely necessary to incorporate the granules, especially at lower rates.

Restrictions: Do not make more than one application per season.

#### **Peanuts**

Use Pilot 15G to control cutworms, lesser comstalk borer, southern com rootworm larvae, suppress wireworms, and inhibit the growth and development of white mold (southern blight) disease caused by *Sclerotium rolfsii*. Pilot 15G will control only those cutworms existing in the soil from the time of application up to 30 days following application.

Application Rates and Pests Controlled

	Banded Applications (Ounces/1,000 feet of row)				
Pests Controlled	At Plant Treatment	Postplant Treatment	Aerial Broadcast (lb/Acre)		
*Preventative Treatments: cutworms lesser cornstalk borer southern com rootworm larvae wireworms white mold (Southern blight)	7.5 to 15	7.5 to 15			
**Rescue Treatments: lesser cornstalk borer	-	7.5 to 15	10 to 13		
Burrowing bug	-	-	13.3		

\*At Plant Preventive Treatment: Apply Pilot 15G in a 6 to 12 inch band over the row behind the planter shoe and in front of the press wheel. Incorporate granules to a depth of 1-inch with tines or chains or other suitable equipment. If the 7.5 oz rate is used at planting time, then another application of 7.5 oz per 1,000 feet of row should be made postplant to extend control.

\*Postplant Preventative Treatment: Apply Pilot 15G to peanuts at early flowering to pegging stage of growth in a 6 to 8 inch band over the row. For extended insect control and continued suppression of white mold (southern blight), a second application of Pilot 15G may be made. Best suppression of white mold (southern blight) is obtained by applying the maximum rate of 15 oz per 1,000 ft of row for each postplant treatment. Irrigation or rain following application is needed to enhance treatment effectiveness for suppression of white mold. Under conditions of heavy white mold pressure, a suitable fungicide may also be required and must be applied separately.

\*\*Band Rescue Treatment: Use Pilot 15G for the control of lesser comstalk borer when the insect first appears, usually just prior to or at pegging. Apply in a 10 to 18 inch band over the fruiting zone.

\*\*Broadcast Rescue Treatment: Use Pilot 15G for the control of lesser cornstalk borer when the insect first appears, usually just prior to or at pegging. Uniformly apply at a rate of 10 to 13 pounds per acre by aerial broadcast application. For control of burrowing bugs, use Pilot 15G when adults first appear, usually at mid to late pegging. Uniformly apply at a rate of 13.3 pounds per acre by aerial broadcast application. Care must be taken to ensure proper equipment calibration and delivery.

Restrictions: Do not apply more than 15 ounces of Pilot 15G per application per 1,000 feet of row or 13.3 pounds of Pilot 15G per acre per application. Do not apply more than 30 oz per 1,000 ft of row or 26.6 pounds per acre per crop season. Do not harvest within 21 days after treatment. Do not feed peanut forage or hay to meat or diary animals. The combined total of preplant and postplant applications of Pilot 4E and Pilot 15G must not exceed 4 pounds of active ingredient per acre per crop season. Aerial application of Pilot 15G to peanuts is prohibited.

#### Soybeans

Application Rates and Pests Controlled

Application rates an	u i esis conincilei	<u> </u>			
	Banded Applications				
	(Ounces per 1,000 feet of row)				
Pests Controlled	T-Band	Band	Postplant Treatment		
lesser cornstalk borer cutworms	. 8	8	8		

At Plant and Postplant Treatments: Use Pilot 15G insecticide to control larvae of the lesser comstalk borer and cutworms by application at planting time or postemergence as a band (row) treatment at the rate of 8 oz per 1,000 feet of row. In the southeast apply 4 to 8 oz per 1,000 feet of row as an at-plant treatment. Equivalent rates of Pilot 15G per acre for various row spacing are given in the accompanying table. When applied at planting time incorporate the granules into the top 1 inch of soil by placing in a 4 to 10 inch band over the row behind the planter shoe and ahead of the press wheel. A drag chain can also be used for incorporation. For postemergence treatment when insects first appear incorporate the granules in a 4 to 10 inch band to a depth of 1/2 to 1 inch using a suitable cultivator. Apply Pilot 15G with equipment that will provide uniform distribution of the granules. Do not apply as an in-furrow treatment. For suppression of fire ants, use Pilot 15G at 8 oz per 1,000 feet of row as an at-plant T-band treatment.

Restrictions: Do not make more than one application per season.

#### **Sugar Beets**

Granular insecticides, including Pilot 15G, may contribute to the stress of the sugar beet plant under certain environmental conditions. This stress may reduce plant stand or interfere with normal plant development. Herbicides used preplant incorporated may interact with insecticides and enhance this stress.

Application Rates and Pests Controlled

	Banded Applications (Ounces per 1,000 feet of row)					
	At Plant Treatments					
Pests Controlled	d T-Band Band					
Sugar beet root maggot	-	4.5 to 9.0	6.5 to 9.0			
cutworms	-	6.6 to 9.0	-			
wireworms (suppression)	1 -	6.5 to 9.0	-			

At Plant Band Treatment: To control sugar beet root maggot larvae and cutworms at planting time, place Pilot 15G in a band 4 to 5 inches wide behind the planter shoe, over the drill row, and in front of the press wheel. Do not apply granules in direct contact with seeds. Apply Pilot 15G at the rate of 4.5 to 9 ounces per 1,000 feet of row (equivalent to 6.75 to 13.5 lb per acre based on a 22 inch row spacing). When root maggot populations are expected to be low, apply Pilot 15G at a rate of 4.5 ounces per 1,000 feet of row (equivalent to 6.75 lb per acre based on 22 inch row spacing). If initial adult fly activity indicates higher than anticipated populations, apply Pilot 4E at or near the time of peak adult emergence to augment control. (Review label for Pilot 4E for recommended use rates, application timing, methods of application, and insecticide resistance management). Incorporate Pilot 15G into the top 1/2 to 1 inch of soil using suitable equipment.

**Postemergence Band Treatment:** For postemergence control of sugar beet maggot larvae, place Pilot 15G in a band 3 to 5 inches wide over the beet row (up to 2 to 4 true leaf stage of plant growth). Apply Pilot 15G at the rate of 6.5 to 9 oz per 1,000 feet of row (equivalent to 10 to 13.5 lb per acre based on a 22 inch row spacing). Incorporate Pilot 15G into the top 1/2 to 1 inch of soil using a suitable incorporation device.

Restrictions: Do not make more than one application per year.

#### **Sunflowers**

Pest Controlled: cutworms

Application Rate: 8 oz per 1,000 feet of row

At Plant Band Treatment: Place the granules in a 7 inch wide band over the row behind the planter shoe in front of the press wheel and incorporate into the top 1 inch of soil using suitable equipment.

Restrictions: Do not make more than one application per season.

#### **Sweet Potatoes**

Pests Controlled: wireworms (conoderus), flea beetles (Systena), and sweet potato beetle

Application Rate: 13.5 lb/acre

Preplant Broadcast Treatment: Use Pilot 15G to reduce the feeding damage caused by populations of the listed pests. Evenly broadcast the granules over the soil surface and then incorporate the granules into the soil to a depth of 4 to 6 inches using a rotary hoe, disc cultivator, or other suitable equipment. Plant the crop in the usual manner no later than 14 days after treatment (any delay in planting will reduce the length of time that Pilot 15G will protect against feeding damage). Pilot 15G will not control false wireworm or whitefringed beetle and other grubs that attack sweet potatoes.

**Restrictions:** Do not make more than one application per season. Do not harvest within 125 days of treatment.

#### Tobacco

Pests Controlled: cutworms, flea beetle larvae, root maggots, and wireworms

Application Rate: 13.5 to 20.0 lb/acre

**Preplant Broadcast Treatment:** Apply Pilot 15G one week before transplanting, using equipment that will evenly distribute the granules over a treated area. Immediately following application, incorporate the granules into the soil to a depth of 2 to 4 inches using suitable equipment. The application of Pilot 15G will also suppress movement of imported fire ant into treated field.

Restrictions: Do not make more than one application per season.

#### Brassica (Cole) Crops

(Bok Choy, Broccoli, Broccoli Raab, Brussels Sprout, Cabbage, Cauliflower, Chinese Broccoli, Chinese Cabbage, Collards, Kale, Kohlrabi, Rutabagas and Turnips)

Pest Controlled: root maggot

Application Rate: 4.6 to 9.2 ounces per 1,000 feet of row

At Plant T-Band Treatment: For direct seeded and transplanted crops, apply Pilot 15G as a 4-inch wide band centered over the row. This application requires a spreader or splitter on the end of the applicator drop tube. Shallow incorporation is necessary. Placement behind the planter shoe and in front of the press wheel is recommended.

**Restrictions:** Do not apply more than 7 1/2 pounds of Pilot 15G\_per acre to crops planted in 40 inch rows or more than 15 pounds of Pilot 15G per acre to crops planted in 20 inch rows (or two rows per bed). Use proportional amounts for other row spacing not to exceed 15 pounds of Pilot 15G per acre. Do not make more than one application per season. Do not use rutabaga tops for food or feed purposes.

#### **Radishes**

Pest Controlled: root maggot

Application Rate: 3.3 ounces per 1,000 feet of row

At Plant In-Furrow Treatment: Place the granules in the seed furrow with

the seed at planting time.

**Restrictions:** Do not apply more than 18.3 pounds of Pilot 15G per acre or make more than one application per season.

Table 1 Application Rates Table-Application Rates/1,000 Ft. of Row and Equivalent/Acre at Different Row Spacing

Amount of Pilot 15G Per 1,000 Feet of Row	Pounds of Pilot 15G Required Per Acre from Various Row Spacing						
	40"	38"	36"	34"	32"	30"	22"
3.7 ounces	3.0	3.2	3.4	3.6	3.8	4.0	5.5
4.0 ounces	3.3	3.4	3.6	3.8	4.1	4.4	5.9
4.5 ounces	3.7	3.9	4.1	4.3	4.6	4.9	-
6.0 ounces	4.9	5.2	5.4	5.8	6.1	6.5	-
6.5 ounces	5.3	5.6	5.9	6.2	•	-	•
7.5 ounces	6.1	6.4	-	<u>-</u>		-	-
8.0 ounces	6.5	-	-	-	-	•	-

#### General Instructions for Calibration of Equipment

Caution: The following chart lists suggested initial gauge settings for application of Pilot 15G with one hopper opening per row. Be sure to check the actual application rate under your operating conditions.

- 1. Fill hopper.
- Attach a plastic bag to tube opening.
- Set your planter to the initial settings shown on chart.
   Measure off 1,000 row feet and drive your planter the pre- measured distance at your desired speed.
- 5. Each bag should contain 6 to 8 ounces (wt.) of granules depending on your desired rate.6. If the result is over or under the desired rate, adjust the settings and repeat the calibration.

Table 2
Equipment Calibration and Calibration Settings for Different Types of Equipment
Application Rate, 8 oz. Per 1,000 row ft.

	Speed (mph)					
	4	5	6	7	8	
Planter Type	Gauge Setting					
Gandy <sup>1</sup>	21.4	23.7	26.0	27.7	30.2	
John Deere <sup>1</sup> Max-Emerge <sup>2</sup>	20	26	30	35	40	
John Deere <sup>1</sup> 7000 Max-Emerge (Odd Nos. on Gate)	14	16	18	19	21	
John Deere <sup>1</sup> 7000 Max-Emerge (Even Nos. on Gate)	17	20	24	26	28	
John Deere <sup>2</sup> 71 Flexi-Planter and Older Planters	<u>1</u> 30	<u>2</u> 5	<u>2</u> 9	<u>2</u> 13	<u>2</u> 16	
Allis Chalmers <sup>3</sup> 70 Series	8	8	8	8	. 8	
Allis Chalmers <sup>4</sup> 78 & 79 Series	<u>1</u> 9.0	<u>2</u> 3.3	<u>2</u> 6.0	<u>2</u> 9.0	<u>3</u> 2.5	
Noble <sup>1</sup> (NEW)	11	14	16	17	19	
White Planter	11	14	16	-17	19	
International Harvester <sup>4</sup>	<u>1</u> 9.0	<u>2</u> 3.3	<u>2</u> 6.0	<u>2</u> 9.0	<u>3</u> 2.5	
Buffalo All-Flex <sup>5</sup> (Fleischer Mfg.)	4 7/8	4 7/8	4 7/8	-	-	

<sup>1</sup> Gauge setting

range

gate

#### **Notice of Warranty and Disclaimer**

Seller warrants that at the time of delivery the product in this container conforms to its chemical description contained hereon and is reasonably fit for its intended purpose under normal conditions of use. This is the only warranty made on this product. Seller expressly disclaims any implied warranties of merchantability or fitness for any particular purpose and, except as set forth above, any other express or implied warranties. Any damages arising from breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid for this product by Buyer, and shall not include incidental or consequential damages such as, but not limited to, loss of profits or values. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of the Seller. To the fullest extent permitted by law, in no event shall Seller be liable for the consequential, special or indirect damages resulting from the use or handling of this product. To the fullest extent permitted by law II such risks shall be assumed by the Buyer. Buyer acknowledges the use of its own independent skill and expertise in the selection and use of the product and does not rely on any oral or written statements or representations.

Revised: [TBA] (IRED Com Mitigation Language, and, PR Notice 2007-4)

Pilot® is a registered trademark of Gharda Chemicals Limited Accent Registered Trademarks of E.I. du Pont de Nemours and Company. Beacon Registered Trademark of Syngenta Crop Protection.

<sup>&</sup>lt;sup>2</sup> Gauge setting with range 1 & 2 - number is notch.

<sup>&</sup>lt;sup>3</sup> Gauge setting is constant regardless of speed.

<sup>&</sup>lt;sup>4</sup> Gauge setting shown with stem gates & dial settings - number shown is dial.

<sup>&</sup>lt;sup>5</sup> Number of turns open on the adjustment nut.