

Pin 25

33688-7

1/4

RESTRICTED USE PESTICIDE

FOR RETAIL SALE TO AND APPLICATION ONLY BY CERTIFIED APPLICATORS OR PERSONNEL UNDER THEIR DIRECT SUPERVISION.

AMITROL T™ brand Liquid Herbicide

CONTROLS QUACKGRASS, CANADA THISTLE, AND OTHER TOUGH ANNUAL AND PERENNIAL WEEDS

ACTIVE INGREDIENT

Amitrole (3-amino-1,2,4-triazole)*21.6%

INERT INGREDIENT78.4%

*Equivalent to 2 pounds per gallon

EPA Reg. No. 264-135-ZA

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN CAUTION

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-800-334-9745

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Induce vomiting immediately. Get medical attention.

IF ON SKIN: Wash thoroughly with soap and water. Remove contaminated clothing and wash before reuse.

IF IN EYES: Flush eyes with plenty of water.

NOTE TO PHYSICIAN

This product may affect thyroid function. Overexposure can be determined by monitoring thyroid function. If worker is under treatment for hypo- or hyperthyroidism, he should be removed from spray crew until the euthyroid state is achieved.

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Avoid contact with eyes, skin, and clothing.

The use of this product may be hazardous to your health. This product contains Amitrole which has been determined to cause cancer in laboratory animals. Wear a respirator during mixing and loading of this product.

When applying this product, wear clean cotton (or cloth) coveralls that cover all parts of the body except the head, hands, and feet. Coveralls must be worn over a long sleeve shirt and long pants. Wear boots or sturdy footwear. When handling the concentrate or when repairing and cleaning the equipment used with this product, chemical resistant gloves must also be worn.

ENVIRONMENTAL HAZARDS

Do not allow spray or spray drift to contaminate edible crops or water intended for irrigation, drinking or other domestic purposes as no chemical residue in crops is permitted. Spray drift from this product may injure cotton, tomatoes, lawns, ornamentals, and other desirable vegetation. Coarse sprays are less likely to drift.

Do not use spray equipment for other purposes.

Do not use in a greenhouse.

Do not allow spray solution to remain in a sprayer any longer than necessary. Rinse all spray equipment with clean water immediately after each use to prevent corrosion of metal parts.

Do not allow livestock to graze or feed in treated non-crop areas.

Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

ACCEPTED
27 17 1991
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 264-135

DIRECTION FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
Read entire label before using this product.

STORAGE AND DISPOSAL

STORAGE

Do not contaminate water, food, or feed by storage or disposal. Do not store near fertilizers, seeds, insecticides, or fungicides.

PESTICIDE DISPOSAL

Open dumping is prohibited. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Do not reuse empty container. Triple rinse or equivalent. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

Do not use more than 4 gallons (8 lbs. active ingredient) of AMITROL-TM per acre per year.

HOW TO APPLY

Apply enough of the spray mixture to wet thoroughly all leaves and stems to ground line. Use a fixed boom sprayer attached to a tractor/truck or railroad wagon that provides uniform and accurate application. Sprayer should be equipped with flat fan nozzles spaced a maximum of 20 inches apart and screens no finer than 50 mesh in the nozzle tips and in-line strainers. Specific application instructions are described under each weed controlled heading.

WOODY PLANTS AND VINES

	GALLONS OF AMITROL-TM	GALLONS OF WATER	WHEN TO APPLY
Poison ivy - poison oak	1	100	Treat from the time foliage is fully developed until plants begin to go dormant. For effective control, all leaves, stems and suckers must be thoroughly wet to ground line. *If regrowth of honeysuckle appears, retreat in August or the following Spring.
Honeysuckle* - Kudzu	2	100	
Salmon berry - Western dewberry (blackberry) - Big leaf maple	2	100	
Wild cherry - Ash-Locust-Sumac	1 1/2	100	

**WEEDS IN NON-CROP AREAS SUCH AS RAILROADS, ROADSIDES,
FENCE ROWS, AND ORNAMENTAL HARDWOOD NURSERIES**

WEEDS	Gallons AMITROL-T™	Gallons Water	Gallons AMITROL-T™	Min. Gals. Water	WHERE TO APPLY
CONTROL: Annual grasses and broadleaf weeds (foxtail, ryegrass, cheatgrass, barnyard grass, annual bluegrass, wild hemp or marijuana, pigweed, kochia, sunflower, wild barley)	1/2	50	1	100	Spray when most weed are 3 to 4 inches tall.
CONTROL: Annual grasses, annual and perennial broadleaf weeds (whitetop, Canada thistle, volunteer alfalfa, milkweed) SUPRESS: Perennial grasses, (ripgut grass, reed canary grass, quackgrass)	1	50	2	100	Spray when most weeds are 6 to 10 inches tall.
Canada thistle Sowthistle	1	50	2	40	Spray when most thistle are up and in the bud to bloom stage. Do not mow treated plants in the Far West, treat during the bloom stage.
Quackgrass (couch, witch)	1	50	2	40	Spray when vigorous young growth is 6 to 8 inches tall.
Whitetop (hoary cress)	1	50	2	40	Spray in Spring when weeds are in the bud to bloom stage. Do not mow treated plants.
Leafy spurge (Western states)	1	50	2	40	
Milkweed	1	50	1	50	
Bermudagrass	2	50	4	50	Treat from Mid-Summer to early Fall when shoots are growing vigorously.
Nutsedge (nut grass)	1	50	2	50	Spray when shoots are 4 to 5 inches tall and growing actively.
Horsenettle	1	50	2	100	In East and Midwest, spray when horsenettle is in the bud to bloom stage. In the Far West treat from Mid-Summer to early frost.
Horsetail rush	1 1/2	50	2	100	Spray when 6 to 8 inches tall and growing actively.
Dock	1/2	50	1	50	Apply anytime when dock is actively growing; preferably before development of bloomstalk.
Wild chrysanthemum in ornamental hardwood nurseries	2	50	4	100	Apply as directed spray on young weed growth 4 to 6 inches tall. Keep spray off foliage of ornamental hardwood nursery stock.

1/ A second treatment may be necessary to control weeds not thoroughly sprayed or weeds coming from dormant seeds and nuts or old roots. Apply when regrowth is young and growing actively,
 2/ INTERMOUNTAIN STATES: Canada thistle, use 4 gals. of AMITROL-T™ in 100 gals. of water per acre. All other weeds, use the rates of AMITROL-T™ per acre as listed in the preceding chart, applied in a minimum of 100 gals. of water per acre.

4/4

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

Rhône-Poulenc Ag Company
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
AMITROL is a trademark of Rhône-Poulenc.

AMITROL T brand Liquid Herbicide (PENDING) Submitted 10/1/91.