

Sanitizing of Food Processing Equipment and Other Hard Surfaces in Food Contact Locations: For sanitizing food processing equipment, dairy equipment, and other hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

Disinfection of Hard Surfaces: For disinfecting hard surfaces, use a solution of **TOTL-150** prepared daily or when the use solution becomes visibly dirty. **No Potable water rinse is required.**

ACCEPTED
with COMMENTS
EPA REG. NO. 33821-8

TOTL-150

**Disinfectant-Sanitizer
Fungicide-Virucide* Deodorizer
with Organic Soil Tolerance
for Hospital, Institutional
Industrial, School, Dairy and Other Farm Use**

registered under EPA Reg. No. 33821-8

Active Ingredients	
Octyl decyl dimethyl ammonium chloride	2.250%
Didecyl dimethyl ammonium chloride	1.125%
Diocetyl dimethyl ammonium chloride	1.125%
Alkyl (C ₁₄ , 50%; C ₁₂ , 40%; C ₁₆ , 10%) dimethyl benzyl ammonium chloride	3.000%
Inert Ingredients	92.500%
	100.000%

KEEP OUT OF REACH OF CHILDREN

DANGER

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if there are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Registration No.

EPA Establishment No.

Net Contents

MANUFACTURED BY
K-1

**For Sale For Use And Storage
By Maintenance Personnel Only**

**Directions for Use
GENERAL CLASSIFICATION**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply **TOTL-150** with a mechanical or mechanical spray device. When applied with a mechanical spray device, surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the use solution becomes visibly dirty.

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions: For disinfecting floors, walls, counter tops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard non-porous surfaces.

Add 35 oz **TOTL-150** to 45 gallons water. Apply to previously cleaned hard surfaces. At this use level, **TOTL-150** is effective against *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella choleraesuis* in the presence of 5% blood serum when evaluated by the AOAC Use-Dilution test.

***Virucidal Performance:** At a 35 oz/45 gallon use level, **TOTL-150** was evaluated in the presence of 5% serum and found to be effective against the following viruses: Herpes simplex, Vaccinia and Influenza A (after 1 hour) on inanimate environmental surfaces.

Disinfection in Institutions, Industry and Schools: Add 35 oz of **TOTL-150** to 45 gallons of water.

Fungicidal Performance: At 0.5 oz to 2.25 gallons of water use level, **TOTL-150** is an effective fungicide against *Trichophyton mentagrophytes* (the athlete's foot fungus) when used on surfaces in areas such as locker room, dressing rooms, shower and bath areas and exercise facilities utilizing the AOAC Fungicidal Test.

Disinfection of Barber Tools: Pre-cleaned barber tools, such as combs, brushes, razors and scissors, can be disinfected by immersing in a 0.5 oz/gallon solution of **TOTL-150**.

Disinfection of Poultry Equipment, Animal Quarters and Kennels: Poultry brooders, watering fountains, feed equipment and other animal quarters, such as stalls in kennel areas, can be disinfected after thorough cleaning by applying a solution of 0.25 oz **TOTL-150** to 45 gallons of water. Small pens should be immersed in the solution. Prior to disinfection, poultry, other animals and their feeds must be removed from the premises. This solution is also effective against ticks and other ectoparasites on animal quarters. Remove all litter and droppings from floors, walls and other surfaces before immersing in the solution.

Disinfection of Poultry Equipment, Animal Quarters and Kennels: Poultry brooders, watering fountains, feed equipment and other animal quarters, such as stalls in kennel areas, can be disinfected after thorough cleaning by applying a solution of 0.25 oz **TOTL-150** to 45 gallons of water. Small pens should be immersed in the solution. Prior to disinfection, poultry, other animals and their feeds must be removed from the premises. This solution is also effective against ticks and other ectoparasites on animal quarters. Remove all litter and droppings from floors, walls and other surfaces before immersing in the solution.

Disinfection of Poultry Equipment, Animal Quarters and Kennels: Poultry brooders, watering fountains, feed equipment and other animal quarters, such as stalls in kennel areas, can be disinfected after thorough cleaning by applying a solution of 0.25 oz **TOTL-150** to 45 gallons of water. Small pens should be immersed in the solution. Prior to disinfection, poultry, other animals and their feeds must be removed from the premises. This solution is also effective against ticks and other ectoparasites on animal quarters. Remove all litter and droppings from floors, walls and other surfaces before immersing in the solution.

Disinfection of Poultry Equipment, Animal Quarters and Kennels: Poultry brooders, watering fountains, feed equipment and other animal quarters, such as stalls in kennel areas, can be disinfected after thorough cleaning by applying a solution of 0.25 oz **TOTL-150** to 45 gallons of water. Small pens should be immersed in the solution. Prior to disinfection, poultry, other animals and their feeds must be removed from the premises. This solution is also effective against ticks and other ectoparasites on animal quarters. Remove all litter and droppings from floors, walls and other surfaces before immersing in the solution.

Disinfection of Poultry Equipment, Animal Quarters and Kennels: Poultry brooders, watering fountains, feed equipment and other animal quarters, such as stalls in kennel areas, can be disinfected after thorough cleaning by applying a solution of 0.25 oz **TOTL-150** to 45 gallons of water. Small pens should be immersed in the solution. Prior to disinfection, poultry, other animals and their feeds must be removed from the premises. This solution is also effective against ticks and other ectoparasites on animal quarters. Remove all litter and droppings from floors, walls and other surfaces before immersing in the solution.

Disinfection of Poultry Equipment, Animal Quarters and Kennels: Poultry brooders, watering fountains, feed equipment and other animal quarters, such as stalls in kennel areas, can be disinfected after thorough cleaning by applying a solution of 0.25 oz **TOTL-150** to 45 gallons of water. Small pens should be immersed in the solution. Prior to disinfection, poultry, other animals and their feeds must be removed from the premises. This solution is also effective against ticks and other ectoparasites on animal quarters. Remove all litter and droppings from floors, walls and other surfaces before immersing in the solution.

Disinfection of Poultry Equipment, Animal Quarters and Kennels: Poultry brooders, watering fountains, feed equipment and other animal quarters, such as stalls in kennel areas, can be disinfected after thorough cleaning by applying a solution of 0.25 oz **TOTL-150** to 45 gallons of water. Small pens should be immersed in the solution. Prior to disinfection, poultry, other animals and their feeds must be removed from the premises. This solution is also effective against ticks and other ectoparasites on animal quarters. Remove all litter and droppings from floors, walls and other surfaces before immersing in the solution.


Disinfection of Poultry Equipment, Animal Quarters and Kennels: Poultry brooders, watering fountains, feed equipment and other animal quarters, such as stalls in kennel areas, can be disinfected after thorough cleaning by applying a solution of 0.25 oz **TOTL-150** to 45 gallons of water. Small pens should be immersed in the solution. Prior to disinfection, poultry, other animals and their feeds must be removed from the premises. This solution is also effective against ticks and other ectoparasites on animal quarters. Remove all litter and droppings from floors, walls and other surfaces before immersing in the solution.

BEST DOCUMENT AVAILABLE

<p>U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDE REGISTRATION WASHINGTON, D.C. 20460</p> <p>NOTICE OF PESTICIDE Label for [Product Name] EPA Registration No. [Number]</p>	<p>EPA REGISTRATION NO. [Number]</p> <p>TERM OF ISSUE [Date]</p>	<p>DATE 09 MAR 1983</p>
<p>NAME AND ADDRESS OF REGISTRANT (Include ZIP)</p> <p>[Name] [Address] [City, State, ZIP]</p>		
<p>NOTE: Changes in labeling formula, different active ingredients, or other information with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.</p>		
<p>A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.</p>		
<p>Registration is in no way to be construed as an endorsement or approval of the product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p>		
<p><input checked="" type="checkbox"/> ATTACHMENT 5 APPLICABLE</p>		
<p>SIGNATURE OF APPROVING OFFICIAL</p>	<p>DATE</p>	

If the conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 4(c). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.



John H. Lee
Product Manager (11)
Disinfectants Branch
Registration Division (TS-757)

Enclosure