

33906-17

4-2-2003

1/8



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg.
Number:

33906-17

Date of Issuance:

APR 2 2003

Term of Issuance:

Conditional

Name of Pesticide Product:

NC-603 WG Herbicide

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Nissan Chemical Industries, Ltd.
c/o Lewis and Harrison LLC
122 C Street, N.W., Suite 740
Washington, DC 20001

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit data.
2. Make the following label changes listed below before you release the product for shipment:
 - a. Change the registration number to "EPA Reg. No. 33906-17".
 - b. In the "Precautionary Statements" and "Agricultural Use Requirements" sections, change "waterproof gloves" to "chemical resistant gloves such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber ≥ 14 mils".
 - c. Replace the language in the "Environmental Hazards" section with the following (to change "this chemical" to the appropriate herbicide):

Carfentrazone-ethyl is very toxic to algae and moderately toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high water mark, except as specified on this label. Do not contaminate water when disposing of equipment wash waters.

Signature of Approving Official:

Date:

4-2-03

Halosulfuron-methyl demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of halosulfuron-methyl in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

c. In the "Pesticide Disposal" section, delete "at a disposal facility" in the sentence that begins "In case of spills...".

d. Replace the first two sentences in the "Tank Mixtures" subsection of the "General Information" section with the following:

NC-603 WG may be tank mixed with other herbicides labeled for use on corn (field, pop, seed, silage and sweet) to control weeds not listed on this label provided that the tank mix product does not prohibit such mixing. Read the manufacturer's label directions, restrictions, and use precautions for the companion herbicide. Follow the most prohibitive restrictions and use precautions for all products used.

e. In the "Adjuvant Use Requirements" section, spell out "urea ammonium nitrate" prior to the abbreviation "UAN".

f. In the "Application Information" section, add the following spray drift subsection:

Spray Drift Mangement

For ground boom applications, apply with a nozzle height of no more than 4 feet above the ground or crop canopy, and when wind speed is 10 mph or less at the application site. Use nozzles that produce a fine or coarser spray.

g. In the "Application Restrictions" section, add the following restriction:

For all herbicide products used during the use season, the total pounds applied of carfentrazone and halosulfuron must not exceed the pounds a.i. per acre per season specified in the table below.

h. In the "Application Restrictions" section, change the maximum amounts applied of product and active ingredients for field and seed corn to the following: 2 oz product, 0.016 lbs a.i. carfentrazone, and 0.064 lbs a.i. halosulfuron. In addition, add a row for silage corn using the the maximum application amounts listed above.

i. In the "For Broadcast Applications" section, change the subheading "Field Corn Production" to "Field Corn and Silage Production" In this subsection, change "adjuvant section" to "adjuvant use requirements section".


j. In the "For Directed Applications" section, change the subheading "Field Corn" to "Field Corn and Silage".

3/8

3. Submit one copy of your final printed labeling before you release the product for shipment. In addition, please submit an electronic pdf copy of this label within one month of receipt of his notice.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.



Jim Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505C)

4/8

NC-603 WG Herbicide

For Agricultural or Commercial Use Only NOT FOR SALE OR USE IN CALIFORNIA

For control of specified weeds on Field Corn,
Popcorn, Seed Corn, Silage Corn and Sweet Corn

EPA Reg. No. 33906-XXXX EPA Est. 11773-IA-01

Active Ingredient:	By Wt.
Carfentrazone-ethyl	12.5
Halosulfuron-methyl	50.0
Inert Ingredients:	37.5%
	100.0%

Contains 62.5% W/W of active ingredient per pound of product
U.S. Patent Numbers 5,125,958 and 4,668,277

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If In Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If Swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

Note to Physician: This product is expected to have low oral and dermal toxicity, and moderate inhalation toxicity. It is expected to be slightly irritating to the skin and minimally irritating to the eyes. This product contains a granular material (sand) that may cause mechanical irritation to the eyes. Treatment is removal of exposure followed by symptomatic and supportive care.

See other panels for additional precautionary information.

Nissan Chemical Industries, Ltd.
7-1, 3-chome, Kanda-Nishiki-cho
Chiyoda-ku, Tokyo 101-0054
Japan
NC-603 WG_3_3-12-2003b

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Caution

Harmful if swallowed, absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This chemical is very toxic to algae and moderately toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high water mark, except as specified on this label. Do not contaminate water when disposing of equipment wash waters.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Physical/Chemical Hazards

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

ACCEPTED
with COMMENTS
In EPA Letter Dated

APR 2 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

33906-17

5/8

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, waterproof gloves, and shoes plus socks.

STORAGE AND DISPOSAL

Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

Pesticide Storage

Not for use or storage in or around the house.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place (below 120°F) and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put granule or dilute material into food or drink containers.

In case of spill, avoid contact, isolate area and keep out unprotected persons and animals. Confine spills.

To confine spill: Dike surrounding area, sweep up spillage. Dispose of in accordance with information given under Pesticide Disposal. Wash spill area with water, absorb with sand, cat litter or commercial clay, sweep up and dispose of in a landfill approved for pesticide disposal in accordance with applicable Federal, state and local procedures, or by such other method as is approved under those procedures. Contact your State Pesticide or Environmental Control Agency, or your local Hazardous Waste Disposal office or the Hazardous Waste Representative at the nearest EPA Regional Office if further guidance is needed. Place damaged container in a larger holding container. Identify contents per required hazardous waste labeling regulations.

Pesticide Disposal

Waste resulting from the use of this product may be disposed of at an approved waste disposal facility.

In case of spills of this product at a disposal facility, refer to the instructions in the Pesticide Storage subsection above for avoiding contact, isolating, confining and cleaning up a spill.

Container Disposal

Plastic containers: Triple rinse (or equivalent). Then offer for approved pesticide container recycling program, or puncture and dispose of in an approved waste disposal facility. Provided on site incineration is allowed by state and local authorities, stay out of smoke.

GENERAL INFORMATION

NC-603 WG is a water dispersible granule formulation. NC-603 WG is to be mixed with water and applied to labeled crops for selective postemergence control of broadleaf weeds. Weed control is best when the product is applied to actively growing weeds up to 4 inches in height. NC-603 WG is a contact and translocated herbicide with minimal residual activity at recommended use rates.

NC-603 WG is rapidly absorbed through the foliage of plants. To avoid significant crop response, applications should not be made within 6 - 8 hours of either rain or irrigation. Within a few hours following application, the foliage of susceptible weeds show signs of desiccation, and in subsequent days, necrosis and death of the plant occur. Some environmental conditions and with certain spray tank additives, herbicidal symptoms may appear on the crop. However, the crop recovers quickly with no loss in yield.

Extremes in environmental conditions such as temperature, moisture, soil conditions, and cultural practices may affect the activity of NC-603 WG. Under warm moist conditions, herbicide

symptoms may be accelerated. While under very dry conditions, the expression of herbicide symptoms is delayed, and weeds hardened off by drought are less susceptible to NC-603 WG.

Tank Mixtures

NC-603 WG may be tank mixed with other herbicides to control weeds not listed on this label. Read and follow all manufacturers' label recommendations for the companion herbicide except for specific recommendations on this label. Tank mixtures of NC-603 WG with EC formulations of other crop protection products, crop oil concentrate, methylated seed oil, silicone based adjuvants, 28% nitrogen or ammonium sulfate may increase crop response.

Adjuvant Use Requirements

Use a non-ionic surfactant (NIS) having at least 80% active ingredient with a minimum of 65% surfactant load, at 0.25% v/v (2 pints per 100 gallons of spray solution). Spray grade ammonium sulfate (AMS) may be used at 2-4 pounds/acre. 28% UAN may be used instead of AMS at 2-4 qts/100 gallons of spray solution. The combination of adjuvants may enhance the level of leaf speckling than either adjuvant alone. Crop oil (COC) or crop oil plus either ammonium sulfate or 28% nitrogen may be used with companion herbicides listed on this label and may be recommended in certain situations.

Mixing and Loading Instructions:

Fill the spray tank 3/4 full with clean water. Make sure the agitation system is operating. Prepare a slurry of NC-603 WG in a clean container using clean water. Slowly add the NC-603 WG/water slurry. Carefully rinse the slurry container adding the rinsate to the spray tank. Complete filling the spray tank to the desired level. The spray tank agitation should be sufficient to ensure uniform spray mixture during application and until the spray tank has been emptied. When tank mixing with other products, NC-603 WG should be mixed first in the spray tank. After the NC-603 WG is thoroughly mixed, add the other products as specified on their label. Ensure the compatibility of other products with NC-603 WG before mixing them together in the spray tank. Avoid the overnight storage of NC-603 WG spray mixtures. Premixing NC-603 WG solutions in nurse tanks is not recommended.

Do not use with tank additives that alter the pH of the spray solution below pH 5 or above pH 8. Buffer spray solution to alter the pH range if appropriate.

Spray Equipment Clean-Out:

After spraying NC-603 WG and before using sprayer equipment for any other applications, the sprayer must be thoroughly cleaned using the following procedure. In addition, users must take appropriate steps to ensure proper equipment clean-out for any other products mixed with NC-603 WG herbicides as required on the other product labels.

1. Drain sprayer tank, hoses, and spray boom. Use a high pressure wash to remove physical sediment and residues from the inside of the sprayer tank and thoroughly rinse. Then thoroughly flush sprayer hoses, boom and nozzles with clean water.
2. Fill the tank 1/2 full with clean water, and add appropriate tank cleaner (follow manufacturer's directions for use). Fill the tank to capacity and operate the sprayer for 15 minutes to flush hoses, boom and nozzles.
3. Drain the sprayer system. Rinse the tank with clean water and flush through the hoses, boom, and nozzles. Remove and clean spray tips and screens separately.
4. Properly dispose of all cleaning solution and rinsate in accordance with Federal, State, and local regulations and guidelines.

APPLICATION INFORMATION

Ground Applications

Utilize a boom and nozzle sprayer equipped with the appropriate nozzles, spray tips and screens and adjusted to provide optimum spray distribution and coverage at appropriate operating pressures. Utilize nozzles which produce minimal amounts of fine spray droplets. Do not exceed 30 psi spray pressure unless otherwise required by the manufacturer of drift reducing nozzles. Apply a minimum of 10 gallons of finished spray per acre. Higher spray volumes are required when there is a dense weed population or crop canopy. Spray booms should be adjusted to position spray tips at the minimum height of 18" above the crop and operated to avoid the application of excessive herbicide rates directly over the rows and/or into the whorl of the crop. Be aware that overlaps and slower ground speeds while starting, stopping or turning while spraying may result in higher application rates and possible crop response.

Additives – Agriculturally approved deposition and retention additives may be added to the spray tank to assist in the application process. Not all additives have been tested with NC-603 WG. Additives that are not susceptible to pump shear degradation have shown to improve spray deposition and retention. Follow manufacturers' label and recommendations.

Application Restrictions

Do not apply this product through any type of irrigation system.

Following application to foliage, allow 30 days before grazing domestic livestock, harvesting forage, or harvesting silage.

Do not apply more than 2 applications per season.

MAXIMUM ALLOWABLE NC-603 WG USE PER ACRE PER SEASON

Total Allowed NC-603 WG Use*

Crop	Ounces NC-603 WG	Pounds a.i. Carfentrazone	Pounds a.i. Halosulfuron
Field Corn	4.0	0.031	0.125
Seed Corn	4.0	0.031	0.125
Sweet Corn	2.0	0.016	0.064
Popcorn	2.0	0.016	0.064

*The total allowed usage volumes include all applications made to the field per calendar year. This includes fallow treatments, burndown treatments and all in-season treatments.

The 2nd treatment of NC-603 WG may be applied only with drop nozzles semi-directed or directed.

CROP ROTATIONAL RESTRICTIONS

Labeled crops may be planted at specified time intervals following application of approved rates of NC-603 WG herbicide. Use the time intervals listed below to determine the required time interval before planting.

TIME INTERVAL BEFORE REPLANTING (Months after treatment with NC-603 WG)

Crop	Months
Field Corn	1
Barley (Winter)	2
Oats (Winter)	2
Rice	2
Seed corn	2
Sorghum	2
Spring cereal crops (wheat, oats, barley)	2
Wheat (winter)	2
Popcorn, Sweetcorn	3
Cotton	4
Soybeans	9
Alfalfa	12
Clovers	12
Cucumbers, Pumpkins, Squash	12
Dry beans	12
Eggplant	12
Field peas	12
Forage Grasses	12
Peanuts	12
Peas	12
Peppers	12
Potatoes	12
Proso Millet	12
Radish	12
Rye (winter)	12
Snap Beans	12
Spring Cereal crops (other)	12
Tomato (transplant)	12
Cabbage	15
Canola	15
Carrot	15
Mint	15
Broccoli, Cauliflower, Collards	18
Leeks, Onions	18
Lettuce crops	18
Sunflowers	18
Sugarbeet (Michigan only)	21
Sugarbeet and Red Beet	24
Spinach	24
Sugarbeet (ND, MN, Red River Valley)*	36

*Also includes other areas where rainfall is sparse or irrigation is required.

Refer to individual product labels to determine rotation crop restrictions when tank mixtures are used.

7/8

CORN

Field Corn, Seed Corn, Popcorn, Corn Silage, and Sweet Corn (Processing and Fresh Market)

Apply NC-603 WG alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply to corn in all tillage systems from crop emergence up to 8-leaf collar corn growth stage. Do not apply when conditions favoring drift exist or wind is above 10 mph.

For best performance, make application to actively growing weeds up to 4 inches high and rosettes less than 3 inches across.

Coverage is essential for good control

To control weeds not listed on this label, NC-603 WG may be tank mixed with other herbicides registered for use in corn. When tank mixing NC-603 WG with other products, be sure the NC-603 WG is mixed in the spray tank water first. For specific mixing instructions, refer to the Mixing and Loading Instructions under the GENERAL INFORMATION section.

For Broadcast Applications

Field Corn Production

Use 1.0 oz NC-603 WG product per acre plus NIS at 0.25 % volume to volume (2 pints per 100 gallons) with or without UAN or AMS as described in the adjuvant section.

Apply NC-603 WG alone or as a tank mixture with other herbicides to emerged and actively growing weeds.

Sweet Corn/ Popcorn Production

NC-603 WG may be applied to sweet corn, however, the user assumes all responsibility for herbicide tolerance with such use. All hybrids/varieties have not been tested for sensitivity to NC-603 WG herbicide nor does Nissan Chemical Industries, Ltd. have access to all seed company or food processor data. Broadcast applications may result in spray being concentrated into the whorl of the plant, which will increase leaf response. To minimize application into the whorl of the plants, drop nozzles or other type directed sprayers must be used to direct the spray to the targeted weeds.

Therefore, any crop response arising from the use of NC-603 WG herbicide on sweet corn is the responsibility of the user. Use NC-603 WG herbicide only under the recommendation of the seed company, food processor, or State Agricultural Extension Service.

If necessary, a sequential treatment of this product at 1 oz of product per acre may be applied only with drop nozzles directed to avoid application into the corn plant whorl.

For Directed Applications

Field corn

NC-603 WG may be applied with drop nozzles or other sprayers capable of directing the spray to the target weeds and away from the whorl of the corn plant. NC-603 WG may be used at 1.0 ounce of formulated product per acre using drop nozzles. Use appropriate rates of adjuvants.

Sweet Corn, Popcorn Production

NC-603 WG may be applied with drop nozzles or other sprayers capable of directing the spray to the target weeds and away from the whorl of the corn plant. NC-603 WG may be used up to a maximum of 1.0 ounce of formulated product per acre using drop nozzles for control of larger weed sizes for those weeds listed below under "Control of Weeds". Use appropriate rates of adjuvants.

Seed Corn Production (Directed Applications Only)

For seed production fields, apply NC-603 WG at 1.0 ounce formulated product per acre using drop nozzles or other equipment to make a directed spray treatment. Avoid directing spray solution into the whorl.

Seed corn inbreds have generally shown good tolerance to NC-603 WG herbicide, however, all inbreds have not been tested.

Weeds controlled with NC-603 WG

Use NC-603 WG at 1.0 ounce/acre (0.008 pound active ingredient carfentrazone + 0.032 pound active ingredient halosulfuron). Applications should be made by ground equipment using a minimum finished spray volume of 10 gallons of spray per acre using nonionic surfactant at 0.25% v/v (2 pt per 100 gallons of water) with or without UAN or AMS as described in the adjuvant section of this label.

When used as directed, NC-603 WG will provide control of listed weeds up to the indicated sizes:

	1 oz NC-603 WG per acre (Broadcast or Directed Spray)	1 oz NC-603 WG per acre plus Banvel/Clarity (Broadcast or Directed Spray)
Cocklebur, Common	9 inches	9 inches
Kochia	3 inches	5 inches
Pokeweed, Common	6 inches	6 inches
Ragweed, Common	9 inches	9 inches
Ragweed, Giant	3 inches	6 inches
Smartweed, Pennsylvania	2 inches	6 inches
Sunflower, Common	12 inches	12 inches
Common Lambsquarters	3 inches	5 inches
Morningglories	2-3 true leaves	2-3 true leaves
Nightshade, Eastern Black	4 inches	4 inches
Pigweed, Redroot	4 inches	8 inches
Pigweed, Amaranth resistant	4 inches	8 inches
Pigweed, other	Suppression	4 inches
Velvetleaf	18 inches	18 inches
Waterhemp, Common	2 inches (with COC at 1% v/v)	4 inches
Waterhemp, Tall	2 inches (with COC at 1% v/v)	4 inches
Nutsedge, Purple	Suppression	Suppression
Nutsedge, Yellow	Suppression	Suppression
Vernice Mallow	3 inches	3 inches
Pinkweed, Barnyard	Suppression	4 inches
Burcucumber	Suppression	4 inches
Field Bindweed	Suppression	4 inches
Jimsonweed	Suppression	3 inches
Spurred Anoda	Suppression	Suppression
Potato, volunteer	3 inches	6 inches
Prickly Side	Suppression	2 inches
Prostrate Spurge	Suppression	Suppression
Blackberry	3 inches	5 inches
Wild Buckwheat	3 inches	6 inches
Barnyard	3 inches	6 inches
Hemp Sesbania	6 inches	6 inches
Common Bean, Copperleaf	2 inches	4 inches
Spurred Anoda	2 inches	6 inches
Dayflower	2 inches	4 inches
Wild Mustard	Suppression	4 inches
Marestail	Suppression	4 inches
Common Purslane	Suppression	4 inches

8/8

Tank Mixtures

NC-603 WG may be tank mixed with other herbicides, labeled for use on corn (field, pop, seed, silage and sweet) to control weeds not listed on this label provided that the tank mix product does not prohibit such mixing. Read the manufacturer's label directions, restrictions, and use precautions for the companion herbicide.

Follow the most prohibitive restrictions and use precautions for all products used. When tank mixing NC-603 WG with other products, be sure NC-603 WG is mixed in the spray tank water first.

For control of additional broadleaf weeds and grasses, NC-603 WG may be tank mixed with 2,4-D (amine), Accent®, Accent Gold®, atrazine, Banvel®, Basise®, Basis Gold®, Beacon®, Callisto™, Clarity™, Distinct®, Exceed®, Hornet®, Liberty®, Lightning®, Marksman®, Northstar™, Permit®, Roundup®, or other glyphosate products, Roundup UltraMax®, Sencor®, Spirit™, Steadfast, Touchdown® and Yukon™.

When tank mixing NC-603 WG with other products, use the adjuvants recommended on the tank mix partner label. These may include nonionic surfactant, crop oil concentrate, 28% nitrogen, ammonium sulfate or combinations of these.

Leaf speckling can occur when NC-603 WG is used with certain crop protection products and adjuvants. Refer to the Tank Mixtures and Recommended Adjuvants sections under General Information.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: Nissan makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product: Nissan's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Accent, Accent Gold, Basis, Basis Gold - trademarks of E.I.

DuPont de Nemours and Company

Beacon, Exceed, Northstar, Spirit, Touchdown - trademarks of Syngenta

Banvel, Clarity, Distinct, Marksman, Lightning - trademarks of BASF Corporation

Hornet- trademark of Dow Agrosciences, LLC

Permit - trademark of Nissan Chemical Industries, Inc.

Roundup, Roundup UltraMax - trademarks of Monsanto Company

Sencor - trademark of Bayer CropScience

Nissan Chemical Industries, Ltd.

7-1, 3-chome, Kanda-Nishiki-cho

Chiyoda-ku, Tokyo 101-0054

Japan