

33906-17

9/25/2003

1/6



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

SEP 25 2003

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Lewis and Harrison, LLC
Attn: Wendy McCombie
122 C. Street NW, Suite 140
Washington, DC 20001

Dear Ms McCombie:

Subject: Label amendment for additional use on grain sorghum
Product: NC-603 WG
EPA Registration Number: 33906-17
Application Dated: 09/02/03

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable. The addition of grain sorghum to the label as well as the other minor amendments have been reviewed and accepted. Amended labeling supercedes all previously accepted labels. A stamped copy of labeling is enclosed for your records.

Sincerely,

A handwritten signature in black ink, appearing to read "Jim Tompkins", is written over the typed name and title.

Jim Tompkins
Team Leader, Team 25
Herbicide Branch
Registration Division (7505C)

Enclosure

ACCEPTED
in EPA Letter Dated
SEP 23 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

33906-17

NC-603 WG Herbicide

INSERT NC-603 WG LOGO.

For Agricultural or Commercial Use Only NOT FOR SALE OR USE IN CALIFORNIA

For control of specified weeds on Field Corn, Popcorn, Seed
Corn, Silage Corn, Sweet Corn, and Grain Sorghum

EPA Reg. No. 33906-17 EPA Est. 11773-IA-01

Active Ingredient:	By Wt.
Carfentrazone-ethyl	12.5
Halosulfuron-methyl	50.0
Inert Ingredients:	37.5%
	100.0%

Contains 62.5% W/W of active ingredient per pound of product
U.S. Patent Numbers 5,125,958 and 4,668,277

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If Swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

Note to Physician: This product is expected to have low oral and dermal toxicity, and moderate inhalation toxicity. It is expected to be slightly irritating to the skin and minimally irritating to the eyes. This product contains a granular material (sand) that may cause mechanical irritation to the eyes. Treatment is removal of exposure followed by symptomatic and supportive care.

See other panels for additional precautionary information.

PRECAUTIONARY STATEMENTS Hazards to Humans (and Domestic Animals) Caution

Harmful if swallowed, absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical resistant gloves such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber ≥ 14 mils, and shoes plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

Carfentrazone-ethyl is very toxic to algae and moderately toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high water mark, except as specified on this label. Do not contaminate water when disposing of equipment wash waters.

Halosulfuron-methyl demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of halosulfuron-methyl in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Physical/Chemical Hazards

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, chemical resistant gloves such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber \geq 14 mils, and shoes plus socks.

STORAGE AND DISPOSAL

Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

Pesticide Storage

Not for use or storage in or around the house.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place (below 120°F) and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put granule or dilute material into food or drink containers.

In case of spill, avoid contact, isolate area and keep out unprotected persons and animals. Confine spills.

To confine spill: Dike surrounding area, sweep up spillage. Dispose of in accordance with information given under Pesticide Disposal. Wash spill area with water, absorb with sand, cat litter or commercial clay, sweep up and dispose of in a landfill approved for pesticide disposal in accordance with applicable Federal, state and local procedures, or by such other method as is approved under those procedures. Contact your State Pesticide or Environmental Control Agency, or your local Hazardous Waste Disposal office or the Hazardous Waste Representative at the nearest EPA Regional Office if further guidance is needed. Place damaged container in a larger holding container. Identify contents per required hazardous waste labeling regulations.

Pesticide Disposal

Waste resulting from the use of this product may be disposed of at an approved waste disposal facility.

In case of spills of this product, refer to the instructions in the Pesticide Storage subsection above for avoiding contact, isolating, confining and cleaning up a spill.

Container Disposal

Plastic containers: Triple rinse (or equivalent). Then offer for approved pesticide container recycling program, or puncture and dispose of in an approved waste disposal facility. Provided on site incineration is allowed by state and local authorities, stay out of smoke.

GENERAL INFORMATION

NC-603 WG is a water dispersible granule formulation. NC-603 WG is to be mixed with water and applied to labeled crops for selective postemergence control of broadleaf weeds. Weed control is best when the product is applied to actively growing weeds up to 4 inches in height. NC-603 WG is a contact and translocated herbicide with minimal residual activity at recommended use rates.

NC-603 WG is rapidly absorbed through the foliage of plants. To avoid significant crop response, applications should not be made within 6 - 8 hours of either rain or irrigation. Within a few hours following application, the foliage of susceptible weeds show signs of desiccation, and in subsequent days, necrosis and death of the

plant occur. Under some environmental conditions and with certain spray tank additives, herbicidal symptoms may appear on the crop. However, the crop recovers quickly with no loss in yield.

Extremes in environmental conditions such as temperature, moisture, soil conditions, and cultural practices may affect the activity of NC-603 WG. Under warm moist conditions, herbicide symptoms may be accelerated. While under very dry conditions, the expression of herbicide symptoms is delayed, and weeds hardened off by drought are less susceptible to NC-603 WG.

Tank Mixtures

NC-603 WG may be tank mixed with other herbicides labeled for use on corn (field, pop, seed, silage and sweet) and grain sorghum to control weeds not listed on this label, provided that the tank mix product does not prohibit such mixing. Read the manufacturer's label directions, restrictions, and use precautions for the companion herbicide. Follow the most prohibitive restrictions and use precautions for all products used. Tank mixtures of NC-603 WG with EC formulations of other crop protection products, crop oil concentrate, methylated seed oil, silicone based adjuvants, 28% nitrogen or ammonium sulfate may increase crop response.

Adjuvant Use Requirements

Use a non-ionic surfactant (NIS) having at least 80% active ingredient with a minimum of 65% surfactant load, at 0.25% v/v (2 pints per 100 gallons of spray solution). Spray grade ammonium sulfate (AMS) may be used at 2-4 pounds/acre. 28% urea ammonium nitrate (UAN) may be used instead of AMS at 2-4 qts/100 gallons of spray solution. The combination of adjuvants may enhance the level of leaf speckling than either adjuvant alone. Crop oil (COC) or crop oil plus either ammonium sulfate or 28% nitrogen may be used with companion herbicides listed on this label and may be recommended in certain situations.

Mixing and Loading Instructions

Fill the spray tank 3/4 full with clean water. Make sure the agitation system is operating. Prepare a slurry of NC-603 WG in a clean container using clean water. Slowly add the NC-603 WG/water slurry to spray tank. Carefully rinse the slurry container adding the rinsate to the spray tank. Complete filling the spray tank to the desired level. The spray tank agitation should be sufficient to ensure uniform spray mixture during application and until the spray tank has been emptied. When tank mixing with other products, NC-603 WG should be mixed first in the spray tank. After the NC-603 WG is thoroughly mixed, add the other products as specified on their label. Ensure the compatibility of other products with NC-603 WG before mixing them together in the spray tank. Avoid the overnight storage of NC-603 WG spray mixtures. Premixing NC-603 WG solutions in nurse tanks is not recommended. Maintain continuous spray solution agitation until all the spray solution has been applied.

Do not use with tank additives that alter the pH of the spray solution below pH 5 or above pH 8. Buffer spray solution to alter the pH range if appropriate.

Spray Equipment Clean-Out

Many pesticides are very active at low rates, especially to sensitive crops. Residues left in mixing equipment, spray tanks, hoses, spray booms and nozzles can cause crop effects if they are not properly cleaned. As soon as possible after spraying NC-603 WG and before using the sprayer equipment for any other applications, the sprayer equipment must be thoroughly cleaned using the following procedure. In addition, users must take appropriate steps to ensure proper equipment clean-out for any other products mixed with NC-603 WG as required on the other product labels. More complete cleaning can be achieved if the spray system is cleaned immediately following the application.

1. Drain sprayer tank, hoses, spray boom and spray nozzles. Use a high pressure detergent wash to remove physical sediment and residues from the inside of the sprayer tank and thoroughly rinse. Then, thoroughly flush sprayer hoses, spray boom and spray nozzles with a clean water rinse. Remove and clean spray tips and all filters and screens (tank, spray hose and spray tips) separately in the ammonia solution of Step 2.

2. Next, prepare a sprayer cleaning solution by adding three gallons of ammonia (containing at least 2% active) per 100 gallons of clean water. Prepare sufficient cleaning solution to allow the operation of the spray system for a minimum of 15 minutes to thoroughly flush hoses, spray boom and spray nozzles.

4/6

3. Convenient and thorough cleaning of the sprayer can be achieved if the ammonia solution or fresh water is left in the spray tank, hoses, spray booms and spray nozzles overnight or during storage.

4. Before using the sprayer, completely drain the sprayer system. Rinse the tank with clean water and flush through the hoses, spray boom, and spray nozzles with clean water. Remove and clean spray tips and all filters and screens (tank, spray hose and spray tip) separately in an ammonia solution.

5. Properly dispose of all cleaning solution and rinsate in accordance with Federal, State, and local regulations and guidelines.

Do not apply sprayer cleaning solutions or rinsate to sensitive crops.

Do not store the sprayer overnight or for any extended period of time with NC-603 WG spray solution remaining in the tank, spray lines, spray boom plumbing, spray nozzles or strainers.

If the sprayer has been stored or idle, purge the spray boom and nozzles with clean water before beginning any application.

Should small quantities of NC-603 WG remain in inadequately cleaned mixing, loading and/or spray equipment, they may be released during subsequent applications potentially causing effects to certain crops and other vegetation. Nissan accepts no liability for any effects due to inadequately cleaned equipment.

APPLICATION INFORMATION

Ground Applications

Utilize a boom and nozzle sprayer equipped with the appropriate nozzles, spray tips and screens and adjusted to provide optimum spray distribution and coverage at appropriate operating pressures. Utilize nozzles which produce minimal amounts of fine spray droplets. Apply a minimum of 10 gallons of finished spray per acre. Higher spray volumes are required when there is a dense weed population or crop canopy. Spray booms should be adjusted to position spray tips at the minimum height of 18" above the crop and operated to avoid the application of excessive herbicide rates directly over the rows and/or into the whorl of the crop. Be aware that overlaps and slower ground speeds while starting, stopping or turning while spraying may result in higher application rates and possible crop response.

Aerial Applications

Use nozzles and arrangements that will provide optimum coverage while producing a minimal amount of fine droplets. Apply at a minimum of 3 gallons of finished spray per acre. Higher spray volumes are required when there is a dense weed population or crop canopy.

Spray Drift Management

Interactions of equipment and environmental factors influence the potential for spray drift. The applicator is responsible for considering all these factors when applying any pesticide. For ground boom applications, apply with a nozzle height of no more than 4 feet above the ground or crop canopy. Drift potential increases at wind speeds < 3 MPH or > 10 MPH. Many factors including droplet size and application equipment determine drift potential at any given wind speed. Use nozzles that are designed for the intended application. Agriculturally approved deposition and retention additives may be added to the spray tank to assist in the application process. Not all additives have been tested with NC-603 WG. Follow manufacturers' label and recommendations. AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

Application Restrictions

Do not apply this product through any type of irrigation system.

Following application to foliage, allow 30 days before grazing domestic livestock, harvesting forage, or harvesting silage.

Do not apply more than 2 applications per season in corn (field, seed, silage, sweet, and pop).

Updated 09-2-03

S:\Nissan\NC-603 WG\Revised Labels\Nissan Priority Label Revisions 2004 09022003.v2.doc

Do not apply more than 1 application per season in grain sorghum.

For all herbicide products used during the use season, the total pounds applied of carfentrazone and halosulfuron must not exceed the pounds a.i. per acre per season specified in the table below.

MAXIMUM ALLOWABLE NC-603 WG USE PER ACRE PER SEASON			
Total Allowed NC-603 WG Use*			
Crop	Ounces NC-603 WG	Pounds a.i. carfentrazone	Pounds a.i. halosulfuron
Field Corn	2.0	0.016	0.064
Seed Corn	2.0	0.016	0.064
Silage Corn	2.0	0.016	0.064
Sweet Corn	2.0	0.016	0.064
Popcorn	2.0	0.016	0.064
Grain Sorghum	1.0	0.008	0.032

*The total allowed usage volumes include all applications made to the field per calendar year. This includes fallow treatments, burndown treatments and all in-season treatments. The 2nd treatment of NC-603 WG may be applied only with drop nozzles semi-directed or directed.

CROP ROTATIONAL RESTRICTIONS

Labeled crops may be planted at specified time intervals following application of approved rates of NC-603 WG. Use the time intervals listed below to determine the required time interval before planting.

TIME INTERVAL BEFORE REPLANTING (Months after treatment with NC-603 WG)

Crop	Months
Field Corn	1
Barley (Winter)	2
Oats (Winter)	2
Rice	2
Seed corn	2
Sorghum	2
Spring cereal crops (wheat, oats, barley)	2
Wheat (winter)	2
Popcorn, Sweetcorn	3
Cotton	4
Soybeans	9
Alfalfa	12
Clovers	12
Cucumbers, Pumpkins, Squash	12
Dry beans	12
Eggplant	12
Field peas	12
Forage Grasses	12
Peanuts	12
Peas	12
Peppers	12
Potatoes	12
Proso Millet	12
Radish	12
Rye (winter)	12
Snap Beans	12
Spring Cereal crops (other)	12
Tomato (transplant)	12
Cabbage	15
Canola	15
Carrot	15
Mint	15
Broccoli, Cauliflower, Collards	18
Leeks, Onions	18
Lettuce crops	18
Sunflowers	18
Sugarbeet (Michigan only)	21
Sugarbeet and Red Beet	24
Spinach	24
Sugarbeet (ND, MN, Red River Valley)*	36

*Also includes other areas where rainfall is sparse or irrigation is required.

Refer to individual product labels to determine rotation crop restrictions when tank mixtures are used.

CORN

Field Corn, Seed Corn, Popcorn, Corn Silage, and Sweet Corn (Processing and Fresh Market)

Apply NC-603 WG alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply to corn in all tillage systems from crop emergence up to 8-leaf collar corn growth stage. Do not apply when wind speed and direction are likely to cause spray drift.

For best performance, make application to actively growing weeds up to 4 inches high and rosettes less than 3 inches across. **Coverage is essential for good control.**

To control weeds not listed on this label, NC-603 WG may be tank mixed with other herbicides registered for use in corn. When tank mixing NC-603 WG with other products, be sure the NC-603 WG is mixed in the spray tank water first. For specific mixing instructions, refer to the Mixing and Loading Instructions under the GENERAL INFORMATION section.

For Broadcast Applications

Field Corn and Silage Production

Use 1.0 oz NC-603 WG product per acre plus NIS at 0.25 % volume to volume (2 pints per 100 gallons) with or without UAN or AMS as described in the adjuvant use requirements section.

Apply NC-603 WG alone or as a tank mixture with other herbicides to emerged and actively growing weeds.

Sweet Corn / Popcorn Production

NC-603 WG may be applied to sweet corn, however, the user assumes all responsibility for herbicide tolerance with such use. All hybrids/varieties have not been tested for sensitivity to NC-603 WG nor does Nissan have access to all seed company or food processor data. Broadcast applications may result in spray being concentrated into the whorl of the plant, which will increase leaf response. To minimize application into the whorl of the plants, drop nozzles or other type directed sprayers must be used to direct the spray to the target weeds and away from the whorl of the plant.

Therefore, any crop response arising from the use of NC-603 WG on sweet corn is the responsibility of the user. Use NC-603 WG only under the recommendation of the seed company, food processor, or State Agricultural Extension Service.

If necessary, a sequential treatment of this product at 1 oz of product per acre may be applied only with drop nozzles directed to avoid application into the corn plant whorl.

For Directed Applications

Field Corn and Silage

NC-603 WG may be applied with drop nozzles or other sprayers capable of directing the spray to the target weeds and away from the whorl of the corn plant. NC-603 WG may be used at 1.0 ounce of formulated product per acre using drop nozzles. Use appropriate rates of adjuvants.

Sweet Corn, Popcorn Production

NC-603 WG may be applied with drop nozzles or other sprayers capable of directing the spray to the target weeds and away from the whorl of the corn plant. NC-603 WG may be used up to a maximum of 1.0 ounce of formulated product per acre using drop nozzles for control of larger weed sizes for those weeds listed below under "Control of Weeds". Use appropriate rates of adjuvants.

Seed Corn Production (Directed Applications Only)

For seed production fields, apply NC-603 WG at 1.0 ounce formulated product per acre using drop nozzles or other equipment to make a directed spray treatment. Avoid directing spray solution into the whorl.

Seed corn inbreds have generally shown good tolerance to NC-603 WG, however, all inbreds have not been tested. Use appropriate rate of adjuvants.

GRAIN SORGHUM

TIMING AND METHOD OF APPLICATION

Refer to General Information and Application Information sections and follow all restrictions.

Apply NC-603 WG alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply to sorghum in all tillage systems from the 2nd to 6th leaf growth stage. For best performance, make applications to actively growing weeds up to 4 inches tall and rosettes less than 3 inches across. Refer to Weeds Controlled with NC-603 WG Chart to determine if a tank mix partner is needed.

Use a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient with a minimum of 65% surfactant load. Postemergence broadcast applications of NC-603 WG herbicide with crop oil concentrate or methylated seed oils are not recommended as increased crop response may occur.

Coverage is essential for good control.

To control weeds not listed on this label, NC-603 WG may be tank mixed with other herbicides registered for use in grain sorghum. When tank mixing NC-603WG with other products, be sure the NC-603 WG is mixed in the spray tank water first. For specific mixing instructions, refer to the Mixing and Loading Instructions under the GENERAL INFORMATION section. Refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions, and rotational cropping restrictions. Sprayers should be adjusted and operated to avoid the application of excessive herbicide rates directly over the row and/or into the whorl of the sorghum plant. Broadcast applications of NC-603 WG to sorghum with wet foliage or application during periods of adverse environmental conditions such as cool, cloudy, wet, or high humidity may cause increased crop response.

Use Rates

Use NC-603 WG at 1.0 ounce per acre. Applications should be made by ground equipment using a minimum finished spray volume of 10 gallons of spray per acre or by air at a minimum finished spray volume of 3 gallons of spray per acre.

For Directed Applications

Grain Sorghum

Drop nozzles are recommended if applications are to be made under any of these conditions to limit the amount of product deposited into sorghum leaves and/or into the sorghum whorl.

When applying NC-603 WG postemergence to sorghum grown for seed, the use of drop nozzles is recommended. Do not apply NC-603 WG to sweet sorghum.

Weeds controlled with NC-603 WG

Use NC-603 WG at 1.0 ounce/acre (0.008 pound active ingredient carfentrazone + 0.032 pound active ingredient halosulfuron). Applications should be made by ground equipment using a minimum finished spray volume of 10 gallons of spray per acre using nonionic surfactant at 0.25% v/v (2 pt per 100 gallons of water) with or without UAN or AMS as described in the adjuvant use requirements section of this label.

When used as directed, NC-603 WG will provide control of listed weeds up to the indicated sizes:

	1 oz NC-603 WG per acre	
Buffalobur	3 inches	6 inches
Burcucumber	Suppression	6 inches
Cocklebur, common	5 inches	14 inches
Common purslane	Suppression	4 inches
Devilsclaw	2 inches	4 inches
Field bindweed	Suppression	4 inches
Hemp sesbania	6 inches	6 inches
Hophornbeam copperleaf	2 inches	4 inches

Jimsonweed	Suppression	4 inches
Kochia	3 inches	6 inches
Lambsquarters, common	3 inches	6 inches
Marestail	Suppression	4 inches
Milkweed, common	Suppression	3 inches
Milkweed, honeyvine	Suppression	6 inches
Morningglories	2-3 true leaves	2-3 true leaves
Nightshade, Eastern black	4 inches	6 inches
Nutsedge, purple	Suppression	Suppression
Nutsedge, yellow	Suppression	Suppression
Pigweed, other	Suppression	4 inches
Pigweed, redroot	4 inches	8 inches
Pigweed, triazine resistant	4 inches	8 inches
Pokeweed, common	6 inches	18 inches
Potato, volunteer	3 inches	6 inches
Prickly sida	Suppression	2 inches
Prostrate spurge	Suppression	Suppression
Ragweed, common	9 inches	12 inches
Ragweed, giant	3 inches	6 inches
Russian thistle	3 inches	5 inches
Smartweed, Pennsylvania	2 inches	6 inches
Spurred anoda	2 inches	6 inches
Sunflower, common	12 inches	15 inches
Velvetleaf	18 inches	18 inches
Venice mallow	3 inches	6 inches
Waterhemp, common	2 inches (with COC at 1% v/v)	6 inches
Waterhemp, tall	2 inches (with COC at 1% v/v)	6 inches
Wild buckwheat	3 inches	6 inches
Wild mustard	Suppression	4 inches

Tank Mixtures

NC-603 WG may be tank mixed with other herbicides, labeled for use on corn (field, pop, seed, silage and sweet) to control weeds not listed on this label provided that the tank mix product does not prohibit such mixing. Read the manufacturer's label directions, restrictions, and use precautions for the companion herbicide. Follow the most prohibitive restrictions and use precautions for all products used. When tank mixing NC-603 WG with other products, be sure NC-603 WG is mixed in the spray tank water first.

For control of additional broadleaf weeds and grasses in corn (field, pop, seed, silage, and sweet), NC-603 WG may be tank mixed with 2,4-D (amine), Accent®, Accent Gold®, atrazine, Banvel®, Basis®, Basis Gold®, Beacon®, Callisto™, Clarity™, Distinct®, Exceed®, Homet®, Liberty®, Lightning®, Marksman®, Northstar™, Permit®, Roundup®, or other glyphosate products, Roundup UltraMAX®, Sencor®, Spirit™, Steadfast®, Touchdown®, and Yukon™.

For control of additional broadleaf weeds in grain sorghum, NC-603 WG may be tank mixed with Clarity™, Marksman®, Banvel®, 2,4-D, atrazine, and other products unless specifically prohibited.

When tank mixing NC-603 WG with other products, use the adjuvants recommended on the tank mix partner label. These may include nonionic surfactant, crop oil concentrate, 28% nitrogen, ammonium sulfate or combinations of these.

Leaf speckling can occur when NC-603 WG is used with certain crop protection products and adjuvants. Refer to the Tank Mixtures and Recommended Adjuvants sections under General Information for details.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: Nissan makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product: Nissan's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Accent, Accent Gold, Basis, Basis Gold - trademarks of E.I. DuPont de Nemours and Company
Beacon, Exceed, Northstar, Spirit, Touchdown - trademarks of Syngenta Crop Protection, Inc.
Banvel, Clarity, Distinct, Marksman, Lightning - trademarks of BASF Corporation
Homet - trademark of Dow Agrosciences, LLC
Permit - trademark of Nissan Chemical Industries, Ltd.
Roundup, Roundup UltraMAX, Roundup WeatherMAX - trademarks of Monsanto Company
Sencor - trademark of Bayer CropScience

Nissan Chemical Industries, Ltd.
7-1, 3-chome, Kanda-Nishiki-cho
Chiyoda-ku, Tokyo 101-0054
Japan
NC-603 WG_3_3-12-2003b