

DIRECTIONS FOR USE ON LAWN OR ATHLETIC FIELD TURF. APPLY THE RECOMMENDED AMOUNT OF CHEMICAL TO AREA SPECIFIED.

Mark off the area to be treated and cross-spray, first from end to end then from side to side. With most knapsack and small tank sprayers, the recommended amount of water will cover the area twice. When spot spraying, avoid overdosage by making no more than 3 quick passes over each clump.

Soil should be moist at time of application. Areas treated in hot, dry weather should be thoroughly watered 3 to 4 days before treatment. Do not water for 3 to 4 days after treatment.

Lawns can be reseeded as soon as crabgrass is dead.

Newly seeded areas may be treated after they have become established and been cut at least twice. In fine-leaved fescue lawns, treat only in early spring or early fall. If more than the recommended rate is applied, most grasses including Zoysia will show temporary discoloration. Do not use on St. Augustine grass.

AMOUNTS TO USE

Seedling crabgrass, foxtail, newly seeded or bent grass lawns:

1 pt. in 8 gal. water to cover 2,000 sq. ft.

1 qt. in 15 gal. water to cover 4,000 sq. ft.

1 gal. in 60 gal. water to cover 16,000 sq. ft.

Repeated applications may be needed as more crabgrass seedlings germinate during the summer.

Branched crabgrass:

1 pt. in 4 gal. water to cover 1,000 sq. ft.

Two or more treatments at 5 to 7 day intervals may be needed for effective control.

Dallisgrass in bermudagrass turf (Southern States)

1 pt. in 3 gal. water to cover 800 sq. ft.

1 qt. in 6 gal. water to cover 1,600 sq. ft.

1 gal. in 24 gal. water to cover 6,400 sq. ft.

Spray from mid-spring to early summer. Mow area 1 to 3 days before spraying. Do not mow, fertilize or water for 48 hours after spraying. For best results, repeat application in one week.

AMCHEM PRODUCTS, INC. AMBLER, PA.

WAG

ACTIVE INGREDIENTS
O O diethyl O 2
phosphorothioate*
Aromatic petroleum d
INERT INGREDIENTS
E.C.

* DIAZINO

WAG
New Or

E.C.

F