directions under normal conditions of use but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extend, to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.

USDA Rel. 0. 577-39 RED

LD-323Vh \*\*

SCLD FOR HERBICIDE PURPOSEL RED

Ma in U.S.A. for ACME INSECTICIDE DIVISION Acme Quality Paints, Inc. Detroit, Michigan 48211

\*\*Insert month and year of current printing e.g.: 7-67

## \*DIRECTIONS FOR USE: Read ALL the Label First!

To be effective, existing weeds must be removed in early Spring or treatment made on freshly cultivated ground. Apply 1 pound per 100 square feet. The contents of this package treats 400 square feet of flower bed. Sift over ground as uniformly as possible. Water lightly after application. Do not treat dry, hard ground, do not cultivate ground after application. A second late Summer or Fall application gives year around weed free plantings. See cautions for plants that should not be treated.

Controls crabgrass, purslane, lambsquarters, red root pigweed, common chickweed, spotted spruge as well as other annual grasses and broad leaf weeds. Ragweed and smartweed are not controlled.

May be used around and under shade trees, evergreens, flowering shrubs and hedge rows. Also in groundcovers such as ivv. and pachysandra. Do not use on dichondra. Does not prevent growth into treated areas by turf grasses.

May be used very effectively in rose beds and in iris and gladiolus plantings.

Treat annual flowers after transplanting from flats or after thinning and initial weeding. Do not treat immediately after seeding petunias, snapdragons and similar flowers.

STRAWBERRIES: New planting — apply overall immediately after transplanting. Established Planting - apply in the Fall and early Spring before weeds emerge. Do not apply after first bloom. Use ½ pound per 60 square feet or entire contents on 480 sq. ft. of bed.

VEGETABLES: Treat snap beans as soon as planted. Use around onion sets and after transplanting cabbage, cauliflower, broccoli, sweet potato siips and tomatoes. On tomatoes, defer treatment 4 to 6 weeks after transplanting, clean cultivate before making application.

\*Cautions: Do not take internally. Avoid contact with eyes. Wash after handling or using. Do not contaminate feed or foodstuffs. Do not store near seeds, fertilizer or pesticides.

The following plants may be injured — Do Not use near these: Bugle weed, button pinks, carnations, dianthus, geum, ice plants (mesembryantheum). pansy, phlox, portulaca, sweet william, telanthera, violets. Use around tulips, amaryllis and hosta is questionable.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the sted on the label when used in accordance 4 POUNDS



**NET WEIGHT** 



Contains 21/2% Dacthal\*

Stops New Weed Growth in Flower Beds, around Trees, Shrubs, Evergreens, **Vegetable Transplants** and Strawberries



The following species of ornamentals, shrubs and flowers can be treated:

Abelia **Alyssum** Aster Arbor-vitae Babys-breath Ash Bellflower Azalea Bleeding-heart Barberry **Blood-leaf** Boxwood Bugloss Birch Candle Larkspur Camellia

Candytufts Chestnut Cinquefoil Cotoneaster Cottonwood Chrysanthemum Columbine Crab Apple Coleus Cypress Coreopsis Deutzia Coral bells Dogwood Cosmos Elaegnus Elm Cuphea Dahtia Euonymus Delphinum

**Evening primrose** Forsyth:a Gum Fernleaf yarrow Hawthorn Feaverfew Holly Forget-me-not Hydrangea Four-O'clock Japanese andromeda **Foxglove** Juniper Gaillardia

Lilac Gladiolus Locust Gold dust Magnolia Golden marguerite

Geranium

Maple Golden tuft Mock orange **Iris** 

Lantana

Mountain-Laurel Lavendercotton

Oak Lily **Pachistima** Lupine Pine Morning glory Pittosporum Marigold **Podocarous** Mother-of-Thyme **Poplar** 

Mourning bride Privet Mourning pinks Redbud Nasturtium Rhododendron Orgine **Peonies** Rose Petunia Russian olive Pink heath Spirea Spruce Poker plant Sycamore Purple coneflower

Tree Peony Salvia Tulip tree Scarlet sage Snapdragon Variegated privet

greens, flowering shrubs and hedge rows. Also in groundcovers such as ivv. and pachysandra. Do not use on dichondra. Does not prevent growth into treated areas by turf grasses.

May be used very effectively in rose beds and in iris and gladiolus plantings.

Treat annual flowers after transplanting from flats or after thinning and initial weeding. Do not treat immediately after seeding petunias, snapdragons and similar flowers.

STRAWBERRIES: New planting — apply overall immediately after transplanting. Established Planting — apply in the Fall and early Spring before weeds emerge. Do not apply after first bloom. Use ½ pound per 60 square feet or entire contents on 480 sq. ft. of bed.

VEGETABLES: Treat snap beans as soon as planted. Use around onion sets and after transplanting cabbage, cauliflower, broccoli, sweet potato slips and tomatoes. On tomatoes, defer treatment 4 to 6 weeks after transplanting, clean cultivate before making application.

\*Cautions: Do not take internally. Avoid contact with eyes. Wash after handling or using. Do not contaminate feed or foodstuffs. Do not store near seeds, fertilizer or pesticides.

The following plants may be injured — Do Not use near these: Bugle weed, button pinks, carnations, dianthus, geum, ice plants (mesembryantheum). pansy, phlox, portulaca, sweet william, telanthera, violets. Use around tulips, amaryllis and hosta is questionable.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Do not reuse container. Destroy when empty.

SOLD FOR HERBICIDE PURPOSES ACME INSECTICIDE DIVISION The Sherwin-Williams Co.

8250 St. Aubin Ave., t. troit Mich. 48211


USDA REGISTRATION NO. 577-474

ACCEPTED

CAUTION: KEEP OUT OF REACH OF CHILDREN SEE SIDE PANELS FOR OTHER CAUTONS

## Contains 21/2% Bacthal\*

## Stops New Weed Growth in Flower Beds, around Trees, Shrubs, Evergreens, **Vegetable Transplants** and Strawberries



## ACTIVE INGREDIENT 2.5% INERT INGREDIENTS ......... 97.5%

TOTAL 100.0%

\*Trade Mark Diamond Alkali Co \*\*U.S. Patent 2,923,634. For use as pre-emergent hurbicide Elaegnus Cosmos Cuphea Elm Dahlia Euonymus Delphinum Forsythia **Evening primrose** Guin Fernleaf yarrow Hawthorn Feaverfew Holly Forget-me-not Hydrangea Four-O'clock Japanese androineda Fexglove Juniper Gaillardia Lantana Geranium Lilac Gladiolus Gold dust Locust

Magnolia Golden marguerite Maple Golden tuft

Mock orange Iris

PORMOOR

Mountain-Laurel Lavendercotton

Oak Lily **Pachistima** Lupine Pine Morning giory **Pittosporum** Marigold

**Podocarous** Mother-of-Thyme Poplar Mourning bride Privet Mourning pinks Redbud Nasturtium Rhododendron Orpine **Peonies** Rose Petunia Russian olive Pink heath Spirea Poker plant Spruce Purple coneflower Sycamore

Tree Peony Salvia Scarlet sage Tulip tree Variegated privet Snapdragon Viburnum Stone Crop Strawflower Walnut Weigela Sundrops Willow Sunflower Yew Sweet Pea Baltic Ivy Verbena

Boston Ivy Virginia Spiderwort English Ivy **Wall Flowers** Honeysuckle Wormwood **Pachysandra** Zinnia

Wilson Ivy Ageratum

F. N. LEDE AND POURLING UP #1T

EL ANDER NO. .... SUITE T - LD-314F2 6有6 展初特 代码 Captainta II.