

(Front Panel)

ONE POUND


ACME  
BORDEAUX  
MIXTURE

NET WEIGHT

07-86

"A Fixed Copper Spray"

"For Disease Control on Peonies, Evergreens,  
Shrubs, Flowers, Fruit and Shade Trees"



1 1/4 Pt. ----- CAUTION: KEEP OUT OF REACH OF CHILDREN ----- 10 Pt.  
SEE LEFT PANEL FOR ADDITIONAL CAUTIONS ----- 6 Pt. Min.

Active Ingredient:

Copper, expressed as metallic (in the form  
of Bordeaux Mixture) .....

12.75%

Inert Ingredients: .....

87.25%

100.00%

(Right-Side Panel)

DIRECTIONS

(8-8-100 is Same as 4-4-50)

APRICOTS, PLUMS, SOUR CHERRIES: Brown rot and leaf spot, use 4 TB. per gallon plus equal amount of lime. Do not use on Japanese plums. Use at shuck split stage and in first and second cover sprays.

PEACH: Leaf curl, use 12 TB/gal. in early spring. Repeat late fall after leaf drop.

PECAN: Scab, use 6 TB/gal. when catkins show. Repeat 3 or 4 times at 3-week intervals. Spanish moss, use 12 TB/gal. Spray during winter.

GRAPES: Black rot and mildew, use 6 TB/gal. when shoots are 18 inches long. Repeat after bloom and 10 days later, again in mid-August.

POTATOES: Early and late blight, use 8 TB/gal. Apply when plants are 3 to 4 inches high, maintain a 10 to 14-day schedule for entire season.

STRAWBERRY: Leaf spot, use 6 TB/gal. at 10 to 14-day intervals, except during harvest. Final spray in late summer.

CITRUS: Lemon brown rot, grapefruit scab, melanose, use 6 TB/gal. Spray just before spring flush. Repeat 1 to 3 weeks after bloom.

TOMATO: Early and late blight, Septoria leaf spot, use 6 TB/gal. Repeat at 7 to 10-day intervals. Alternate with Zineb 75W if Anthracnose is a problem.

(continued)

(Right-Side Panel - cont'd)

MIXING TABLE

Use level spoonfuls in measuring. 3 teaspoonfuls equal 1 tablespoonful. Use higher dosage if can has been shaken to fluff the powder.

Amount of Spray Desired	Amount of Bordeaux Mixture Required	
	4-4-100	8-8-100
1 Pint	3/4 to 1 tablespoon	1 to 1 1/2 tablespoons
1 Quart	1 1/2 to 1-3/4 tablespoons	2 to 3 tablespoons
1 Gallon	6 to 7 tablespoons	8 to 12 tablespoons
3 Gallons	1-1/8 to 1 1/2 cupfuls	1 1/2 to 2 1/2 cupfuls

(Left-Side Panel)

Dry Bordeaux Mixture is made from equal parts of Copper Sulphate and Lime. Eight pounds of this product, mixed with 50 gallons of water, is equivalent to a 4-4-50 Bordeaux Mixture formula. This strength is sometimes referred to as 8-8-100.

Disease control is a matter of prevention. Application should be made beforehand or as soon as first signs of disease appear.

APPLES: To control late scab, black rot, bitter rot and blotch, use 8 TBS/gal. plus 8 TBS. of hydrated lime. May russet certain varieties.

PEARS: Fire blight, use 4 TBS/gal. in full bloom. Do not add insecticides.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

CAUTION: HARMFUL IF SWALLOWED

Do not reuse container

Destroy when empty

(Back Panel)

SYCAMORES, LINDEN, MAPLE, OAK, TULIP TREE, ELLI: Control of Anthracnose, Leaf Spot and Elm Leaf Curl, use 8 TBS/gal. as leaves uncurl in spring. Repeat in 2 to 3 weeks.

(continued)

(Back Panel--cont'd)

JUNIPER, CEDAR, CYPRESS, ARBORVITAE, SPRUCE, PINE, YEW: For Needle Cast, Sphaeropsis blight, twig blight, Coryneum blight, apply fall and spring using 8 TBS/gal. Spray Cedar and Junipers July-August for Cedar Apple Rust.

IVY: Leaf Spot, use 6 TBS/gal. in spring. Repeat in 14 days or at first sign of disease.

LAUREL, LILAC, RHODODENDRON: Leaf Spots and blight, use 4 TBS/gal. in early spring. Again after bloom.

DOGWOOD, BOXWOOD, CAMELLIAS: Dieback, Leaf Spot and Nectria Canker, use 8 TBS/gal. as growth starts in early spring.

DAHLIAS, TULIPS, GLADIOLUS, LILIES, AND IRIS: For Leaf Spot and Botrytis blight (gray mold), use 6 TBS/gal. in early spring. Repeat in 7 to 10 days.

PEONY: Leaf Blotch, Botrytis blight (gray mold), use 6 TBS/gal. Spray before shoots are one foot tall, repeat in two weeks. If any sign of bud blast, spray again. Remove any wilted shoots by cutting below ground. Place in paper bag and burn. Remove old stalks and leaves in fall and burn - do not compost.

DELPHINIUM, GERANIUM: Leaf Spots, use 4 TBS/gal. at first sign of disease.

PALMS: Anthracnose, Scab, Leaf Spot, use 6 TBS/gal. Spray at first sign of disease.

USDA Registration No. 577-97

LD-11,772 8-68

Made in U.S.A.

ACME INSECTICIDE DIVISION  
THE SHERWIN-WILLIAMS CO.  
8250 St. Aubin Ave., Detroit, Mich. 48211

Sold for Fungicide Purposes

Draft 3/27/69  
Typed 6/10/69