ACME®


37-86

GARDEN *

Contains 1% ROTENONE

CAUTION: Keep Out of Reach of Children
See Side Panels for Additional Caution Statements.

ACTIVE INGREDIENTS:
Rotenone 1.00%
Other Cube Resins 1.60%
INERT INGREDIENTS 97.40%



DIRECTIONS FOR USE

As a dust — apply with a dust gun or sift from the shaker top canister. Dust plant thoroughly, covering both upper and lower sides of foliage, covering all insects to be controlled. Make repeat applications weekly to cover new growth or until the pests are under control.

ACME GARDEN GUARDS, when applied as directed, may be used on edible portions of vegetables. Wash to remove residues before eating FOR CONTROL OF . . .

Asparagus: Asparagus beetles.

Beans: Mexican bean beetle.
bean leaf beetle.

Cabbage & Cauliflower: Cabbage looper, vegetable weevil, Harlequin bug.

Celery: To control celery leaf tier, dust at first sign of infestation in celery hearts and repeat within 24 hours.

Cucumbers: Striped cucumber beetle. Squash Vine borer.

Currants: Imported Currantworm.

Dust as soon as feeding is observed on leaves.

Eggplants & Peppers: Flea beetles, pepper maggot.

Lettuce: Garden webworm.

Potatoes: Colorado potato beetle.

Tomatoes: Flea beetles.

flea beetles.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and conditions of application equipment will vary, Seller makes no claims other than those stated on this label.



ACME[®]

37-86

GARDEN GRARD°

Contains 1% ROTENONE

CAUTION: Keep Out of Reach of Children

See Side Panels for Additional Caution Statements.

ACTIVE INGREDIENTS:

Rotenone 1.00%
Other Cube Resins 1.60%
INERT INGREDIENTS 97.40%
TOTAL 100.00%



FLOWERS: For control of aphids and exposed flower thrips on Asters, Delphiniums, Hollyhocks Roses Snapdragons, and Zinnias, dust lightly just as flower buds open Repeat once each week

DOGS & CATS: Effective against fleas and lice on dogs and cats. Do not use on pet birds or poultry. Repeat application when fleas or line reappear. Dust all body areas, especially back of head and neck, Brush into the fur Treat animal sleeping quarters and bedding.

CAUTION: Harmful if taken internally Do not breathe dust. May cause irritation of nose and throat. Do not apply to crops within one day of harvest.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DO NOT RE-USE CONTAMICER. DESTROY WHEN EMPTY.

EPA REG. NO. 33955-270-AA EPA EST. NO. 2217-KS-1 554-975

NET WEIGHT ONE POUND

ACCEPTED OCT 23 1975

Under the Federal Irse time, Fungicide, one is not permitted to a permitted recipied and a second to the permitted recipied and a second to the permitted to th